

Groupoid Models for Diagrams of Groupoid Correspondences

Joanna Ko

joint with Celso Antunes, and Ralf Meyer

Masarykova Univerzita

Higher structures in Noncommutative Geometry and
Quantum Algebra

Motivation

Motivation

Idea

- many C^* -algebras can be expressed as groupoid C^* -algebras

Motivation

Idea

- many C^* -algebras can be expressed as groupoid C^* -algebras
- e.g. C^* -algebras associated to group actions, higher-rank graphs, and self-similar groups

Motivation

Idea

- many C^* -algebras can be expressed as groupoid C^* -algebras
- e.g. C^* -algebras associated to group actions, higher-rank graphs, and self-similar groups

Albandik and Meyer has given an interpretation of higher-rank graphs as dynamical systems

Motivation

Idea

- many C^* -algebras can be expressed as groupoid C^* -algebras
- e.g. C^* -algebras associated to group actions, higher-rank graphs, and self-similar groups

Albandik and Meyer has given an interpretation of higher-rank graphs as dynamical systems

\rightsquigarrow by studying a general kind of dynamical system, to recover the constructions of these groupoids and their C^* -algebras

Groupoid correspondences

Groupoid correspondences

Muhly, Renault, Williams: Morita equivalences of groupoids

Groupoid correspondences

Muhly, Renault, Williams: Morita equivalences of groupoids

Hilsum, Skandalis: morphisms of groupoids

Groupoid correspondences

Muhly, Renault, Williams: Morita equivalences of groupoids

Hilsum, Skandalis: morphisms of groupoids

\rightsquigarrow C^* -correspondences **X**

Groupoid correspondences

Muhly, Renault, Williams: Morita equivalences of groupoids

Hilsum, Skandalis: morphisms of groupoids

\rightsquigarrow C^* -correspondences **X**

Idea

- groupoid correspondences are variants of these

Groupoid correspondences

Muhly, Renault, Williams: Morita equivalences of groupoids

Hilsum, Skandalis: morphisms of groupoids

\rightsquigarrow C^* -correspondences \times

Idea

- groupoid correspondences are variants of these
- they are spaces with commuting actions of two groupoids

Groupoid correspondences

Muhly, Renault, Williams: Morita equivalences of groupoids

Hilsum, Skandalis: morphisms of groupoids

\rightsquigarrow C^* -correspondences **X**

Idea

- groupoid correspondences are variants of these
- they are spaces with commuting actions of two groupoids
- taking groupoid C^* -algebras induces C^* -correspondences 😊

Groupoid correspondences

Definition

An *étale groupoid*:

Groupoid correspondences

Definition

An *étale groupoid*: range and source maps $r, s: \mathcal{G} \rightrightarrows \mathcal{G}^0$, multiplication, and inverse maps are local homeomorphisms

Groupoid correspondences

Definition

An *étale groupoid*: range and source maps $r, s: \mathcal{G} \rightrightarrows \mathcal{G}^0$, multiplication, and inverse maps are local homeomorphisms

An étale groupoid is *locally compact*: \mathcal{G}^0 is locally compact Hausdorff

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X}

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X} with a continuous *anchor map* $s: \mathcal{X} \rightarrow \mathcal{G}^0$,

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X} with a continuous *anchor map* $s: \mathcal{X} \rightarrow \mathcal{G}^0$, and a continuous map

$$\cdot: \mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X},$$

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X} with a continuous *anchor map* $s: \mathcal{X} \rightarrow \mathcal{G}^0$, and a continuous map

$$\cdot: \mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X},$$

such that

$$(1) \quad s(x \cdot g) = s(g) \text{ for } x \in \mathcal{X}, g \in \mathcal{G} \text{ with } s(x) = r(g)$$

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X} with a continuous *anchor map* $s: \mathcal{X} \rightarrow \mathcal{G}^0$, and a continuous map

$$\cdot: \mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X},$$

such that

- (1) $s(x \cdot g) = s(g)$ for $x \in \mathcal{X}$, $g \in \mathcal{G}$ with $s(x) = r(g)$
- (2) $(x \cdot g_1) \cdot g_2 = x \cdot (g_1 \cdot g_2)$ $g_1, g_2 \in \mathcal{G}$ with $s(x) = r(g_1)$,
 $s(g_1) = r(g_2)$

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X} with a continuous *anchor map* $s: \mathcal{X} \rightarrow \mathcal{G}^0$, and a continuous map

$$\cdot: \mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X},$$

such that

- (1) $s(x \cdot g) = s(g)$ for $x \in \mathcal{X}$, $g \in \mathcal{G}$ with $s(x) = r(g)$
- (2) $(x \cdot g_1) \cdot g_2 = x \cdot (g_1 \cdot g_2)$ $g_1, g_2 \in \mathcal{G}$ with $s(x) = r(g_1)$,
 $s(g_1) = r(g_2)$
- (3) $x \cdot s(x) = x$

Groupoid correspondences

Definition (Groupoid actions)

A *right \mathcal{G} -space*: \mathcal{X} with a continuous *anchor map* $s: \mathcal{X} \rightarrow \mathcal{G}^0$, and a continuous map

$$\cdot: \mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X},$$

such that

- (1) $s(x \cdot g) = s(g)$ for $x \in \mathcal{X}$, $g \in \mathcal{G}$ with $s(x) = r(g)$
- (2) $(x \cdot g_1) \cdot g_2 = x \cdot (g_1 \cdot g_2)$ $g_1, g_2 \in \mathcal{G}$ with $s(x) = r(g_1)$,
 $s(g_1) = r(g_2)$
- (3) $x \cdot s(x) = x$

Notation

Write $r: \mathcal{X} \rightarrow \mathcal{G}^0$ for left anchor maps.

Groupoid correspondences

Definition (Free, proper, and basic actions)

Groupoid correspondences

Definition (Free, proper, and basic actions)

A right \mathcal{G} -space is *free* if and only if the map

$$\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X} \times \mathcal{X}, \quad (x, g) \mapsto (x \cdot g, x).$$

is injective.

Groupoid correspondences

Definition (Free, proper, and basic actions)

A right \mathcal{G} -space is *free* if and only if the map

$$\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X} \times \mathcal{X}, \quad (x, g) \mapsto (x \cdot g, x).$$

is injective.

It is *proper* if and only if this map is proper.

Groupoid correspondences

Definition (Free, proper, and basic actions)

A right \mathcal{G} -space is *free* if and only if the map

$$\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{G} \rightarrow \mathcal{X} \times \mathcal{X}, \quad (x, g) \mapsto (x \cdot g, x).$$

is injective.

It is *proper* if and only if this map is proper.

It is *basic* if and only if this map is a homeomorphism onto its image.

Groupoid correspondences

Relationship between basic and free and proper actions:

Groupoid correspondences

Relationship between basic and free and proper actions:

Proposition

Let \mathcal{X} be a right \mathcal{G} -space.

Groupoid correspondences

Relationship between basic and free and proper actions:

Proposition

Let \mathcal{X} be a right \mathcal{G} -space. TFAE:

- (1) the action of \mathcal{G} on \mathcal{X} is basic and the orbit space \mathcal{X}/\mathcal{G} is Hausdorff;*
- (2) the action of \mathcal{G} on \mathcal{X} is free and proper.*

Groupoid correspondences

Definition (Groupoid correspondence)

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X}

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X} with commuting actions of \mathcal{H} on the left and \mathcal{G} on the right:

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X} with commuting actions of \mathcal{H} on the left and \mathcal{G} on the right:

for $s(h) = r(x)$ and $s(x) = r(g)$

$$(1) \quad s(h \cdot x) = s(x)$$

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X} with commuting actions of \mathcal{H} on the left and \mathcal{G} on the right:

for $s(h) = r(x)$ and $s(x) = r(g)$

$$(1) \quad s(h \cdot x) = s(x)$$

$$(2) \quad r(x \cdot g) = r(x)$$

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X} with commuting actions of \mathcal{H} on the left and \mathcal{G} on the right:

for $s(h) = r(x)$ and $s(x) = r(g)$

$$(1) \quad s(h \cdot x) = s(x)$$

$$(2) \quad r(x \cdot g) = r(x)$$

$$(3) \quad (h \cdot x) \cdot g = h \cdot (x \cdot g)$$

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X} with commuting actions of \mathcal{H} on the left and \mathcal{G} on the right:

for $s(h) = r(x)$ and $s(x) = r(g)$

$$(1) \quad s(h \cdot x) = s(x)$$

$$(2) \quad r(x \cdot g) = r(x)$$

$$(3) \quad (h \cdot x) \cdot g = h \cdot (x \cdot g)$$

such that $s: \mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism and the right \mathcal{G} -action is *basic*.

Groupoid correspondences

Definition (Groupoid correspondence)

A *groupoid correspondence* $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is a space \mathcal{X} with commuting actions of \mathcal{H} on the left and \mathcal{G} on the right:

for $s(h) = r(x)$ and $s(x) = r(g)$

$$(1) \quad s(h \cdot x) = s(x)$$

$$(2) \quad r(x \cdot g) = r(x)$$

$$(3) \quad (h \cdot x) \cdot g = h \cdot (x \cdot g)$$

such that $s: \mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism and the right \mathcal{G} -action is *basic*.

When \mathcal{G} and \mathcal{H} are locally compact, and \mathcal{X}/\mathcal{G} is Hausdorff, then it is a *locally compact groupoid correspondence*.

Groupoid correspondences

Remark

In the case of a locally compact groupoid correspondence, \mathcal{X} is automatically locally compact.

Groupoid correspondences

Remark

In the case of a locally compact groupoid correspondence, \mathcal{X} is automatically locally compact.

Definition (Proper and tight correspondences)

Groupoid correspondences

Remark

In the case of a locally compact groupoid correspondence, \mathcal{X} is automatically locally compact.

Definition (Proper and tight correspondences)

A groupoid correspondence $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is *proper*:

$$r_*: \mathcal{X}/\mathcal{G} \rightarrow \mathcal{H}^0$$

induced by $r: \mathcal{X} \rightarrow \mathcal{H}^0$ is proper.

Groupoid correspondences

Remark

In the case of a locally compact groupoid correspondence, \mathcal{X} is automatically locally compact.

Definition (Proper and tight correspondences)

A groupoid correspondence $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is *proper*:

$$r_*: \mathcal{X}/\mathcal{G} \rightarrow \mathcal{H}^0$$

induced by $r: \mathcal{X} \rightarrow \mathcal{H}^0$ is proper.

It is *tight*: r_* is a homeomorphism.

Examples of groupoid correspondences

Examples of groupoid correspondences

Example (Topological correspondences)

Examples of groupoid correspondences

Example (Topological correspondences)

Let \mathcal{G} and \mathcal{H} be locally compact spaces.

Examples of groupoid correspondences

Example (Topological correspondences)

Let \mathcal{G} and \mathcal{H} be locally compact spaces.

Then the groupoid actions on \mathcal{X} are simply continuous maps on \mathcal{X} .

Examples of groupoid correspondences

Example (Topological correspondences)

Let \mathcal{G} and \mathcal{H} be locally compact spaces.

Then the groupoid actions on \mathcal{X} are simply continuous maps on \mathcal{X} .

By the proposition, \mathcal{X} is locally compact Hausdorff.

Examples of groupoid correspondences

Example (Topological correspondences)

Let \mathcal{G} and \mathcal{H} be locally compact spaces.

Then the groupoid actions on \mathcal{X} are simply continuous maps on \mathcal{X} .

By the proposition, \mathcal{X} is locally compact Hausdorff.

A groupoid correspondence $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is the same as a locally compact Hausdorff \mathcal{X} with a continuous map $r: \mathcal{X} \rightarrow \mathcal{H}$ and a local homeomorphism $s: \mathcal{X} \rightarrow \mathcal{G}$.

Examples of groupoid correspondences

Example (Topological correspondences)

Let \mathcal{G} and \mathcal{H} be locally compact spaces.

Then the groupoid actions on \mathcal{X} are simply continuous maps on \mathcal{X} .

By the proposition, \mathcal{X} is locally compact Hausdorff.

A groupoid correspondence $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is the same as a locally compact Hausdorff \mathcal{X} with a continuous map $r: \mathcal{X} \rightarrow \mathcal{H}$ and a local homeomorphism $s: \mathcal{X} \rightarrow \mathcal{G}$.

\rightsquigarrow a topological correspondence

Examples of groupoid correspondences

Example (Topological correspondences)

Let \mathcal{G} and \mathcal{H} be locally compact spaces.

Then the groupoid actions on \mathcal{X} are simply continuous maps on \mathcal{X} .

By the proposition, \mathcal{X} is locally compact Hausdorff.

A groupoid correspondence $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is the same as a locally compact Hausdorff \mathcal{X} with a continuous map $r: \mathcal{X} \rightarrow \mathcal{H}$ and a local homeomorphism $s: \mathcal{X} \rightarrow \mathcal{G}$.

\rightsquigarrow a topological correspondence

If furthermore $\mathcal{H} = \mathcal{G}$, it is a *topological graph* as in Katsura's work.

Examples of groupoid correspondences

Example ('Group correspondences')

Let \mathcal{G} and \mathcal{H} be discrete groups.

Examples of groupoid correspondences

Example ('Group correspondences')

Let \mathcal{G} and \mathcal{H} be discrete groups.

Since $s: \mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is discrete.

Examples of groupoid correspondences

Example ('Group correspondences')

Let \mathcal{G} and \mathcal{H} be discrete groups.

Since $s: \mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is discrete.

In this case, the action is basic \iff free.

Examples of groupoid correspondences

Example ('Group correspondences')

Let \mathcal{G} and \mathcal{H} be discrete groups.

Since $s: \mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is discrete.

In this case, the action is basic \iff free.

A groupoid correspondence $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ is the same as a set \mathcal{X} with commuting actions, where the right \mathcal{G} -action is free.

Examples of groupoid correspondences

Example ('Group correspondences')

Examples of groupoid correspondences

Example ('Group correspondences')

Write $A := \mathcal{X}/\mathcal{G}$.

Examples of groupoid correspondences

Example ('Group correspondences')

Write $A := \mathcal{X}/\mathcal{G}$. Since the right \mathcal{G} -action is free, we have

$$A \times \mathcal{G} \cong \mathcal{X}$$

Examples of groupoid correspondences

Example ('Group correspondences')

Write $A := \mathcal{X}/\mathcal{G}$. Since the right \mathcal{G} -action is free, we have

$$A \times \mathcal{G} \cong \mathcal{X}$$

So $A \times \mathcal{G} \cong \mathcal{X}$ is a groupoid correspondence $\mathcal{H} \leftarrow \mathcal{G}$ isomorphic to \mathcal{X} .

Examples of groupoid correspondences

Example ('Group correspondences')

Write $A := \mathcal{X}/\mathcal{G}$. Since the right \mathcal{G} -action is free, we have

$$A \times \mathcal{G} \cong \mathcal{X}$$

So $A \times \mathcal{G} \cong \mathcal{X}$ is a groupoid correspondence $\mathcal{H} \leftarrow \mathcal{G}$ isomorphic to \mathcal{X} .

Left \mathcal{H} -action on $A \times \mathcal{G}$:

$$h \cdot (x, g) = (\pi_h(x), \varphi(h, x) \cdot g)$$

for $\pi_h: A \rightarrow A$, $\varphi: \mathcal{H} \times A \rightarrow \mathcal{G}$

Examples of groupoid correspondences

Example ('Group correspondences')

Left \mathcal{H} -action on $A \times \mathcal{G}$:

$$h \cdot (x, g) = (\pi_h(x), \varphi(h, x) \cdot g)$$

Examples of groupoid correspondences

Example ('Group correspondences')

Left \mathcal{H} -action on $A \times \mathcal{G}$:

$$h \cdot (x, g) = (\pi_h(x), \varphi(h, x) \cdot g)$$

(1) $h \mapsto \pi_h$ is a group action on A

Examples of groupoid correspondences

Example ('Group correspondences')

Left \mathcal{H} -action on $A \times \mathcal{G}$:

$$h \cdot (x, g) = (\pi_h(x), \varphi(h, x) \cdot g)$$

- (1) $h \mapsto \pi_h$ is a group action on A
- (2) $\varphi: \mathcal{H} \times A \rightarrow \mathcal{G}$ is a 1-cocycle

Examples of groupoid correspondences

Example ('Group correspondences')

Left \mathcal{H} -action on $A \times \mathcal{G}$:

$$h \cdot (x, g) = (\pi_h(x), \varphi(h, x) \cdot g)$$

(1) $h \mapsto \pi_h$ is a group action on A

(2) $\varphi: \mathcal{H} \times A \rightarrow \mathcal{G}$ is a 1-cocycle

$$\varphi(h_1 h_2, x) = \varphi(h_1, h_2 x) \cdot \varphi(h_2, x)$$

Examples of groupoid correspondences

Example ('Group correspondences')

Left \mathcal{H} -action on $A \times \mathcal{G}$:

$$h \cdot (x, g) = (\pi_h(x), \varphi(h, x) \cdot g)$$

(1) $h \mapsto \pi_h$ is a group action on A

(2) $\varphi: \mathcal{H} \times A \rightarrow \mathcal{G}$ is a 1-cocycle

$$\varphi(h_1 h_2, x) = \varphi(h_1, h_2 x) \cdot \varphi(h_2, x)$$

Furthermore, any isomorphism $\mathcal{X} \cong A \times \mathcal{G}$ is unique up to

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

$$h \cdot (x, g) \mapsto h \cdot (x, \psi(x) \cdot g) = (\pi_h(x), \varphi(h, x) \cdot \psi(x) \cdot g)$$

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

$$h \cdot (x, g) \mapsto h \cdot (x, \psi(x) \cdot g) = (\pi_h(x), \varphi(h, x) \cdot \psi(x) \cdot g)$$

\parallel

$$(\pi_h(x), \varphi(h, x) \cdot g)$$

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

$$h \cdot (x, g) \mapsto h \cdot (x, \psi(x) \cdot g) = (\pi_h(x), \varphi(h, x) \cdot \psi(x) \cdot g)$$

||

$$(\pi_h(x), \varphi(h, x) \cdot g) \mapsto (\pi_h(x), \psi(\pi_h(x)) \cdot \varphi^\psi(h, x) \cdot g)$$

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

$$h \cdot (x, g) \mapsto h \cdot (x, \psi(x) \cdot g) = (\pi_h(x), \varphi(h, x) \cdot \psi(x) \cdot g)$$

||

$$(\pi_h(x), \varphi(h, x) \cdot g) \mapsto (\pi_h(x), \psi(\pi_h(x)) \cdot \varphi^\psi(h, x) \cdot g)$$

$$\varphi^\psi(h, x) := \psi(\pi_h(x))^{-1} \cdot \varphi(h, x) \cdot \psi(x)$$

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

$$h \cdot (x, g) \mapsto h \cdot (x, \psi(x) \cdot g) = (\pi_h(x), \varphi(h, x) \cdot \psi(x) \cdot g)$$

\parallel

$$(\pi_h(x), \varphi(h, x) \cdot g) \mapsto (\pi_h(x), \psi(\pi_h(x)) \cdot \varphi^\psi(h, x) \cdot g)$$

The assignment of 'new' 1-cocycle

$$\psi \mapsto \varphi^\psi$$

is a right action.

Examples of groupoid correspondences

Example ('Group correspondences')

$$(x, g) \xrightarrow{k} (x, \psi(x) \cdot g)$$

$$h \cdot (x, g) \mapsto h \cdot (x, \psi(x) \cdot g) = (\pi_h(x), \varphi(h, x) \cdot \psi(x) \cdot g)$$

\parallel

$$(\pi_h(x), \varphi(h, x) \cdot g) \mapsto (\pi_h(x), \psi(\pi_h(x)) \cdot \varphi^\psi(h, x) \cdot g)$$

The assignment of 'new' 1-cocycle

$$\psi \mapsto \varphi^\psi$$

is a right action.

isomorphism classes of $\mathcal{H} \leftarrow \mathcal{G} \rightleftarrows$ equivalence classes of (A, φ)

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

$\mathcal{H} \leftarrow \mathcal{G}$ is proper $\iff A \cong \mathcal{X}/\mathcal{G}$ is finite,

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

$\mathcal{H} \leftarrow \mathcal{G}$ is proper $\iff A \cong \mathcal{X}/\mathcal{G}$ is finite,

It is tight $\iff A$ is a singleton.

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

$\mathcal{H} \leftarrow \mathcal{G}$ is proper $\iff A \cong \mathcal{X}/\mathcal{G}$ is finite,

It is tight $\iff A$ is a singleton.

Tight:

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

$\mathcal{H} \leftarrow \mathcal{G}$ is proper $\iff A \cong \mathcal{X}/\mathcal{G}$ is finite,

It is tight $\iff A$ is a singleton.

Tight:

$\mathcal{X} \cong \mathcal{G}$ with a base point in \mathcal{X} .

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

$\mathcal{H} \leftarrow \mathcal{G}$ is proper $\iff A \cong \mathcal{X}/\mathcal{G}$ is finite,

It is tight $\iff A$ is a singleton.

Tight:

$\mathcal{X} \cong \mathcal{G}$ with a base point in \mathcal{X} .

Left action of \mathcal{H} :

$$h \cdot g = \varphi(h) \cdot g$$

for a group homomorphism $\varphi: \mathcal{H} \rightarrow \mathcal{G}$.

Examples of groupoid correspondences

Example ('Group correspondences')

Since \mathcal{H}^0 is a singleton,

$\mathcal{H} \leftarrow \mathcal{G}$ is proper $\iff A \cong \mathcal{X}/\mathcal{G}$ is finite,

It is tight $\iff A$ is a singleton.

Tight:

$\mathcal{X} \cong \mathcal{G}$ with a base point in \mathcal{X} .

Left action of \mathcal{H} :

$$h \cdot g = \varphi(h) \cdot g$$

for a group homomorphism $\varphi: \mathcal{H} \rightarrow \mathcal{G}$.

If we pick another base point, φ is replaced by

$$\mathrm{Ad}_g^{-1} \circ \varphi$$

Examples of groupoid correspondences

Example ('Group correspondences')

Tight:

isomorphism classes of tight $\mathcal{H} \leftarrow \mathcal{G} \rightleftarrows$ equivalence classes of

$$\varphi: \mathcal{H} \rightarrow \mathcal{G}$$

Examples of groupoid correspondences

Example ('Group correspondences')

Tight:

isomorphism classes of tight $\mathcal{H} \leftarrow \mathcal{G} \rightleftarrows$ equivalence classes of

$$\varphi: \mathcal{H} \rightarrow \mathcal{G}$$

If $\varphi: \mathcal{H} \rightarrow \mathcal{G}$ is injective, and $\mathcal{G} = \mathcal{H}$, we can get Stammer's notion.

Examples of groupoid correspondences

Example ('Group correspondences')

Tight:

isomorphism classes of tight $\mathcal{H} \leftarrow \mathcal{G} \rightleftarrows$ equivalence classes of

$$\varphi: \mathcal{H} \rightarrow \mathcal{G}$$

If $\varphi: \mathcal{H} \rightarrow \mathcal{G}$ is injective, and $\mathcal{G} = \mathcal{H}$, we can get Stammeier's notion.

If φ is injective, and both \mathcal{G} and \mathcal{H} are Abelian, we recover Cuntz and Vershik's notion.

Examples of groupoid correspondences

Example ('Group correspondences')

Proper:

Examples of groupoid correspondences

Example ('Group correspondences')

Proper:

isomorphism classes of proper $\mathcal{H} \leftarrow \mathcal{G} \rightleftarrows$ Nekrashevych's covering
permutational bimodules

Examples of groupoid correspondences

Example ('Group correspondences')

Proper:

isomorphism classes of proper $\mathcal{H} \leftarrow \mathcal{G} \rightsquigarrow$ Nekrashevych's covering
permutational bimodules

where covering permutational bimodules describe self-similarities of
groups.

Examples of groupoid correspondences

Example ('Group correspondences')

Proper:

isomorphism classes of proper $\mathcal{H} \leftarrow \mathcal{G} \rightsquigarrow$ Nekrashevych's covering
permutational bimodules

where covering permutational bimodules describe self-similarities of
groups.

\rightsquigarrow a proper $\mathcal{G} \leftarrow \mathcal{G}$ can be viewed as a self-similarity of \mathcal{G}

The bicategory \mathfrak{Gr}_{1c}

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$.

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$.

Let \mathcal{G} act on $\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{Y}$ by the diagonal action

$$g \cdot (x, y) := (x \cdot g^{-1}, g^{-1} \cdot y)$$

with $s(g) = r(y) = s(x)$.

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$.

Let \mathcal{G} act on $\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{Y}$ by the diagonal action

$$g \cdot (x, y) := (x \cdot g^{-1}, g^{-1} \cdot y)$$

with $s(g) = r(y) = s(x)$.

Denote by

$$\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$$

the orbit space,

The bicategory \mathfrak{Gr}_{lc}

Construction (Composition of groupoid correspondences)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$.

Let \mathcal{G} act on $\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{Y}$ by the diagonal action

$$g \cdot (x, y) := (x \cdot g^{-1}, g^{-1} \cdot y)$$

with $s(g) = r(y) = s(x)$.

Denote by

$$\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$$

the orbit space, and by $[x, y]$ the image of (x, y) .

The bicategory \mathfrak{Gr}_{lc}

Construction (Composition of groupoid correspondences)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$.

Let \mathcal{G} act on $\mathcal{X} \times_{s, \mathcal{G}^0, r} \mathcal{Y}$ by the diagonal action

$$g \cdot (x, y) := (x \cdot g^{-1}, g^{-1} \cdot y)$$

with $s(g) = r(y) = s(x)$.

Denote by

$$\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$$

the orbit space, and by $[x, y]$ the image of (x, y) .

Set $r(x, y) := r(x)$ and $s(x, y) := s(y)$.

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

We set

$$h \cdot [x, y] := [h \cdot x, y] \qquad [x, y] \cdot k := [x, y \cdot k]$$

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

We set

$$h \cdot [x, y] := [h \cdot x, y] \quad [x, y] \cdot k := [x, y \cdot k]$$

We have:

Proposition (Albandik, Antunes, K., Meyer)

The actions of \mathcal{H} and \mathcal{K} on $\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$ are well-defined and turn $\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$ into a groupoid correspondence

$$\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}: \mathcal{H} \leftarrow \mathcal{K}$$

The bicategory \mathfrak{Gr}_{1c}

Construction (Composition of groupoid correspondences)

We set

$$h \cdot [x, y] := [h \cdot x, y] \quad [x, y] \cdot k := [x, y \cdot k]$$

We have:

Proposition (Albandik, Antunes, K., Meyer)

The actions of \mathcal{H} and \mathcal{K} on $\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$ are well-defined and turn $\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$ into a groupoid correspondence

$$\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}: \mathcal{H} \leftarrow \mathcal{K}$$

If both \mathcal{X} and \mathcal{Y} are proper or tight, then so is $\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}$.

The bicategory \mathfrak{Gr}_{1c}

Definition (Equivariant maps)

The bicategory \mathfrak{Gr}_{1c}

Definition (Equivariant maps)

A continuous map $f: \mathcal{X} \rightarrow \mathcal{Y}$ between two right \mathcal{G} -spaces is

\mathcal{G} -equivariant:

$$s(f(x)) = s(x)$$

for all x ,

The bicategory \mathfrak{Gr}_{lc}

Definition (Equivariant maps)

A continuous map $f: \mathcal{X} \rightarrow \mathcal{Y}$ between two right \mathcal{G} -spaces is

\mathcal{G} -equivariant:

$$s(f(x)) = s(x)$$

for all x , and

$$f(x \cdot g) = f(x) \cdot g$$

for all x and g with $s(x) = r(g)$.

The bicategory \mathfrak{Gr}_{lc}

Proposition (Albandik, Antunes, K., Meyer)

The following data constitute a bicategory \mathfrak{Gr}_{lc} :

The bicategory \mathfrak{Gr}_{lc}

Proposition (Albandik, Antunes, K., Meyer)

The following data constitute a bicategory \mathfrak{Gr}_{lc} :

- *objects: **locally compact** étale groupoids $\mathcal{H}, \mathcal{G}, \dots$*

The bicategory \mathfrak{Gr}_{lc}

Proposition (Albandik, Antunes, K., Meyer)

The following data constitute a bicategory \mathfrak{Gr}_{lc} :

- *objects: **locally compact** étale groupoids $\mathcal{H}, \mathcal{G}, \dots$*
- *1-morphisms: **locally compact** groupoid correspondences $\mathcal{X}, \mathcal{Y}: \mathcal{H} \rightleftarrows \mathcal{G}$*

The bicategory \mathfrak{Gr}_{lc}

Proposition (Albandik, Antunes, K., Meyer)

The following data constitute a bicategory \mathfrak{Gr}_{lc} :

- *objects: **locally compact** étale groupoids $\mathcal{H}, \mathcal{G}, \dots$*
- *1-morphisms: **locally compact** groupoid correspondences $\mathcal{X}, \mathcal{Y}: \mathcal{H} \rightleftarrows \mathcal{G}$*
- *2-morphisms: injective, \mathcal{H}, \mathcal{G} -equivariant continuous $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$*

The bicategory \mathfrak{Gr}_{lc}

Proposition (Albandik, Antunes, K., Meyer)

The following data constitute a bicategory \mathfrak{Gr}_{lc} :

- objects: *locally compact* étale groupoids $\mathcal{H}, \mathcal{G}, \dots$
- 1-morphisms: *locally compact* groupoid correspondences $\mathcal{X}, \mathcal{Y}: \mathcal{H} \rightleftarrows \mathcal{G}$
- 2-morphisms: injective, \mathcal{H}, \mathcal{G} -equivariant continuous $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$

with unitors given by

$$\begin{aligned}\mathcal{H} \circ_{\mathcal{H}} \mathcal{X} &\rightarrow \mathcal{X} & \mathcal{X} \circ_{\mathcal{G}} \mathcal{G} &\rightarrow \mathcal{X} \\ [h, x] &\mapsto h \cdot x & [x, g] &\mapsto x \cdot g\end{aligned}$$

The bicategory \mathfrak{Gr}_{lc}

Proposition (Albandik, Antunes, K., Meyer)

The following data constitute a bicategory \mathfrak{Gr}_{lc} :

- objects: *locally compact* étale groupoids $\mathcal{H}, \mathcal{G}, \dots$
- 1-morphisms: *locally compact* groupoid correspondences $\mathcal{X}, \mathcal{Y}: \mathcal{H} \rightleftarrows \mathcal{G}$
- 2-morphisms: injective, \mathcal{H}, \mathcal{G} -equivariant continuous $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$

with unitors given by

$$\begin{aligned}\mathcal{H} \circ_{\mathcal{H}} \mathcal{X} &\rightarrow \mathcal{X} & \mathcal{X} \circ_{\mathcal{G}} \mathcal{G} &\rightarrow \mathcal{X} \\ [h, x] &\mapsto h \cdot x & [x, g] &\mapsto x \cdot g\end{aligned}$$

and associator given by

$$\begin{aligned}\mathcal{X}_1 \circ_{\mathcal{G}_1} (\mathcal{X}_2 \circ_{\mathcal{G}_2} \mathcal{X}_3) &\rightarrow (\mathcal{X}_1 \circ_{\mathcal{G}_1} \mathcal{X}_2) \circ_{\mathcal{G}_2} \mathcal{X}_3 \\ [x_1, [x_2, x_3]] &\mapsto [[x_1, x_2], x_3]\end{aligned}$$

C^* -correspondences

C^* -correspondences

Definition

A C^* -*correspondence* $A \leftarrow B$ is a Hilbert right B -module \mathcal{E} ,

C^* -correspondences

Definition

A C^* -*correspondence* $A \leftarrow B$ is a Hilbert right B -module \mathcal{E} , equipped with a non-degenerate left action of A ,

C^* -correspondences

Definition

A C^* -*correspondence* $A \leftarrow B$ is a Hilbert right B -module \mathcal{E} , equipped with a non-degenerate left action of A , i.e. a $*$ -homomorphism

$$\varphi: A \rightarrow \mathcal{L}(\mathcal{E})$$

Constructing C^* -correspondences

Constructing C^* -correspondences

Construction

Denote by $\mathfrak{S}(\mathcal{G})$ the \mathbb{C} -vector space of quasi-continuous functions on \mathcal{G} .

Constructing C^* -correspondences

Construction

Denote by $\mathfrak{S}(\mathcal{G})$ the \mathbb{C} -vector space of quasi-continuous functions on \mathcal{G} .

Equip $\mathfrak{S}(\mathcal{G})$ with a $*$ -algebra structure:

$$\begin{aligned}\xi * \eta(g) &= \sum_{\substack{h \\ r(h)=r(g)}} \xi(h)\eta(h^{-1}g) \\ \xi^*(g) &= \overline{\xi(g^{-1})}\end{aligned}$$

Constructing C^* -correspondences

Construction

Denote by $\mathfrak{S}(\mathcal{G})$ the \mathbb{C} -vector space of quasi-continuous functions on \mathcal{G} .

Equip $\mathfrak{S}(\mathcal{G})$ with a $*$ -algebra structure:

$$\begin{aligned}\xi * \eta(g) &= \sum_{\substack{h \\ r(h)=r(g)}} \xi(h)\eta(h^{-1}g) \\ \xi^*(g) &= \overline{\xi(g^{-1})}\end{aligned}$$

It can be shown that $\mathfrak{S}(\mathcal{G})$ has the largest C^* -norm.

Constructing C^* -correspondences

Construction

Denote by $\mathfrak{S}(\mathcal{G})$ the \mathbb{C} -vector space of quasi-continuous functions on \mathcal{G} .

Equip $\mathfrak{S}(\mathcal{G})$ with a $*$ -algebra structure:

$$\begin{aligned}\xi * \eta(g) &= \sum_{\substack{h \\ r(h)=r(g)}} \xi(h)\eta(h^{-1}g) \\ \xi^*(g) &= \overline{\xi(g^{-1})}\end{aligned}$$

It can be shown that $\mathfrak{S}(\mathcal{G})$ has the largest C^* -norm.

Definition ($C^*(\mathcal{G})$)

The *groupoid C^* -algebra* $C^*(\mathcal{G})$ of \mathcal{G} is the completion of $\mathfrak{S}(\mathcal{G})$ in the largest C^* -norm.

Constructing C^* -correspondences

Construction

Consider $\mathfrak{S}(\mathcal{X})$ for $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$.

Constructing C^* -correspondences

Construction

Consider $\mathfrak{S}(\mathcal{X})$ for $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$.

The left and right actions on $\mathfrak{S}(\mathcal{X})$ can be defined just like the convolution.

Constructing C^* -correspondences

Construction

Consider $\mathfrak{S}(\mathcal{X})$ for $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$.

The left and right actions on $\mathfrak{S}(\mathcal{X})$ can be defined just like the convolution.

Equip $\mathfrak{S}(\mathcal{X})$ with the $(\mathfrak{S}(\mathcal{G}))$ -valued inner product

$$\langle \xi \mid \eta \rangle(g) := \sum_{\substack{x \\ s(x)=r(g)}} \overline{\xi(x)} \eta(x \cdot g)$$

Constructing C^* -correspondences

Construction

Consider $\mathfrak{S}(\mathcal{X})$ for $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$.

The left and right actions on $\mathfrak{S}(\mathcal{X})$ can be defined just like the convolution.

Equip $\mathfrak{S}(\mathcal{X})$ with the $(\mathfrak{S}(\mathcal{G}))$ -valued inner product

$$\langle \xi \mid \eta \rangle(g) := \sum_{\substack{x \\ s(x)=r(g)}} \overline{\xi(x)} \eta(x \cdot g)$$

Define the norm on $\mathfrak{S}(\mathcal{X})$ as

$$\|\xi\| := \|\langle \xi \mid \xi \rangle\|_{C^*(\mathcal{G})}^{\frac{1}{2}}$$

Constructing C^* -correspondences

Construction

Consider $\mathfrak{S}(\mathcal{X})$ for $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$.

The left and right actions on $\mathfrak{S}(\mathcal{X})$ can be defined just like the convolution.

Equip $\mathfrak{S}(\mathcal{X})$ with the $(\mathfrak{S}(\mathcal{G}))$ -valued inner product

$$\langle \xi \mid \eta \rangle(g) := \sum_{\substack{x \\ s(x)=r(g)}} \overline{\xi(x)} \eta(x \cdot g)$$

Define the norm on $\mathfrak{S}(\mathcal{X})$ as

$$\|\xi\| := \|\langle \xi \mid \xi \rangle\|_{C^*(\mathcal{G})}^{\frac{1}{2}}$$

Notation $(C^*(\mathcal{X}))$

Denote by $C^*(\mathcal{X})$ the completion of $\mathfrak{S}(\mathcal{X})$ in this norm.

Cuntz-Pimsner algebras of C^* -correspondences

Cuntz-Pimsner algebras of C^* -correspondences

For any C^* -correspondence $\mathcal{E}: A \leftarrow A$, we can associate a C^* -algebra, called the *Cuntz-Pimsner algebra* $\mathcal{O}_{\mathcal{E}}$:

Cuntz-Pimsner algebras of C^* -correspondences

For any C^* -correspondence $\mathcal{E}: A \leftarrow A$, we can associate a C^* -algebra, called the *Cuntz-Pimsner algebra* $\mathcal{O}_{\mathcal{E}}$:

Idea

Consider covariant Toeplitz representations $(A, \mathcal{E}) \rightarrow B$:

Cuntz-Pimsner algebras of C^* -correspondences

For any C^* -correspondence $\mathcal{E}: A \leftarrow A$, we can associate a C^* -algebra, called the *Cuntz-Pimsner algebra* $\mathcal{O}_{\mathcal{E}}$:

Idea

Consider covariant Toeplitz representations $(A, \mathcal{E}) \rightarrow B$:
a $*$ -homomorphism $A \rightarrow B$ and a linear map $\tau: \mathcal{E} \rightarrow B$ satisfying some conditions.

Cuntz-Pimsner algebras of C^* -correspondences

For any C^* -correspondence $\mathcal{E}: A \leftarrow A$, we can associate a C^* -algebra, called the *Cuntz-Pimsner algebra* $\mathcal{O}_{\mathcal{E}}$:

Idea

Consider covariant Toeplitz representations $(A, \mathcal{E}) \rightarrow B$:
a $*$ -homomorphism $A \rightarrow B$ and a linear map $\tau: \mathcal{E} \rightarrow B$ satisfying some conditions.

The universal one, $(A, \mathcal{E}) \rightarrow \mathcal{O}_{\mathcal{E}}$, gives the Cuntz-Pimsner algebra.

Examples of $C^*(\mathcal{X})$

Examples of $C^*(\mathcal{X})$

Example (Topological graph C^* -algebras)

Examples of $C^*(\mathcal{X})$

Example (Topological graph C^* -algebras)

Let \mathcal{G} be a locally compact space.

Examples of $C^*(\mathcal{X})$

Example (Topological graph C^* -algebras)

Let \mathcal{G} be a locally compact space.

Recall that a groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ in this case is a topological graph.

Examples of $C^*(\mathcal{X})$

Example (Topological graph C^* -algebras)

Let \mathcal{G} be a locally compact space.

Recall that a groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ in this case is a topological graph.

We obtain $C^*(\mathcal{G}) = C_0(\mathcal{G})$.

Examples of $C^*(\mathcal{X})$

Example (Topological graph C^* -algebras)

Let \mathcal{G} be a locally compact space.

Recall that a groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ in this case is a topological graph.

We obtain $C^*(\mathcal{G}) = C_0(\mathcal{G})$.

As \mathcal{X} is Hausdorff, we have $\mathfrak{S}(\mathcal{X}) = C_c(\mathcal{X})$.

Examples of $C^*(\mathcal{X})$

Example (Topological graph C^* -algebras)

Let \mathcal{G} be a locally compact space.

Recall that a groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ in this case is a topological graph.

We obtain $C^*(\mathcal{G}) = C_0(\mathcal{G})$.

As \mathcal{X} is Hausdorff, we have $\mathfrak{S}(\mathcal{X}) = C_c(\mathcal{X})$.

The norm completion $C^*(\mathcal{X})$ is the C^* -correspondence in Katsura's work used to define topological graph C^* -algebras.

Examples of $C^*(\mathcal{X})$

Examples of $C^*(\mathcal{X})$

Example (Cuntz-Pimsner algebras of self-similar groups)

Examples of $C^*(\mathcal{X})$

Example (Cuntz-Pimsner algebras of self-similar groups)

Recall that a proper groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ for a group \mathcal{G} is the covering permutational bimodule of a self-similar group.

Examples of $C^*(\mathcal{X})$

Example (Cuntz-Pimsner algebras of self-similar groups)

Recall that a proper groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ for a group \mathcal{G} is the covering permutational bimodule of a self-similar group.

Here, $\mathfrak{S}(\mathcal{G})$ is the group ring $\mathbb{C}[\mathcal{G}]$

Examples of $C^*(\mathcal{X})$

Example (Cuntz-Pimsner algebras of self-similar groups)

Recall that a proper groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ for a group \mathcal{G} is the covering permutational bimodule of a self-similar group.

Here, $\mathfrak{S}(\mathcal{G})$ is the group ring $\mathbb{C}[\mathcal{G}]$ and $\mathfrak{S}(\mathcal{X})$ is the vector space $\mathbb{C}[\mathcal{X}]$ with basis \mathcal{X} .

Examples of $C^*(\mathcal{X})$

Example (Cuntz-Pimsner algebras of self-similar groups)

Recall that a proper groupoid correspondence $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$ for a group \mathcal{G} is the covering permutational bimodule of a self-similar group.

Here, $\mathfrak{S}(\mathcal{G})$ is the group ring $\mathbb{C}[\mathcal{G}]$ and $\mathfrak{S}(\mathcal{X})$ is the vector space $\mathbb{C}[\mathcal{X}]$ with basis \mathcal{X} .

And the Cuntz-Pimsner algebra $\mathcal{O}_{C^*(\mathcal{X})}$ is Nekrashevych's universal Cuntz-Pimsner algebra $\mathcal{O}_{(\mathcal{G}, \mathcal{X})}$ of the self-similar group.

The bicategory \mathbf{Corr}

The bicategory $\mathcal{C}orr$

The bicategory $\mathcal{C}orr$ of C^* -correspondences consists of

The bicategory \mathfrak{Corr}

The bicategory \mathfrak{Corr} of C^* -correspondences consists of

- objects: C^* -algebras A, B, \dots

The bicategory \mathfrak{Corr}

The bicategory \mathfrak{Corr} of C^* -correspondences consists of

- objects: C^* -algebras A, B, \dots
- 1-morphisms: C^* -correspondences $\mathcal{E}, \mathcal{F}: A \rightleftarrows B$

The bicategory \mathfrak{Corr}

The bicategory \mathfrak{Corr} of C^* -correspondences consists of

- objects: C^* -algebras A, B, \dots
- 1-morphisms: C^* -correspondences $\mathcal{E}, \mathcal{F}: A \ltimes B$
- 2-morphisms: isometric A, B -bimodule maps $\gamma: \mathcal{E} \rightarrow \mathcal{F}$

The bicategory \mathfrak{Corr}

The bicategory \mathfrak{Corr} of C^* -correspondences consists of

- objects: C^* -algebras A, B, \dots
- 1-morphisms: C^* -correspondences $\mathcal{E}, \mathcal{F}: A \ltimes B$
- 2-morphisms: isometric A, B -bimodule maps $\gamma: \mathcal{E} \rightarrow \mathcal{F}$
- composition: completed tensor product

Constructing $\mathcal{G}r_{lc} \rightarrow \mathcal{C}orr$

Constructing $\mathcal{G}r_{lc} \rightarrow \mathcal{C}orr$

Construction

Recall that a 2-morphism in \mathcal{G} is an injective \mathcal{H}, \mathcal{G} -equivariant continuous map $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$.

Constructing $\mathcal{G}r_{lc} \rightarrow \mathcal{C}orr$

Construction

Recall that a 2-morphism in \mathcal{G} is an injective \mathcal{H}, \mathcal{G} -equivariant continuous map $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$.

It can be shown that α is a homeomorphism onto an open subset of \mathcal{Y} ,

Constructing $\mathcal{G}r_{lc} \rightarrow \mathcal{C}orr$

Construction

Recall that a 2-morphism in \mathcal{G} is an injective \mathcal{H} , \mathcal{G} -equivariant continuous map $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$.

It can be shown that α is a homeomorphism onto an open subset of \mathcal{Y} , and therefore extension by zero gives an injective map

$$\tilde{\alpha}: \mathfrak{S}(\mathcal{X}) \rightarrow \mathfrak{S}(\mathcal{Y})$$

Constructing $\mathcal{G}r_{lc} \rightarrow \mathcal{C}orr$

Construction

Recall that a 2-morphism in \mathcal{G} is an injective \mathcal{H}, \mathcal{G} -equivariant continuous map $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$.

It can be shown that α is a homeomorphism onto an open subset of \mathcal{Y} , and therefore extension by zero gives an injective map

$$\tilde{\alpha}: \mathfrak{S}(\mathcal{X}) \rightarrow \mathfrak{S}(\mathcal{Y})$$

which preserves the bimodule structure and the inner product, and is an isometry.

Constructing $\mathfrak{Gr}_{lc} \rightarrow \mathfrak{Corr}$

Construction

Recall that a 2-morphism in \mathcal{G} is an injective \mathcal{H}, \mathcal{G} -equivariant continuous map $\mathcal{X} \xrightarrow{\alpha} \mathcal{Y}$.

It can be shown that α is a homeomorphism onto an open subset of \mathcal{Y} , and therefore extension by zero gives an injective map

$$\tilde{\alpha}: \mathfrak{S}(\mathcal{X}) \rightarrow \mathfrak{S}(\mathcal{Y})$$

which preserves the bimodule structure and the inner product, and is an isometry.

\rightsquigarrow extends uniquely to an isometric $C^*(\mathcal{H}), C^*(\mathcal{G})$ -bimodule map

$$C^*(\alpha): C^*(\mathcal{X}) \rightarrow C^*(\mathcal{Y})$$

Constructing $\mathcal{G}r_{lc} \rightarrow \mathcal{C}orr$

The functoriality constraint is guaranteed by

Constructing $\mathsf{Gr}_{\text{lc}} \rightarrow \mathsf{Corr}$

The functoriality constraint is guaranteed by

Proposition (Albandik, Antunes, K., Meyer)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$.

Constructing $\mathsf{Gr}_{lc} \rightarrow \mathsf{Corr}$

The functoriality constraint is guaranteed by

Proposition (Albandik, Antunes, K., Meyer)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$. There is

$$\mu_{\mathcal{X}, \mathcal{Y}}^0: \mathfrak{S}(\mathcal{X}) \otimes \mathfrak{S}(\mathcal{Y}) \rightarrow \mathfrak{S}(\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y})$$

$$\mu_{\mathcal{X}, \mathcal{Y}}^0(f_1 \otimes f_2)([x, y]) = \sum_{\substack{g \\ r(g)=s(x)}} f_1(xg) \cdot f_2(g^{-1}y)$$

Constructing $\mathsf{Gr}_{\mathrm{lc}} \rightarrow \mathsf{Corr}$

The functoriality constraint is guaranteed by

Proposition (Albandik, Antunes, K., Meyer)

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$ and $\mathcal{Y}: \mathcal{G} \leftarrow \mathcal{K}$. There is

$$\begin{aligned}\mu_{\mathcal{X},\mathcal{Y}}^0: \mathfrak{S}(\mathcal{X}) \otimes \mathfrak{S}(\mathcal{Y}) &\rightarrow \mathfrak{S}(\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y}) \\ \mu_{\mathcal{X},\mathcal{Y}}^0(f_1 \otimes f_2)([x, y]) &= \sum_{\substack{g \\ r(g)=s(x)}} f_1(xg) \cdot f_2(g^{-1}y)\end{aligned}$$

which extends to an isomorphism of $C^(\mathcal{K}), C^*(\mathcal{G})$ -correspondences*

$$\mu_{\mathcal{X},\mathcal{Y}}: C^*(\mathcal{X}) \otimes_{C^*(\mathcal{G})} C^*(\mathcal{Y}) \rightarrow C^*(\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y})$$

Constructing $\mathsf{Gr}_{\text{lc}} \rightarrow \mathsf{Corr}$

Proposition (Albandik, Antunes, K., Meyer)

The assignments

$$\mathcal{G} \mapsto C^*(\mathcal{G})$$

$$\mathcal{X} \mapsto C^*(\mathcal{X})$$

$$\alpha \mapsto C^*(\alpha)$$

Constructing $\mathfrak{Gr}_{lc} \rightarrow \mathfrak{Corr}$

Proposition (Albandik, Antunes, K., Meyer)

The assignments

$$\mathcal{G} \mapsto C^*(\mathcal{G})$$

$$\mathcal{X} \mapsto C^*(\mathcal{X})$$

$$\alpha \mapsto C^*(\alpha)$$

together with

$$C^*(1_{\mathcal{G}}) = 1_{C^*(\mathcal{G})}$$

$$\mu_{\mathcal{X}, \mathcal{Y}}: C^*(\mathcal{X}) \otimes_{C^*(\mathcal{G})} C^*(\mathcal{Y}) \rightarrow C^*(\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y})$$

Constructing $\mathfrak{Gr}_{lc} \rightarrow \mathfrak{Corr}$

Proposition (Albandik, Antunes, K., Meyer)

The assignments

$$\mathcal{G} \mapsto C^*(\mathcal{G})$$

$$\mathcal{X} \mapsto C^*(\mathcal{X})$$

$$\alpha \mapsto C^*(\alpha)$$

together with

$$C^*(1_{\mathcal{G}}) = 1_{C^*(\mathcal{G})}$$

$$\mu_{\mathcal{X}, \mathcal{Y}}: C^*(\mathcal{X}) \otimes_{C^*(\mathcal{G})} C^*(\mathcal{Y}) \rightarrow C^*(\mathcal{X} \circ_{\mathcal{G}} \mathcal{Y})$$

form a (strictly unital) pseudofunctor $\mathfrak{Gr}_{lc} \rightarrow \mathfrak{Corr}$.

Actions of C^* -correspondences

Actions of C^* -correspondences

An action of a C^* -correspondence is equivalent to an action of $(\mathbb{N}_0, +)$.

Actions of C^* -correspondences

An action of a C^* -correspondence is equivalent to an action of $(\mathbb{N}_0, +)$.

What about a general monoid M other than $(\mathbb{N}_0, +)$?

Actions of C^* -correspondences

An action of a C^* -correspondence is equivalent to an action of $(\mathbb{N}_0, +)$.

What about a general monoid M other than $(\mathbb{N}_0, +)$?

\rightsquigarrow assembling several C^* -correspondences in a certain way

Actions of C^* -correspondences

An action of a C^* -correspondence is equivalent to an action of $(\mathbb{N}_0, +)$.

What about a general monoid M other than $(\mathbb{N}_0, +)$?

\rightsquigarrow assembling several C^* -correspondences in a certain way

Fowler's *product systems (of C^* -correspondences) over monoids*

Actions of C^* -correspondences

An action of a C^* -correspondence is equivalent to an action of $(\mathbb{N}_0, +)$.

What about a general monoid M other than $(\mathbb{N}_0, +)$?

\rightsquigarrow assembling several C^* -correspondences in a certain way

Fowler's *product systems (of C^* -correspondences) over monoids*

Indeed,

$$M \rightarrow \mathfrak{Gr}_{lc} \xrightarrow{\text{red}} \mathfrak{Gr} \rightarrow \mathfrak{Corr}$$

is equivalent to a product system over M .

Actions of C^* -correspondences

An action of a C^* -correspondence is equivalent to an action of $(\mathbb{N}_0, +)$.

What about a general monoid M other than $(\mathbb{N}_0, +)$?

\rightsquigarrow assembling several C^* -correspondences in a certain way

Fowler's *product systems (of C^* -correspondences) over monoids*

Indeed,

$$M \rightarrow \mathfrak{Gr}_{lc} \xrightarrow{\text{red}} \mathfrak{Gr} \rightarrow \mathfrak{Corr}$$

is equivalent to a product system over M .

\rightsquigarrow we should study diagrams of groupoid correspondences

Diagrams of groupoid correspondences

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in $\mathfrak{G}\mathfrak{r}$, i.e., $\mathcal{C} \rightarrow \mathfrak{G}\mathfrak{r}$, consists of

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in \mathfrak{Gr} , i.e., $\mathcal{C} \rightarrow \mathfrak{Gr}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in \mathfrak{Gr} , i.e., $\mathcal{C} \rightarrow \mathfrak{Gr}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$
- (2) groupoid correspondences $\mathcal{X}: \mathcal{G}_x \leftarrow \mathcal{G}_y$ for each $g: x \leftarrow y$

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in \mathfrak{Gr} , i.e., $\mathcal{C} \rightarrow \mathfrak{Gr}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$
- (2) groupoid correspondences $\mathcal{X}: \mathcal{G}_x \leftarrow \mathcal{G}_y$ for each $g: x \leftarrow y$
- (3) isomorphisms $\mu_{g,h}: \mathcal{X}_g \circ_{\mathcal{G}_y} \mathcal{X}_h \xrightarrow{\cong} \mathcal{X}_{gh}$ for composable g, h

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in \mathfrak{Gr} , i.e., $\mathcal{C} \rightarrow \mathfrak{Gr}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$
 - (2) groupoid correspondences $\mathcal{X}: \mathcal{G}_x \leftarrow \mathcal{G}_y$ for each $g: x \leftarrow y$
 - (3) isomorphisms $\mu_{g,h}: \mathcal{X}_g \circ_{\mathcal{G}_y} \mathcal{X}_h \xrightarrow{\cong} \mathcal{X}_{gh}$ for composable g, h
- satisfying

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in \mathfrak{Gr} , i.e., $\mathcal{C} \rightarrow \mathfrak{Gr}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$
 - (2) groupoid correspondences $\mathcal{X}: \mathcal{G}_x \leftarrow \mathcal{G}_y$ for each $g: x \leftarrow y$
 - (3) isomorphisms $\mu_{g,h}: \mathcal{X}_g \circ_{\mathcal{G}_y} \mathcal{X}_h \xrightarrow{\cong} \mathcal{X}_{gh}$ for composable g, h
- satisfying

- \mathcal{X}_x for $x \in \text{ob } \mathcal{C}$ is the identity on \mathcal{G}_x

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in $\mathbb{G}\mathbf{r}$, i.e., $\mathcal{C} \rightarrow \mathbb{G}\mathbf{r}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$
 - (2) groupoid correspondences $\mathcal{X}: \mathcal{G}_x \leftarrow \mathcal{G}_y$ for each $g: x \leftarrow y$
 - (3) isomorphisms $\mu_{g,h}: \mathcal{X}_g \circ_{\mathcal{G}_y} \mathcal{X}_h \xrightarrow{\cong} \mathcal{X}_{gh}$ for composable g, h
- satisfying

- \mathcal{X}_x for $x \in \text{ob } \mathcal{C}$ is the identity on \mathcal{G}_x
- compatibility conditions concerning the unitor and the associator

Diagrams of groupoid correspondences

Definition

A \mathcal{C} -shaped diagram in \mathfrak{Gr} , i.e., $\mathcal{C} \rightarrow \mathfrak{Gr}$, consists of

- (1) groupoids \mathcal{G}_x for each $x \in \text{ob } \mathcal{C}$
 - (2) groupoid correspondences $\mathcal{X}: \mathcal{G}_x \leftarrow \mathcal{G}_y$ for each $g: x \leftarrow y$
 - (3) isomorphisms $\mu_{g,h}: \mathcal{X}_g \circ_{\mathcal{G}_y} \mathcal{X}_h \xrightarrow{\cong} \mathcal{X}_{gh}$ for composable g, h
- satisfying

- \mathcal{X}_x for $x \in \text{ob } \mathcal{C}$ is the identity on \mathcal{G}_x
- compatibility conditions concerning the unitor and the associator

Notation

We can describe a \mathcal{C} -shaped diagram by $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$.

Actions of diagrams of groupoid correspondences

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space.

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

- (1) a left \mathcal{G} -action on Y with anchor $r_Y: Y \rightarrow \mathcal{G}^0$

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

- (1) a left \mathcal{G} -action on Y with anchor $r_Y: Y \rightarrow \mathcal{G}^0$
- (2) an open, continuous, surjective map

$$\begin{aligned}\alpha: \mathcal{X} \times_{s, \mathcal{G}^0, r_Y} Y &\rightarrow Y \\ (\gamma, y) &\mapsto \gamma \cdot y\end{aligned}$$

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

- (1) a left \mathcal{G} -action on Y with anchor $r_Y: Y \rightarrow \mathcal{G}^0$
- (2) an open, continuous, surjective map

$$\begin{aligned}\alpha: \mathcal{X} \times_{s, \mathcal{G}^0, r_Y} Y &\rightarrow Y \\ (\gamma, y) &\mapsto \gamma \cdot y\end{aligned}$$

satisfying

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

- (1) a left \mathcal{G} -action on Y with anchor $r_Y: Y \rightarrow \mathcal{G}^0$
- (2) an open, continuous, surjective map

$$\begin{aligned}\alpha: \mathcal{X} \times_{s, \mathcal{G}^0, r_Y} Y &\rightarrow Y \\ (\gamma, y) &\mapsto \gamma \cdot y\end{aligned}$$

satisfying

- $x \cdot (g \cdot y) = (x \cdot g) \cdot y$

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

- (1) a left \mathcal{G} -action on Y with anchor $r_Y: Y \rightarrow \mathcal{G}^0$
- (2) an open, continuous, surjective map

$$\begin{aligned}\alpha: \mathcal{X} \times_{s, \mathcal{G}^0, r_Y} Y &\rightarrow Y \\ (\gamma, y) &\mapsto \gamma \cdot y\end{aligned}$$

satisfying

- $x \cdot (g \cdot y) = (x \cdot g) \cdot y$
- $r(x \cdot y) = r(x)$ and $g \cdot (x \cdot y) = (g \cdot x) \cdot y$

Actions of diagrams of groupoid correspondences

Definition

Let $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$, and Y be a space. An *action of \mathcal{X} on Y* consists of

- (1) a left \mathcal{G} -action on Y with anchor $r_Y: Y \rightarrow \mathcal{G}^0$
- (2) an open, continuous, surjective map

$$\alpha: \mathcal{X} \times_{s, \mathcal{G}^0, r_Y} Y \rightarrow Y$$
$$(\gamma, y) \mapsto \gamma \cdot y$$

satisfying

- $x \cdot (g \cdot y) = (x \cdot g) \cdot y$
- $r(x \cdot y) = r(x)$ and $g \cdot (x \cdot y) = (g \cdot x) \cdot y$
- if $\gamma \cdot y = \gamma' \cdot y'$, then there is $\eta \in \mathcal{G}$ so that

$$\gamma' = \gamma \cdot \eta \quad \text{and} \quad y = \eta \cdot y'$$

Actions of diagrams of groupoid correspondences

Actions of diagrams of groupoid correspondences

Note that the disjoint union $\mathcal{H} \sqcup \mathcal{G}$ is also an étale groupoid.

Actions of diagrams of groupoid correspondences

Note that the disjoint union $\mathcal{H} \sqcup \mathcal{G}$ is also an étale groupoid.

Definition

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$.

Actions of diagrams of groupoid correspondences

Note that the disjoint union $\mathcal{H} \sqcup \mathcal{G}$ is also an étale groupoid.

Definition

Let $\mathcal{X}: \mathcal{H} \leftarrow \mathcal{G}$. An action of \mathcal{X} on Y is defined as an action of

$$\mathcal{H} \sqcup \mathcal{G} \leftarrow \mathcal{H} \sqcup \mathcal{G}$$

on Y .

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Definition (F -actions)

Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$ of groupoid correspondences.

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Definition (F -actions)

Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$ of groupoid correspondences. An *F -action on Y* consists of

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Definition (F -actions)

Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$ of groupoid correspondences. An *F -action on Y* consists of

- (1) a partition $Y = \bigsqcup_{x \in \text{ob } \mathcal{C}} Y_x$

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Definition (F -actions)

Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$ of groupoid correspondences. An *F -action on Y* consists of

- (1) a partition $Y = \bigsqcup_{x \in \text{ob } \mathcal{C}} Y_x$
- (2) continuous $r_{Y_x}: Y_x \rightarrow \mathcal{G}_x^0$

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Definition (F -actions)

Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$ of groupoid correspondences. An **F -action on Y** consists of

- (1) a partition $Y = \bigsqcup_{x \in \text{ob } \mathcal{C}} Y_x$
- (2) continuous $r_{Y_x}: Y_x \rightarrow \mathcal{G}_x^0$
- (3) open, continuous, surjective maps

$$\alpha_g: \mathcal{X}_g \times_{s, \mathcal{G}_x^0, r} Y_x \rightarrow Y_{x'}$$

for $g: x' \leftarrow x$

Actions of diagrams of groupoid correspondences

We are ready to define actions of general diagrams of groupoid correspondences.

Definition (F -actions)

Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram $(\mathcal{G}_x, \mathcal{X}_g, \mu_{g,h})$ of groupoid correspondences. An F -action on Y consists of

- (1) a partition $Y = \bigsqcup_{x \in \text{ob } \mathcal{C}} Y_x$
- (2) continuous $r_{Y_x}: Y_x \rightarrow \mathcal{G}_x^0$
- (3) open, continuous, surjective maps

$$\alpha_g: \mathcal{X}_g \times_{s, \mathcal{G}_x^0, r} Y_x \rightarrow Y_{x'}$$

$$\text{for } g: x' \leftarrow x$$

satisfying evident conditions.

F -actions and groupoid models

F -actions and groupoid models

Definition

A continuous map $\varphi: Y \rightarrow Y'$ between two spaces with F -actions is *F -equivariant*:

F -actions and groupoid models

Definition

A continuous map $\varphi: Y \rightarrow Y'$ between two spaces with F -actions is *F -equivariant*:

$$\varphi(Y_x) \subseteq Y'_x$$

for all $x \in \mathcal{C}$,

F -actions and groupoid models

Definition

A continuous map $\varphi: Y \rightarrow Y'$ between two spaces with F -actions is *F -equivariant*:

$$\varphi(Y_x) \subseteq Y'_x$$

for all $x \in \mathcal{C}$, and

$$r(\varphi(y)) = r(y) \quad \text{and} \quad \varphi(\gamma \cdot y) = \gamma \cdot \varphi(y)$$

for all $g \in \text{mor } \mathcal{C}$, $y \in Y_{s(g)}$, $\gamma \in \mathcal{X}_g$ with $s(\gamma) = r(y)$.

F -actions and groupoid models

F -actions and groupoid models

Definition

An F -action Ω is *universal*:

F -actions and groupoid models

Definition

An F -action Ω is *universal*: for any space Y with an F -action, there is a unique F -equivariant map $Y \rightarrow \Omega$.

F -actions and groupoid models

Definition

An F -action Ω is *universal*: for any space Y with an F -action, there is a unique F -equivariant map $Y \rightarrow \Omega$.

Remark

There is a category of F -actions and F -equivariant maps.

F -actions and groupoid models

Definition

An F -action Ω is *universal*: for any space Y with an F -action, there is a unique F -equivariant map $Y \rightarrow \Omega$.

Remark

There is a category of F -actions and F -equivariant maps.

A universal F -action is the terminal object in this category.

F -actions and groupoid models

Definition

An F -action Ω is *universal*: for any space Y with an F -action, there is a unique F -equivariant map $Y \rightarrow \Omega$.

Remark

There is a category of F -actions and F -equivariant maps.

A universal F -action is the terminal object in this category.

Definition (Groupoid models)

A *groupoid model* for F -actions is an étale groupoid \mathcal{U} with natural bijections

$$\{\mathcal{U}\text{-actions on } Y\} \leftrightarrow \{F\text{-actions on } Y\}$$

for all spaces Y .

F -actions and groupoid models

In other words, groupoid models encode diagrams of groupoid correspondences:

action by a diagram of
groupoid correspondences \leftrightarrow action by some groupoid

F -actions and groupoid models

In other words, groupoid models encode diagrams of groupoid correspondences:

action by a diagram of
groupoid correspondences \leftrightarrow action by some groupoid

Indeed, in favourable cases, the Cuntz-Pimsner algebra associated to a diagram of groupoid correspondences is the groupoid C^* -algebra of the groupoid model.

F -actions and groupoid models

In other words, groupoid models encode diagrams of groupoid correspondences:

action by a diagram of
groupoid correspondences \leftrightarrow action by some groupoid

Indeed, in favourable cases, the Cuntz-Pimsner algebra associated to a diagram of groupoid correspondences is the groupoid C^* -algebra of the groupoid model.

\rightsquigarrow for a groupoid C^* -algebra to be defined, the groupoid has to be *locally compact*.

Existence of groupoid models

Questions:

Existence of groupoid models

Questions:

- (1) Does groupoid model always exists for any diagram of groupoid correspondences?

Existence of groupoid models

Questions:

- (1) Does groupoid model always exists for any diagram of groupoid correspondences?
- (2) Is the resulting groupoid model locally compact?

Existence of groupoid models

Questions:

- (1) Does groupoid model always exists for any diagram of groupoid correspondences?
- (2) Is the resulting groupoid model locally compact? Under what *extra* conditions?

Existence of groupoid models

As one may guess, it is possible to relate groupoid models with universal F -actions.

Existence of groupoid models

As one may guess, it is possible to relate groupoid models with universal F -actions.

Idea

An action of a diagram of groupoid correspondences can be described as action of *partial homeomorphisms* of the space.

Existence of groupoid models

As one may guess, it is possible to relate groupoid models with universal F -actions.

Idea

An action of a diagram of groupoid correspondences can be described as action of *partial homeomorphisms* of the space.

The partial homeomorphisms of Y form an *inverse semi-group* $I(Y)$.

Existence of groupoid models

As one may guess, it is possible to relate groupoid models with universal F -actions.

Idea

An action of a diagram of groupoid correspondences can be described as action of *partial homeomorphisms* of the space.

The partial homeomorphisms of Y form an *inverse semi-group* $I(Y)$. Similarly, we can encode a diagram F of groupoid correspondences by an inverse semi-group $I(F)$.

Existence of groupoid models

Idea

The groupoid model can be expressed as a *transformation groupoid* of $I(F)$.

Existence of groupoid models

Idea

The groupoid model can be expressed as a *transformation groupoid* of $I(F)$.

Definition

Let S be an inverse semi-group, and Y be a space with an action $\vartheta: S \rightarrow I(Y)$.

Existence of groupoid models

Idea

The groupoid model can be expressed as a *transformation groupoid* of $I(F)$.

Definition

Let S be an inverse semi-group, and Y be a space with an action $\vartheta: S \rightarrow I(Y)$. The *transformation groupoid* $S \ltimes Y$ consists of

Existence of groupoid models

Idea

The groupoid model can be expressed as a *transformation groupoid* of $I(F)$.

Definition

Let S be an inverse semi-group, and Y be a space with an action $\vartheta: S \rightarrow I(Y)$. The *transformation groupoid* $S \ltimes Y$ consists of

- objects: elements of Y

Existence of groupoid models

Idea

The groupoid model can be expressed as a *transformation groupoid* of $I(F)$.

Definition

Let S be an inverse semi-group, and Y be a space with an action $\vartheta: S \rightarrow I(Y)$. The *transformation groupoid* $S \ltimes Y$ consists of

- objects: elements of Y
- morphisms:

$$y \xrightarrow{(s,y)} \vartheta_s(y)$$

is an equivalence class of pairs (s, y)

Existence of groupoid models

Idea

The groupoid model can be expressed as a *transformation groupoid* of $I(F)$.

Definition

Let S be an inverse semi-group, and Y be a space with an action $\vartheta: S \rightarrow I(Y)$. The *transformation groupoid* $S \ltimes Y$ consists of

- objects: elements of Y
- morphisms:

$$y \xrightarrow{(s,y)} \vartheta_s(y)$$

is an equivalence class of pairs (s, y) , where

$$(s_1, y) \sim (s_2, y) \iff \exists \text{ idempotent } e : s_1 e = s_2 e$$

Existence of groupoid models

Definition

- composition:

Existence of groupoid models

Definition

- composition:

$$\begin{array}{ccc} y & \xrightarrow{(s,y)} \vartheta_s(y) & \xrightarrow{(u,\vartheta_s(y))} \vartheta_u(\vartheta_s(y)) \\ & \parallel & \\ y & \xrightarrow{(us,y)} & \vartheta_{us}(y) \end{array}$$

Existence of groupoid models

Proposition (Meyer)

Let Ω be a universal F -action.

Existence of groupoid models

Proposition (Meyer)

Let Ω be a universal F -action. The transformation groupoid

$$I(F) \ltimes \Omega$$

is a groupoid model for F .

Existence of groupoid models

Proposition (Meyer)

Let Ω be a universal F -action. The transformation groupoid

$$I(F) \ltimes \Omega$$

is a groupoid model for F .

\rightsquigarrow showing the existence of universal F -actions will imply groupoid models exist

Existence of groupoid models

Existence of groupoid models

A general result from Category Theory states that

Existence of groupoid models

A general result from Category Theory states that

Lemma

Let D be a locally small cocomplete category.

Existence of groupoid models

A general result from Category Theory states that

Lemma

Let D be a locally small cocomplete category. Assume that there is a set $\Phi \subseteq \text{ob } D$ such that for any $d \in \text{ob } D$ there is $d \rightarrow c$ where $c \in \Phi$.

Existence of groupoid models

A general result from Category Theory states that

Lemma

Let D be a locally small cocomplete category. Assume that there is a set $\Phi \subseteq \text{ob } D$ such that for any $d \in \text{ob } D$ there is $d \rightarrow c$ where $c \in \Phi$. Then D has a terminal object.

Existence of groupoid models

A general result from Category Theory states that

Lemma

Let D be a locally small cocomplete category. Assume that there is a set $\Phi \subseteq \text{ob } D$ such that for any $d \in \text{ob } D$ there is $d \rightarrow c$ where $c \in \Phi$. Then D has a terminal object.

By applying the lemma to the category of F -actions, we attain

Existence of groupoid models

A general result from Category Theory states that

Lemma

Let D be a locally small cocomplete category. Assume that there is a set $\Phi \subseteq \text{ob } D$ such that for any $d \in \text{ob } D$ there is $d \rightarrow c$ where $c \in \Phi$. Then D has a terminal object.

By applying the lemma to the category of F -actions, we attain

Theorem (K., Meyer)

Any diagram $F: \mathcal{C} \rightarrow \mathbb{G}\mathbf{x}$ of groupoid correspondences has a universal F -action.

Existence of groupoid models

Since a groupoid model can be built from the universal F -action, we immediately obtain

Existence of groupoid models

Since a groupoid model can be built from the universal F -action, we immediately obtain

Corollary (K., Meyer)

Any diagram of groupoid correspondences has a groupoid model.

Existence of locally compact groupoid models

Next, we would like to know if the groupoid model is *locally compact*, in order to have applications in C^* -algebras.

Counterexamples

Counterexamples

Example

Consider the monoid $(\mathbb{N}_0, +)$ and the trivial groupoid \mathcal{G} (one object and one arrow).

Counterexamples

Example

Consider the monoid $(\mathbb{N}_0, +)$ and the trivial groupoid \mathcal{G} (one object and one arrow).

Then a diagram $F: (\mathbb{N}_0, +) \rightarrow \mathfrak{Gr}$ is determined by a single $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$.

Counterexamples

Example

Consider the monoid $(\mathbb{N}_0, +)$ and the trivial groupoid \mathcal{G} (one object and one arrow).

Then a diagram $F: (\mathbb{N}_0, +) \rightarrow \mathbb{G}\mathbf{r}$ is determined by a single $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$.

Since $\mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is a discrete set.

Counterexamples

Example

Consider the monoid $(\mathbb{N}_0, +)$ and the trivial groupoid \mathcal{G} (one object and one arrow).

Then a diagram $F: (\mathbb{N}_0, +) \rightarrow \mathbb{G}\mathbf{r}$ is determined by a single $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$.

Since $\mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is a discrete set.

It can be shown that Ω is given by

$$\prod_{n \in \mathbb{N}_0} \mathcal{X}$$

Counterexamples

Example

Consider the monoid $(\mathbb{N}_0, +)$ and the trivial groupoid \mathcal{G} (one object and one arrow).

Then a diagram $F: (\mathbb{N}_0, +) \rightarrow \mathbb{G}r$ is determined by a single $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$.

Since $\mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is a discrete set.

It can be shown that Ω is given by

$$\prod_{n \in \mathbb{N}_0} \mathcal{X}$$

If \mathcal{X} is infinite, then Ω is not locally compact.

Counterexamples

Example

Consider the monoid $(\mathbb{N}_0, +)$ and the trivial groupoid \mathcal{G} (one object and one arrow).

Then a diagram $F: (\mathbb{N}_0, +) \rightarrow \mathfrak{Gr}$ is determined by a single $\mathcal{X}: \mathcal{G} \leftarrow \mathcal{G}$.

Since $\mathcal{X} \rightarrow \mathcal{G}^0$ is a local homeomorphism, \mathcal{X} is a discrete set.

It can be shown that Ω is given by

$$\prod_{n \in \mathbb{N}_0} \mathcal{X}$$

If \mathcal{X} is infinite, then Ω is not locally compact.

\rightsquigarrow conditions on groupoid correspondences ensuring 'compactness'

Existence of locally compact groupoid models

Idea

Existence of locally compact groupoid models

Idea

We want a groupoid model to be locally compact

Existence of locally compact groupoid models

Idea

We want a groupoid model to be locally compact

\implies the universal F -action should take place on a locally compact Hausdorff space

Existence of locally compact groupoid models

Idea

We want a groupoid model to be locally compact

\implies the universal F -action should take place on a locally compact Hausdorff space

\rightsquigarrow for each F -action, we find a unique corresponding F -action on a locally compact Hausdorff space

Existence of locally compact groupoid models

Definition (Relative Stone-Čech compactification)

Existence of locally compact groupoid models

Definition (Relative Stone-Čech compactification)

Let B be a locally compact Hausdorff space, and X be a B -space, i.e. there is a continuous

$$X \rightarrow B$$

Existence of locally compact groupoid models

Definition (Relative Stone-Čech compactification)

Let B be a locally compact Hausdorff space, and X be a B -space, i.e. there is a continuous

$$X \rightarrow B$$

The *relative Stone-Čech compactification* $\beta_B X$ of X over B is the spectrum of

$$C_b(X) \cdot r^*(C_0(B)) \subseteq C_b(X)$$

Existence of locally compact groupoid models

Definition (Relative Stone-Čech compactification)

Let B be a locally compact Hausdorff space, and X be a B -space, i.e. there is a continuous

$$X \rightarrow B$$

The *relative Stone-Čech compactification* $\beta_B X$ of X over B is the spectrum of

$$C_b(X) \cdot r^*(C_0(B)) \subseteq C_b(X)$$

Example

When B is a singleton, the above notion recovers the usual *Stone-Čech compactification*.

Existence of locally compact groupoid models

Definition (Relative Stone-Čech compactification)

Let B be a locally compact Hausdorff space, and X be a B -space, i.e. there is a continuous

$$X \rightarrow B$$

The *relative Stone-Čech compactification* $\beta_B X$ of X over B is the spectrum of

$$C_b(X) \cdot r^*(C_0(B)) \subseteq C_b(X)$$

Example

When B is a singleton, the above notion recovers the usual *Stone-Čech compactification*.

Remark

The induced $\beta_B X \rightarrow B$ is a proper map.

Existence of locally compact groupoid models

Theorem (K., Meyer)

The relative Stone-Čech compactification β_B is left adjoint to the inclusion of proper Hausdorff B -spaces.

$$B\text{-Space} \begin{array}{c} \xrightarrow{\beta_B} \\ \xleftarrow{\perp} \end{array} B\text{-Space-Pro}$$

Existence of locally compact groupoid models

Theorem (K., Meyer)

The relative Stone-Čech compactification β_B is left adjoint to the inclusion of proper Hausdorff B -spaces.

$$B\text{-Space} \begin{array}{c} \xrightarrow{\beta_B} \\ \xleftarrow{\perp} \end{array} B\text{-Space-Pro}$$

Remark

In other words, the category of proper Hausdorff B -spaces is a *reflective subcategory* of the category of B -spaces.

Existence of locally compact groupoid models

Idea

Existence of locally compact groupoid models

Idea

Let F be a diagram of *proper locally compact* groupoid correspondences.

Existence of locally compact groupoid models

Idea

Let F be a diagram of *proper locally compact* groupoid correspondences.

For any F -action on a space Y , we extend uniquely to an F -action on $\beta_{\mathcal{G}_0} Y$.

Existence of locally compact groupoid models

Idea

Let F be a diagram of *proper locally compact* groupoid correspondences.

For any F -action on a space Y , we extend uniquely to an F -action on $\beta_{\mathcal{G}^0} Y$.

This construction 'lifts' the previous adjunction to

$$\begin{array}{ccc} \text{Category of } F\text{-actions} & \begin{array}{c} \xrightarrow{L} \\ \xleftarrow{\perp} \end{array} & \begin{array}{l} \text{Category of } F\text{-actions on proper} \\ \text{Hausdorff } \mathcal{G}^0\text{-spaces} \end{array} \end{array}$$

Existence of locally compact groupoid models

Idea

Let F be a diagram of *proper locally compact* groupoid correspondences.

For any F -action on a space Y , we extend uniquely to an F -action on $\beta_{\mathcal{G}^0} Y$.

This construction 'lifts' the previous adjunction to

$$\begin{array}{ccc} \text{Category of } F\text{-actions} & \begin{array}{c} \xrightarrow{L} \\ \xleftarrow{\perp} \end{array} & \begin{array}{l} \text{Category of } F\text{-actions on proper} \\ \text{Hausdorff } \mathcal{G}^0\text{-spaces} \end{array} \end{array}$$

\rightsquigarrow reflective subcategory

Existence of locally compact groupoid models

Then we apply the following general result from Category Theory:

Existence of locally compact groupoid models

Then we apply the following general result from Category Theory:

Lemma

The inclusion of a reflective (full) subcategory

$$D \hookrightarrow C$$

creates all limits that C admits.

Existence of locally compact groupoid models

Then we apply the following general result from Category Theory:

Lemma

The inclusion of a reflective (full) subcategory

$$D \hookrightarrow C$$

creates all limits that C admits.

This means if C has a limit, then D also has a limit, which is isomorphic to the limit in C .

Existence of locally compact groupoid models

Then we apply the following general result from Category Theory:

Lemma

The inclusion of a reflective (full) subcategory

$$D \hookrightarrow C$$

creates all limits that C admits.

This means if C has a limit, then D also has a limit, which is isomorphic to the limit in C .

In particular, terminal objects are a kind of limits.

Existence of locally compact groupoid models

Recall that universal F -actions are terminal objects.

Existence of locally compact groupoid models

Recall that universal F -actions are terminal objects.

Theorem (K., Meyer)

*Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram of **proper (locally compact)** groupoid correspondences.*

Existence of locally compact groupoid models

Recall that universal F -actions are terminal objects.

Theorem (K., Meyer)

*Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram of **proper (locally compact)** groupoid correspondences.*

The universal F -action takes place on a space Ω that is locally compact Hausdorff.

Existence of locally compact groupoid models

Recall that universal F -actions are terminal objects.

Theorem (K., Meyer)

*Let $F: \mathcal{C} \rightarrow \mathfrak{Gr}_{lc}$ be a diagram of **proper (locally compact)** groupoid correspondences.*

The universal F -action takes place on a space Ω that is locally compact Hausdorff.

Therefore, the groupoid model $I(F) \ltimes \Omega$ of F is locally compact.

Thank you!



Suliman Albandik,

A colimit construction for groupoids, Ph.D. Thesis, Georg-August Universität Göttingen, 2015.



Celso Antunes, Joanna Ko, and Ralf Meyer,

The bicategory of groupoid correspondences, New York J. Math. 28 (2022) 1329–1364.



Alcides Buss, Ralf Meyer, and Chenchang Zhu,

A higher category approach to twisted actions on C^ -algebras*, Proc. Edinb. Math. Soc. (2) 56 (2013), no. 2, 387–426.



Takeshi Katsura,

A class of C^ -algebras generalizing both graph algebras and homeomorphism C^* -algebras. I. Fundamental results*, Trans. Amer. Math. Soc. 356 (2004), no. 11, 4287–4322.



Joanna Ko, and Ralf Meyer,

Existence of groupoid models for diagrams of groupoid correspondences, Theory Appl. Categ. 41(13) (2024) 449–469.



Ralf Meyer,

Groupoid models for diagrams of groupoid correspondences, (2022), arXiv:2203.12068.



Volodymyr Nekrashevych,

C^ -algebras and self-similar groups*, J. Reine Angew. Math. 630 (2009), 59–123.