

PREDICTING MUMBAI'S AIR QUALITY INDEX BY MACHINE LEARNING

Dr. Kalpana Chaudhari

*Department of Electronics And
Telecommunication Engineering
Shah And Anchor Kutchhi
Engineering College
Mumbai,India
kalpanachaudhari@sakec.ac.in*

Dhruvi Joshi

*Department of Computer Science
Shah And Anchor Kutchhi
Engineering College
Mumbai,India
dhruvivipjoshi24@gmail.com*

Pratik Harugade

*Department of Electronics And
Computer Science
Shah And Anchor Kutchhi
Engineering College
Mumbai,India
harugadepratikpramod@gmail.com*

Kritik Jambusariya

*Department of Electronics And
Telecommunication Engineering
Shah And Anchor Kutchhi
Engineering College
Mumbai,India
kritikjambusariya@gmail.com*

Vaibhav Tiwari

*Department of Electronics And
Telecommunication Engineering
Shah And Anchor Kutchhi
Engineering College
Mumbai,India
tvaibhav541@gmail.com*

Abstract — The purity of air plays a major role on mortal health of people. Poor quality of air leads to many forms of diseases, most probably in the kids. Looking at the vulnerability of air quality standards the government further decides to take important acts for prevention of such diseases. Ancient ways of tackling diseases have a short range of success due to no access to huge sets of data. For this project, we have considered using machine's algorithm to augur Air quality pointer for megacity Mumbai. Our created model can analyze closer to 93 percent of aqi (air quality index), it further also predicts many oxides of carbon, nitrogen, sulfur and oxygen. Therefore, we as a country feel the need to have a machine to read the pollution levels for us to maintain the environment and to start taking respective precautions needed. In numerous artificial and civic regions at the moment, balancing the air index for human health is the biggest task right now. The burning of fossil energies, business patterns, and artificial variables all have a big impact on air pollution. We need to apply models that will record information regarding the attention of air pollutants because of the rising pollutant situations. The quality of people's lives is being impacted by the buildup of these dangerous chemicals in the air, particularly in metropolitan areas. Several lab tests have lately started using the

Big Data Analytics fashion as per rise of pollutants in air.

Keywords — *air quality indicator, direct retrogression model, mumbai, vaticination.*

I. INTRODUCTION

The contaminants and granules that make up the air around us are all veritably minute and, for the utmost part, undetectable to the naked eye. They could get into our more sensitive systems like the lungs or eyes. Those unnoticeable items from ground position ozone to patches constantly beget serious health issues. Only 10 percent of people on earth are safe from pollution, all others are victims of pollution, and an army of people die directly from it. To check air purity a device named aqi is used. It provides information on how dirty the girding air is and could become in the near future. Also, it informs you of any implicit negative health impacts. It substantially concentrates on the negative health impacts you could have many hours or days after breathing polluted air. For people to understand the range of air quality, an "air quality index" measuring unit was generated. It transforms the intricate information about the numerous adulterants and poisons in the air into a numerical number (indicator value), vocabulary, and color. Rising nations like India are facing huge pollution, due to the rapid population growth and profitable growth in metropolises. Mortal health is directly impacted by air pollution. In our nation, there has been a rise in public knowledge

of the same. Thus we as society must aim at improving the quality of air. A gas called sulfur dioxide, is one of the primary air adulterants.

It has an unwelcome, harsh smell and is tintless. It readily reacts with other chemicals to produce dangerous composites like sulfuric acid, sulfurous acid and affects humans when it's breathed in. In Section 1 we give a brief preface about the content we've chosen. In section 2, we outline our exploration and talk over our motives, outlining the possible benefits of being suitable to directly anticipate air quality. In Section 3, we present a critical reassessment of before applicable work. In Section 4, algorithms of ml are described, specifically Linear Regression. In sections 5 and 6, we talk more about the comparison of Kanpur City's AQI before and after COVID- 19. The dataset that we have used for every topic is mentioned in section 6. We describe the setting for the trial as well as the issues in Section 7 of our donation. And our paper ends in section 8.

Ethics for serviceable air quality situations have been established by public associations similar as the EPA, EU, and numerous others. The situations of the air's criteria adulterants are shown by the air quality indicator(AQI). The loftiest AQI reading for each of the individual criterion adulterants makes up the total AQI. The health enterprises related to exposure to the specific air quality are also indicated by AQI situations. The medical symptoms may appear soon after being exposed to polluted air or they may develop over time.

II. BACKGROUND AND MOTIVATION

A. In order to lower the climatic change, the dangerous chemicals in the air has to be reduced. Greenhouse elements can potentially harm the air condition by direct liberation in the air. If the rise of such chemicals are minimized it can help the overall Earth's temperature to be closer to 20 degree celsius.

B. Pollution affects our body. Even Though we can't see it, we are breathing in bad air. Tiny particles can get into our bodies and make us sick. Many people die each year, the cause is pollution. Huge number of humans don't even know that pollution is hurting them.

C. Children suffer the side effects of dirty air. To inhale oxygen rich air is our birthright. Just 7 % of the total, are breathing oxygen rich air. In the current situation even the one day olds are breathing not so good quality of air. These are one

of the major reasons for a child to suffer from the illness of the brain.

D. The most vulnerable suffer the most. The poorest and most marginalized communities are more likely to work and live in highly polluted and densely populated areas. An index called AQI is used to track daily quality and percent of chemical contents in air. It provides information on how polluted the present air and how far it could become more polluted later. To help people understand the current state of air quality a scale named the Air Quality Index was created. Converts complex information about all pollutants in the air into numbers (index values), and colors. In emerging countries like India, rapid population growth and the growth of profitable metropolises are creating problems for all types of pollution. People's health is directly affected by the toxins present in the air. In many countries, public knowledge regarding keeping the air clean is increasing. Predicted air quality forecasts can mitigate the impact of peak exposures on people and terrain. Therefore, one of society's top priorities is to improve predictions of air quality. One gas is sulfur dioxide. It readily reacts with other chemicals to form toxic compounds such as sulfuric acid, sulfites, which affect humans when inhaled.

E. We must keep our country clean and clear from pollution. People nowadays prefer car travel over trains and public services. It is believed by 2060, 70% of pollution will be generated in cities. Not only the air we breathe, but our health, safety, and general well-being as city dwellers are all affected by the way cities are built and moved.

III. PREVIOUS AND RELATED WORK

We reviewed similar research papers and summarized them in an attempt to identify their problems, how they work, and their shortcomings.

[1] This paper explores how greatly Air Quality Index affects human health. It specifically describes how it contributes to child-led health deterioration. The method used is SVR (Support Vector Regression) for diagnosis of purity of air. RBF (Radial Basis Function) is also used because it gets the best results in combination with SVR. This model predicts AQI with 93% accuracy.

[2] The authors talk about serious diseases caused by air quality. And giving us a general solution to these problems.

[3] Big data is used as a methodology, and various ml models, air purity ratios, help in knowing the worldly increasing air intoxicants. They use data from 3 cities in China to evaluate the model.

[4] The authors focus on building a project for regular 60 minutes detection of the toxins concentration in the air. In this study, a model named RNN-LSTM takes lively data with respect to time for knowing the presence of a quantity of chemicals. Timely data analysis done for the 12 air elements measurements in Visakhapatnam city were used for lab analysis. The study results share the idea that deep learning practices help in better performances. Simultaneous multivariate modeling of all 12 contaminants could improve this study. The proposed paradigm can also be extended by a two-way approach.

IV. LINEAR REGRESSION MODEL

Within reinforcement methods linear regression is a topic, its purpose is to diagnose a particular component and their correlation, this process is used in ict global medical care and making predictions for stock prices, weather prediction and landslides predictions amongst several other things.

An air closed-circuit television has to report the waste product levels over time so as to derive the AQI. The methodologies to translate close air concentrations to AQIs vary looking at the contaminants and also the country. There's an outline, a color code, and categories for every variety of major pollutants. Every close air quality space has prescribed medical consultations coupled with it. There appear to be 2 main stages in the development of Air Quality understanding. For each stuff, sub-indexes should be created. These subindexes are concatenated within the following stages (breakpoints).

A contaminant's index worth is basically its concentration, given as a proportion of the concentration, expressed as a proportion of the relevant the best potential variety for every waste product sub-index is employed to represent the site's total AQI. Supported whether or not a private contamination will negatively have an effect on population, terminus concentrations area unit calculable victimization statistics from data-based studies and NAAQS. Many marked area units are so contaminant-dependent.

There appear to be six AQI categories as per India's AQI. Pollution levels may be good, fair, moderately unhealthy, poor, or severe. The mathematical formula for regression toward the mean is: $y = a_0 + a_1x$, wherever y is that variable quantity.

x = unrelated variable

Line intersection at a_0 .

Together with the variable x , the variable quantity y increases too.

The principal methodology for computing constant values

- 1] Initialize the parameters.
- 2] Having used the parameter, compute the amount present.
- 3] Then compute the individual difference present in intervals of parameters..
- 4] The exogenous variables with reference to a_0 and a_1 need to be entered.
- 5] Add the worth comparable to every variety.
- 6] Modify a_0 and a_1 's values.

Contaminants	Subfix.	Assemblage
X1	$I_1 = f_1(X_1)$	$I = f$
X2	$I_2 = f_2(X_2)$	$(I_1, I_2, \dots, I_n)$
Xn	$I_n = f_n(X_n)$	
	Step1	Step2

fig.1. AQI Development

V. COMPARING THE AIR QUALITY INDEX OF KANPUR CITY BEFORE AND AFTER COVID-19

In India, Uttar Pradesh state has a man-made megacity named Kanpur. Kanpur, which was established in 1207, rose to elevation among British India's service and marketable posts. Furthermore, Kanpur serves as Uttar Pradesh's pecuniary cynosure. We have made analysis for the kanpur city's aqi of before covid-19 and after covid-19.

The model has been trained using instantaneous on-screen records gathered from the Control of Pollution Board which has been appointed by the Central Government as well as World Air Quality Index of Kanpur. The dataset for the 1st 21 days of lockdown is considered from Mar 25th, 2020 to Apr 14, 2020. We've made the analysis of air quality of Kanpur metropolis beforehand and after Covid- 19.

Pandas lib has been used for analysis. In the dataset collected pm10 values were not defined. Multiple rows were empty and these rows were filled by null value. For this we had to pull data from dates Mar,2020 to Apr,2020. For this we made a new variable and called the function(date)> '2020-03-24'. This would respond in collection of 7 days of March and 14 days of April. Sorted the data and included only the needed 14 days data of April,2020.




Fig 2. Line graph for value PM2.5

Generate a new index and allocate the dates from 2020-03-05 to 2020-03-25. This is done to root the dates past lockdown. Further we're comparing the PM2.5 values for beforehand and after Covid- 19 for air quality analysis. Generate a new variable and input all values of PM2.5 beforehand and after Covid- 19. Plot the graph for PM2.5. Consequently, from our analysis the air pollution after lockdown has dropped staggeringly.



Fig 3. Line graph for value PM2.5 before and after covid-19.

VI. DATA DESCRIPTION

The used recordsets were maintained by the CPCB and have been stored safely in their data libraries. Few of them are mentioned below. Hourly enthrallment of fine granular substance having radius of lower or equitable to 2.5 micrometers (PM2.5) and 10 micrometers or lower (PM10, according to data from the CPCB in India) are included in the data (PM10). On the basis of these figures, they calculate the Air Quality Index. Data splitting was done as follows: 20 percent was dogged for testing and the remaining 80 percent data was being fixed for training purposes. Preprocessing and trait selection, only Maharashtra state's data were examined and algorithms were used.

VII. EXPERIMENTAL STUDY

The model has been trained using instantaneous on-screen records gathered from the Control of Pollution Board which has been appointed by the Central Government specially for analyzing the Air Quality Index of Mumbai City from Jan 1, 2022 to Apr 17, 2022. Mumbai (Bombay) is the most populated megacity in India. The collected information namely: Average visibility (VV), Average Wind Speed (V), Average temperature (T), Total downfall or snowmelt (PP), Average relative moisture (H), Maximum temperature (TM), Maximum sustained wind speed (VM), Minimal temperature (tm), Maximum speed of wind (VG), and Atmospheric Pressure at Ocean Position (SLP). Also, in addition to the above specifications the rows which were assigned to the predictor variable P.M. 2.5 bearing null properties have been removed. The pollutants within the air and their units present are employed in this

exploration disquisition are displayed in Tables 1

Sno	Parameters	Unit
1	P.M. 2.5 (Fine Particulate matter 2.5)	ug/m3
2	P.M. 2.5-10 (Particulate matter value lying between 2.5 to 10)	ug/m3
3	Ozone	ug/m3
4	SO2 (Sulfur di-oxide)	ug/m3
5	NO2(Nitrogen di-oxide)	ug/m3

Table 1.Different chemical compounds present in air.

The predictive model's performance is impacted by uncleaned data, hence data from the above mentioned source must be preprocessed. Missing values increase the data's noise level. The mean values of the corresponding parameter were used to impute missing values for the proposed model. As a result, the final dataset taken into consideration has a small number of feature samples. Our objective is to create a model with highest accuracy for hourly detection of concentration of air pollutants.Of the complete dataset 75% is used in training purpose .To locate Mumbai city on the map we have made use of a cartopy library.




Fig.4 Location of Mumbai with AQI on world map.

Air pollutants and their probable amount in the atmosphere of Mumbai




Fig. 5. Pie diagram for air measurements




Fig.6. Heatmap for air components.




Fig. 7. Histogram for every air measurements

VIII. CONCLUSION

We now talk over our findings and vital takeaways. We've learnt to make an ML model to foretell the AQI at the position. And also, we introduce how important AQI predicting is. Also, We learned many new modules and concepts regarding machine learning libraries. In turn we also got to know how technology plays a part in the development of mortal society and lead them towards a bright future. The cost function that computes the accuracy level determines the life of the model.

The cost function helps in knowing the counterplot within the input variable and affair variable. This mapping variable is referred to as the thesis function. Air quality is a major factor, which is of concern for each and every mortal and to get accurate AQI is the need of the moment. Technological, or profitable(or request- grounded) results make up the maturity of enterprise to reduce the pollution of air. Regulatory Alternatives try to decrease the air pollution through supervision and penalties include legislation and the establishment of governmental associations.

Pre-warmed chemicals change the transformers, gasoline, and the pollution emitted from them. SUVs square captures a high value of advancements in tech fields. These choices, that square measure driven by request problems, provide businesses the liberty to settle on the foremost cost-efficient results to fulfill set environmental goals. This study aims to manufacture a system for checking an hourly air

impurity immersion exploitation. The effectiveness of the results from direct retrogression should be objectively assessed. We are able to take action for metropolises like Pune and Bombay wherever the attention of so2 is rising, so as to anticipate the values of so2. As these patches square measure to blame for a range of health consequences, such things need to be proved within the future for health guarantee. The requirement for mortal commerce is reduced by the aptitude of cubic centimeter models to mechanically overlook huge quantities of knowledge and select vital attributes. Given massive datasets, cubic centimeter models could also be additionally correct than standard applied math designs. Similar Models are very helpful for AQI prognosticating tasks for an extended time.

ACKNOWLEDGEMENT

We the authors wholeheartedly thank University of Naples Federico II, Italy and to our college Shah And Anchor Kutchhi Engineering College for this amazing opportunity to work with the Italian professors and learn about urban development problems and to tackle them using machine learning. Further thanking our mentor Dr. Kalpana Chaudhari Ma'am for their guidance and support. We are deeply grateful for their support and contribution to our project and paper.

REFERENCES

- [1] Samayan Bhattacharya, Sk Shahnawaz, "Using Machine Learning to Predict Air Quality Index in New Delhi", Cornell University, 2021.
- [2] T. Madan, S. Sagar and D. Virmani, "Air Quality Prediction using Machine Learning Algorithms –A Review," 2020 2nd International Conference on Advances in Computing, Communication Control and Networking (ICACCCN), 2020.
- [3] Gaganjot Kaur Kang, Jerry Zeyu Gao, Sen Chiao, Shengqiang Lu, and Gang Xie, "Air Quality Prediction: Big Data and Machine Learning Approaches", International Journal of Environmental Science and Development, Vol. 9, No. 1, January 2018.
- [4] K Srinivasa Rao, G. Lavanya Devi, N. Ramesh, "Air Quality Prediction in Visakhapatnam with LSTM based Recurrent Neural Networks", International Journal of Intelligent Systems and Applications (IJISA), Vol. 11, No. 2, pp. 18-24, 2019.

- [5] H. Drucker, C. J. Burges, L. Kaufman, A. Smola, V. Vapnik et al., "Support vector regression machines," *Advances in neural information processing systems*, vol. 9, pp. 155–161, 1997.
- [6] R. Sharda and R. Patil, "Neural networks as forecasting experts: an empirical test," in *Proceedings of the International Joint Conference on Neural Networks*, vol. 2. IEEE, 1990, pp. 491–494.
- [7] C.-M. Vong, W.-F. Ip, P.-k. Wong, and J.-y. Yang, "Short-term prediction of air pollution in macau using support vector machines," *Journal of Control Science and Engineering*, vol. 2012, 2012.
- [8] L. Cagliero, T. Cerquitelli, S. Chiusano, P. Garza, G. Ricupero, and X. Xiao, "Modeling correlations among air pollution-related data through generalized association rules," in *2016 IEEE International Conference on Smart Computing (SMARTCOMP)*. IEEE, 2016, pp. 1–6.
- [9] W. Li, D. Kong, and J. Wu, "A new hybrid model fpa-svm considering cointegration for particular matter concentration forecasting: a case study of kunming and yuxi, china," *Computational intelligence and neuroscience*, vol. 2017, 2017.
- [10] D. C. Payne-Sturges, M. A. Marty, F. Perera, M. D. Miller, M. Swanson, K. Ellickson, D. A. Cory-Slechta, B. Ritz, J. Balmes, L. Anderko et al., "Healthy air, healthy brains: advancing air pollution policy to protect children's health," *American journal of public health*, vol. 109, no. 4, pp. 550–554, 2019.
- [11] Y. Gonzalez, C. Carranza, M. Iniguez, M. Torres, R. Quintana, A. R. Osornio-Vargas, C. Gardner, S. Sarkar, and S. Schwander, "Inhaled air pollution particulate matter in alveolar macrophages alters local pro-inflammatory cytokine and peripheral production in response to mycobacterium tuberculosis," in *B17. MYCOBACTERIAL HOST DEFENSES*. American Thoracic Society, 2017, pp. A2901–A2901.