

# PowerNumbers.jl: a fast approach to automatic asymptotics

SHEEHAN OLVER, Imperial College

MATTHEW REES, University College London

We develop a scheme for quick arithmetic on asymptotic series, evaluated to just one or two terms. We give a full description of the algebraic rules for these "Power Numbers", with justification of sufficient equivalence to asymptotic algebra. Some example applications follow, including evaluation of rational functions at infinity.

CCS Concepts: • **Mathematics of computing** → **Solvers**.

Additional Key Words and Phrases: none

## ACM Reference Format:

Sheehan Olver and Matthew Rees. 2019. PowerNumbers.jl: a fast approach to automatic asymptotics. 1, 1 (September 2019), 4 pages. <https://doi.org/10.1145/nnnnnnn.nnnnnnn>

## 1 INTRODUCTION

## 2 DESCRIPTION

*Definition 2.1.* Let  $K$  be a field. The set of functions of  $\epsilon \in [0, \infty)$ ,

$$\mathbf{PN}_\epsilon^K = \{a\epsilon^\alpha + b\epsilon^\beta : a, b \in K; \alpha, \beta \in \mathbb{R} \cup \{\infty\}; \alpha \leq \beta\}$$

is called the set of **Power Numbers**.

In the case  $\alpha = \beta$ , we write only one term  $(a + b)\epsilon^\beta$ . Notationally,  $\epsilon^0$  is omitted. We enforce the equality  $0\epsilon^\alpha + b\epsilon^\beta = b\epsilon^\beta$ . However, in general,  $a\epsilon^\alpha + 0\epsilon^\beta \neq a\epsilon^\alpha$ . By defining  $(+, *)$  below, we acquire the double monoid

$$\mathbf{PN}_\epsilon^K = (\mathbf{PN}_\epsilon^K, +, *)$$

---

Authors' addresses: Sheehan Olver, [s.olver@imperial.ac.uk](mailto:s.olver@imperial.ac.uk), Imperial College, London, England, SW7 2AZ; Matthew Rees, [matthew.rees.16@ucl.ac.uk](mailto:matthew.rees.16@ucl.ac.uk), University College London, London, England, WC1E 6BT.

---

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from [permissions@acm.org](mailto:permissions@acm.org).

© 2019 Association for Computing Machinery.

XXXX-XXXX/2019/9-ART \$15.00

<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

## 2.1 Addition

$+$  :  $\text{PN}_\epsilon^K \times \text{PN}_\epsilon^K \rightarrow \text{PN}_\epsilon^K$  is described by an algorithm.

**Data:**  $a\epsilon^\alpha + b\epsilon^\beta, c\epsilon^\gamma + d\epsilon^\delta \in \text{PN}_\epsilon^K$ ; assume WLOG that  $\beta \leq \delta$

**Result:**  $(a\epsilon^\alpha + b\epsilon^\beta) + (c\epsilon^\gamma + d\epsilon^\delta) = p\epsilon^\zeta + q\epsilon^\eta \in \text{PN}_\epsilon^K$

**if**  $\beta = \delta$  **then**

**if**  $\gamma = \beta$  **then**

$p = a, q = b + c + d, \zeta = \alpha, \eta = \beta$

**else if**  $\alpha < \gamma < \beta$  **then**

$p = a, q = c, \zeta = \alpha, \eta = \gamma$

**else if**  $\gamma = \alpha$  **then**

$p = a + c, q = b + d, \zeta = \alpha, \eta = \beta$

**else**

$p = c, q = a, \zeta = \gamma, \eta = \alpha$

**end**

**else**

**if**  $\beta < \gamma$  **then**

$p = a, q = b, \zeta = \alpha, \eta = \beta$

**else if**  $\gamma = \beta$  **then**

$p = a, q = b + c, \zeta = \alpha, \eta = \beta$

**else if**  $\alpha < \gamma < \beta$  **then**

$p = a, q = c, \zeta = \alpha, \eta = \gamma$

**else if**  $\gamma = \alpha$  **then**

$p = a + c, q = b, \zeta = \alpha, \eta = \beta$

**else**

$p = c, q = a, \zeta = \gamma, \eta = \alpha$

**end**

**end**

### Algorithm 1: Summing Power Numbers

The additive identity for Power Numbers is  $0\epsilon^\infty$ .

## 2.2 Multiplication

$*$  :  $\text{PN}_\epsilon^K \times \text{PN}_\epsilon^K \rightarrow \text{PN}_\epsilon^K$  can be most simply expressed as addition of Power Numbers:

$$(a\epsilon^\alpha + b\epsilon^\beta) * (c\epsilon^\gamma + d\epsilon^\delta) = (ac\epsilon^{\alpha+\gamma} + ad\epsilon^{\alpha+\delta}) + (bc\epsilon^{\beta+\gamma} + bd\epsilon^{\beta+\delta}) = p\epsilon^\zeta + q\epsilon^\eta \in \text{PN}_\epsilon^K$$

The multiplicative identity for Power Numbers is  $1 + 0\epsilon^\infty$ .

## 2.3 Pseudo-Negation

While  $\text{PN}_\epsilon^K$  is a monoid under both addition and multiplication, we can define two operations that have some of the properties we expect for subtraction and division.

$-$  :  $\text{PN}_\epsilon^K \rightarrow \text{PN}_\epsilon^K$  is defined as:

$$-(a\epsilon^\alpha + b\epsilon^\beta) = (-a)\epsilon^\alpha + (-b)\epsilon^\beta$$

Naturally,  $-$  :  $\text{PN}_\epsilon^K \times \text{PN}_\epsilon^K \rightarrow \text{PN}_\epsilon^K$  is defined as  $A + (-B)$ , where  $A, B \in \text{PN}_\epsilon^K$ . Note that for  $\beta \neq \infty$ , we have  $(a\epsilon^\alpha + b\epsilon^\beta) - (a\epsilon^\alpha + b\epsilon^\beta) = 0\epsilon^\beta \neq 0\epsilon^\infty$ .

## 2.4 Pseudo-Inverse

The multiplicative pseudo-inverse is slightly more complicated.

**Data:**  $a\epsilon^\alpha + b\epsilon^\beta \in \text{PN}_\epsilon^K$

**Result:**

$$\frac{1}{a\epsilon^\alpha + b\epsilon^\beta} = p\epsilon^\zeta + q\epsilon^\eta \in \text{PN}_\epsilon^K$$

**if**  $\alpha = \beta$  **then**

$p = 0, q = \frac{1}{a+b}, \zeta = -\alpha, \eta = -\alpha$

**else**

$p = \frac{1}{a+b}, q = -\frac{b}{a^2}, \zeta = -\alpha, \eta = \beta - 2\alpha$

**end**

### Algorithm 2: Multiplicative Inversion

This matches the Dual Numbers case, where  $\alpha = 0, \beta = 1$ . Similarly to subtraction,

$$\frac{a\epsilon^\alpha + b\epsilon^\beta}{a\epsilon^\alpha + b\epsilon^\beta} \neq 1 + 0\epsilon^\infty$$

## 2.5 Exponentiation & Analytic Functions

The definition of  $A^m$  for  $A \in \text{PN}_\epsilon^K, m \in \mathbb{Z}$  follows naturally from the above definitions of multiplication and division.

Since we will generally consider  $\epsilon$  to be small, we can extend any analytic function  $h : K \rightarrow K$  to  $h : \text{PN}_\epsilon^K \rightarrow \text{PN}_\epsilon^K$  for  $\alpha \geq 0$ . This is via Taylor series:

$$\begin{aligned} h(a\epsilon^0 + b\epsilon^\beta) &= h(a + b\epsilon^\beta) = h(a) + b\epsilon^\beta h'(a) \\ h(a\epsilon^\alpha + b\epsilon^\beta) &= h(0) + a\epsilon^\alpha h'(0) \quad \text{where } \alpha \neq 0 \end{aligned}$$

## 2.6 Asymptotic Series

Consider an asymptotic series that is also a Hahn series; that is,

$$S(z) = \sum_{n=1}^{\infty} a_n z^{\alpha_n} \quad \text{as } z \rightarrow 0 \quad (1)$$

where the  $\alpha_n \in \mathbb{R}$  are strictly increasing. The primary significance of Power Numbers comes from the following

**PROPOSITION 2.2.** *Let  $f(z) = az^\alpha + bz^\beta + o(z^\beta), g(z) = cz^\gamma + dz^\delta + o(z^\delta)$  be series of the form in (1). Define the associated Power Numbers,  $F = a\epsilon^\alpha + b\epsilon^\beta, G = c\epsilon^\gamma + d\epsilon^\delta$ . Then, where  $\cdot$  is one of  $(+, -, *, /)$  we have that*

$$f(z) \cdot g(z) = pz^\zeta + qz^\eta + o(z^\eta)$$

$$\Leftrightarrow$$

$$F \cdot G = p\epsilon^\zeta + q\epsilon^\eta$$

Furthermore, for any analytic function  $h : K \rightarrow K$  we have

$$h(f(z)) = pz^\zeta + qz^\eta + o(z^\eta)$$

$$\Leftrightarrow$$

$$h(F) = p\epsilon^\zeta + q\epsilon^\eta$$

The proof is trivial.

### **3 EXAMPLES**

#### **3.1 Rational Functions At Infinity**

### **4 CONCLUSION**