


Position × residue enrichment, coloured by significance (positions 4–12)  
colour =  $-\log_{10} p$ , chi-square as reported in the workbook · cell text = fold change · deeper red = more significant



n = 20 P/G-D/E/T substrates vs n = 193 non-substrate controls carrying the same motif. Rows are ordered by chemical class — Basic K H R · Acidic D E · Aromatic F Y W · Aliphatic G P A M · Hydrophobic V L I · Polar S T C N Q.

Cell text is the fold change (substrates / controls) of every cell at p < 0.05, under the star bin of its uncorrected p: \* p < 0.05, \*\* p < 0.01, \*\*\* p < 0.001. Grey = residue absent from both groups, no test possible.

No depleted cell reaches p < 0.05 anywhere in this dataset, so every coloured cell above is an enrichment. 12 of the 179 testable cells shown here reach p < 0.05 uncorrected, against 9 expected by chance alone — and only R at 7, R at 8 survive BH-FDR (q < 0.05) — the rest is not separable from that expectation.

This panel is the N-terminal window only — positions 4–12, out to the 12th amino acid of the 24-mer. The crop is a display crop: every cell's p is unchanged, and q is still BH-FDR across all 416 cells of the full 4–24 matrix rather than recomputed inside the window — recomputing it would be choosing the region after seeing the data.