


## Chemical class × position, coloured by significance (positions 4–12)

colour =  $-\log_{10} p$ , chi-square as reported in the workbook · Basic K H R · Acidic D E · Aromatic F Y W · Aliphatic G P A M · Hydrophobic V L I · Polar S T C N Q



n = 20 P/G-D/E/T substrates vs n = 193 non-substrate controls carrying the same motif.

Cell text is the fold change (substrates / controls) of every cell at  $p < 0.05$ , under the star bin of its uncorrected p: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

No depleted cell reaches  $p < 0.05$  anywhere in this dataset, so every coloured cell above is an enrichment. 4 of the 54 testable cells shown here reach  $p < 0.05$  uncorrected, against 3 expected by chance alone — and only Basic at 8, Basic at 5, Basic at 7 survive BH-FDR ( $q < 0.05$ ) — the rest is not separable from that expectation.

This panel is the N-terminal window only — positions 4–12, out to the 12th amino acid of the 24-mer. The crop is a display crop: every cell's p is unchanged, and q is still BH-FDR across all 126 cells of the full 4–24 matrix rather than recomputed inside the window — recomputing it would be choosing the region after seeing the data.