

Northern Humboldt model validation

Field	Entry
Region	Humboldt Current, LME_013. Selected-model evaluation of the northern Peruvian food web. Regional workbook
Catch source	Sea Around Us, 1950–2019; confidence reference: 2019 landings. All 218 exact labels, including zero catch and unidentified categories. Catch and classic inputs Reference arithmetic
Selected article	Chiaverano et al. (2018). Evaluating the role of large jellyfish and forage fishes as energy pathways, and their interplay with fisheries, in the Northern Humboldt Current System. Progress in Oceanography 164, 28–36. DOI 10.1016/j.pocean.2018.04.009. Source PDF DOI
Other known articles	Neira et al. (2026), Chilean Patagonia 1980–2020 food-web modelling, DOI 10.1016/j.pocean.2025.103631. Its static 1980 model is retained as previous-selection evidence. Tam et al. (2008), Northern Humboldt trophic linkages under La Niña and El Niño, is a predecessor; its stocks cannot replace this model without a verified crosswalk. Selected-model review
Selected model	Chiaverano2018_detailed_Northern_Humboldt_1995_1998; selected audited detailed Northern Humboldt baseline, 1995–1998. 39 ecological stocks plus two source fleet nodes, over 165,000 km ² at 4°S–16°S and up to 111 km offshore. The paper counts 36 living groups and three detritus pools. Anchovy eggs (36) are a nonfeeding pool in the supplement, giving four operational detritus pools in this translation. The runtime adds one synthetic diet_import source. Canonical source JSON Computational input
Other models from the same article	The aggregated baseline has 24 ecological groups plus one pooled fishery node (25 source nodes), over the same area and period. It merges source groups and uses rounded printed values. Full detailed detritus/fleet routing is retained only in the resolved version. The four structural scenarios alter jellyfish consumption, forage-fish consumption or fishing rates; they are separate scenarios, not replacement static baselines.
Selection rationale	Article - Requested source review to test the local source evidence and its applicability to Northern Humboldt catches. Model - Prioritize the resolved supplement because its distinct taxa, hake stages, two fleets and detailed diets/routing permit a more faithful audit than the aggregated alternative. The user selected this audited model after auto returned finite, nonnegative SPPR in all three methods. Selection does not establish scientific validity or researcher validation.

Field	Entry
Model extraction	The PDF, XLS supplement and Word figures/tables are present and verified; historical unavailable-file metadata is obsolete. Printed Chrysaora plocamia (7) diet sums to 1.045 and is preserved in canonical JSON and imports; runtime normalization is separate. Gelatinous zooplankton (6) source biomass and P/B imply production 0.00530 t/km²/year, far below its predation demand. Sardine landings are 5.6513425 in the supplement but 1.4 in the paper's stated balanced baseline. Neither conflict has a supported numerical repair. GS is derived from source assimilation efficiency; explicit living BA=0 follows the stated steady state. Runtime detritus EE is forced to one and pool BA is derived. Internal fishery discards are retained in source evidence but their returned-flow ancestry is unsupported in the SPPR translation. The runtime loader also replaces eight native detritus-fate cells with pool self-identity; the exact changes remain explicit. Fresh reconstruction audit Source conflicts Loader matrix changes
GE diagnostics	FAIL overall: input FAIL; convergence FAIL; SPPR balance FAIL. No negative SPPR entries across any basal-source column or recipient, including unfished groups. rho_living = 0.7000 Unpooled recycling b = 1.5918; it does not converge. Auto pools four detritus columns; pooled b = 0.7659. Detritus SPPR: Anchovy eggs (36) = 6.1037; Fishery offal (37) = 6.1037; Pelagic detritus (38) = 6.1037; Benthic detritus (39) = 6.1037. Strict model balance and SPPR balance both fail. PP gap 16.4%. Full GE return Labeled GE matrix
TE diagnostics	FAIL overall: input FAIL; convergence WARN; SPPR balance FAIL. No negative SPPR entries across any basal-source column or recipient, including unfished groups. rho_living = 0.7316 Detritus SPPR: Anchovy eggs (36) = 0.0000; Fishery offal (37) = 0.0000; Pelagic detritus (38) = 0.3043; Benthic detritus (39) = 0.0000. Strict model balance and SPPR balance both fail. PP gap 9.4%. The engine cannot recredit EE=0 losses with four operational detritus pools; auto does not change this TE formulation. Full TE return Labeled TE matrix
SPPR calculation	[Researcher calculation choices and notes]
Geographic fit	Approximate A. Region covered by study: 6.0–8.0%. Approximate B. Study covered by region: 90–100%. The northern shelf domain leaves most of this LME outside the model. Boundary assessment
Temporal fit	Static 1995–1998 baseline applied to catch years 1950–2019; 2019 is 21–24 years later. Fixed coefficients and source catch/biomass allocations do not represent annual ecosystem or catch-composition changes. Publication year 2018 is separate from model period.

Field	Entry
Other	All direct calls explicitly use <code>det_collapse_mode='auto'</code> with the audited constructor, normalization, <code>EE=0</code> fix and closed-detritus settings unchanged. With Egestion remains overall FAIL: input FAIL, convergence FAIL, SPPR balance WARN; <code>rho_living</code> = 0.5600; unpooled <code>b</code> = 1.1098; pooled <code>b</code> = 0.8579; pooled detritus SPPR = 11.0035; PP gap 4.9%. All its SPPR entries are nonnegative, including unfished groups. Auto eliminates the previous negative coefficients but does not resolve source production conflicts, native-flow translation limits or strict balance failures. Selected-model regional arithmetic is displayed provisionally with production eligibility false, pending researcher review. Full direct report Candidate calculations
Open issues and next action	
Review and reproducibility	Researcher name: [name] review date: [dd/mm/yyyy]

Taxon mapping and coverage

Reference: 2019 landings; all recorded labels, all model groups and no taxon exclusions.

39 source ecological groups plus two fleet records and one computational import; 218 catch taxa.

Simple-chain PPR totals 277,099,551.43 t C in 2019.

34 labels lack classic TL/coefficient inputs; all have zero 2019 landings and contribute zero annual PPR. Their coefficients remain unavailable.

Method: independent simple trophic chain using the saved regional coefficients.

[Excel taxon appendix](#) | [Descriptive Sources sheet](#)

Overall confidence	Taxa (n)	Catch (%)	Simple-chain PPR (%)
High	70	87%	71%
Medium	104	9.4%	24%
Low	17	0.24%	0.26%
Very low	27	3.2%	4.9%
Unresolved	0	0%	0%

Overall confidence uses the weaker required membership or allocation component.

Group assignment rules

Plain-language rule	Confidence	PPR percentage
Explicit inclusion of the taxon in the source model	High	71%
Supported extension from the source's listed representative taxa	Medium	7.3%
Source membership connected through a verified synonym or documented spelling correction	High	6.7%
Supported ecological assignment requiring a group-boundary interpretation	Medium	6.6%
Closest represented ecological or taxonomic analogue	Very low	3.3%
Assumed eligible group set supported by taxonomy and ecology	Medium	3.2%
Approximation of a broad reporting category across plausible model pools	Very low	1.6%
Partial or competing ecological fit	Low	0.26%
Unambiguous fit to a documented taxonomic or ecological group	High	0.15%

Membership evidence: exact supplement group definitions, source taxonomic lists and authoritative synonym/habitat records. [Complete mapping evidence](#)

Allocation weight rules

Plain-language rule	Confidence	PPR percentage
One group; no allocation split required	High	86%
Allocation proportional to native source-model landings	Medium	14%
Complete source-model biomass fallback after unusable landings	Medium	0.046%

Weights use measured composition only when applicable. The source-model landings proxy is used before biomass; assumed proportions are fixed across catch years, with stage, fishery and geographic transfers recorded. [Allocation evidence](#)

Each percentage is the independent simple-chain PPR associated with taxa using that rule.

Unresolved taxa: none. 27 Very low decisions account for 3.2% of catch and 4.9% of independent simple-chain PPR.

Very low decisions

Affected taxa	Why confidence is very low
Actinopterygii; Gadiformes; Marine finfishes not identified; Marine fishes not identified; Marine groundfishes not identified; Marine pelagic fishes not identified; Perciformes; Scorpaeniformes	Broad fish reporting labels have unknown composition across named and residual pools, historical taxonomic scope, and potentially unrepresented deep-water members; native landings approximate their mixtures.
Batoidea; Chondrichthyes; Elasmobranchii; Myliobatidae	Broad/historical cartilaginous labels cross benthic and water-column pools with unrepresented chimaera or filter-feeding ray components.
Callorhynchus callorynchus	Benthic chimaera uses an elasmobranch analogue across the Holocephali/Elasmobranchii boundary.
Decapoda; Malacostraca; Miscellaneous marine crustaceans	Benthic/pelagic mixture uses macrozooplankton/benthos biomass because native landings are zero; a 2–20 mm size mismatch remains.
Dissostichus eleginoides; Hoplostethus atlanticus	Deep, cold-water fishes use a medium-demersal analogue with population, depth and size mismatches.
Galatheidae; Grimothea monodon	Squat lobsters lack a source pool; macrozooplankton/benthos biomass weights retain the 2–20 mm size mismatch.
Galaxias maculatus	Amphidromous galaxiid lacks a corresponding source population; a coastal small-fish analogue is assumed.
Macruronus magellanicus; Merluccius australis; Merluccius gayi; Micromesistius australis	Non-Peruvian hake/cod-like fishes use Peruvian hake groups; species, southern ecology and size boundaries differ, and native stage landings are transferred.
Mobula birostris	Pelagic filter-feeding manta uses the closest chondrichthyan pool despite the source apex-predator correspondence.
Ophichthus	Snake eel mapped to unnamed conger pool despite taxonomic identity differing from the regional congrio interpretation.


Geographic evidence

The study covers a northern coastal shelf segment of the target LME. The shaded source boundary was traced approximately against its graticule; reported area and traced area differ, so the A/B ranges express boundary uncertainty.

Region area

Target region boundary

Humboldt Current boundary and Northern Humboldt study



Humboldt Current LME_013 Sea Around Us boundary and approximate study trace in common geographic coordinates. [Boundary evidence](#)

Study area in the article

Study boundary for the selected model




Fig. 1. Map of the study area (shaded area) covering from 4°S to 16°S, and up to 111 km (60 nm) from the coastline, in the Northern Humboldt Current Ecosystem.

Chiaverano et al. (2018), original Figure 1, PDF page 2 / printed page 29. Hatching marks the 4°S–16°S coastal domain, up to 111 km offshore. [Source PDF](#)