

Appendix S3. Table S3

Basic input parameters of the Mediterranean marine ecosystem model for the 1990s period. O=Other; f.=fish; NC=Non commercial; OC=Other commercial.

# FG	Group name	Trophic level	Biomass (t/km ²)	Production / biomass (/year)	Consumption / biomass (/year)	Ecotrophic Efficiency	Production / consumption (/year)
1	Bottlenose dolphin	4.48	0.002	0.067	22.50	0.05	0.00
2	Striped dolphin	4.35	0.003	0.065	35.99	0.65	0.00
3	Common dolphin	4.29	0.000	0.045	29.24	0.01	0.00
4	Fin whale	3.15	0.009	0.022	7.25	0.00	0.00
5	Sperm whale	4.60	0.003	0.050	8.15	0.32	0.01
6	Deep Sea cetaceans	4.54	0.004	0.075	16.56	0.40	0.00
7	Monk seal	4.71	0.000	0.082	27.32	0.58	0.00
8	Pelagic Seabird	3.72	0.000	0.395	114.61	0.59	0.00
9	Gull&Cormorant	3.17	0.000	0.307	158.17	0.20	0.00
10	Tern	3.25	0.000	0.284	141.76	0.33	0.00
11	Loggerhead turtle	3.08	0.002	0.270	2.52	0.95	0.11
12	Green turtle	2.12	0.000	0.200	9.84	0.99	0.02
13	Pelagic shark	4.50	0.082	0.098	1.38	0.02	0.07
14	NC large pelagic	3.64	0.088	0.112	1.23	0.99	0.09
15	Bluefin tuna	4.03	0.102	0.688	3.12	0.41	0.22
16	Swordfish	4.08	0.013	0.604	3.98	0.96	0.15