

Sine Saloum thesis candidate taxon mapping

2019 landings excluding discards. Independent simple trophic chain in tonnes C. Descending unrounded PPR then taxon. Unknown last.

37 source groups from thesis Table 6.5, 1991–1992. Candidate assignments remain unadopted. Model SPPR unavailable because group 20 diet identity is unresolved.

512 taxa. Landings 5,421,499.96 t. Simple-chain PPR 244,709,206.37 t C.

Missing inputs display ?. Zero recorded catch contributes zero PPR even with an unavailable coefficient. Detailed source and allocation evidence are on Sources.

Taxon name	TL	Catch (t)	Simple-chain PPR (t C)	Mapped group names and weights	Confidence level	Reason
Marine fishes not identified	3.28	1,190,215.05	25,198,978.07	Sphyraena guanchancho (0.331162%); Scomberomorus tritor (0.540899%); Polydactylus quadrifilis (0.287007%); Galeoides decadactylus (4.00706%); Pseudotolithus elongatus (2.66034%); Pseudotolithus brachygnathus (1.01556%); Elops lacerta (0.61817%); Arius latiscutatus (0.993487%); Pomadasys jubelini (1.92074%); Chloroscombrus chrysurus (1.69997%); Caranx hippos (0.209736%); Trachinotus teraia (0.496744%); Citarichthys stampflii (0.320124%); Cynoglossus senegalensis (0.430511%); Eucinostomus melanopterus (1.0818%); Drepane africana (1.57854%); Ilisha africana (4.70251%); Brachydeuterus auritus (2.07528%); Epinephelus aeneus (2.1084%); Hemiramphus brasiliensis (0.386356%); Pellonula leonensis (2.85904%); Trichiurus lepturus (0.298046%); Monodactylus sebae (0.26493%); Ethmalosa fimbriata (6.95441%); Sardinella maderensis (26.6365%); Liza grandisquamis (4.70251%); Sarotherodon melanotheron (30.0144%); Tilapia guineensis (0.805828%)	Very low	The reported broad bony-fish assemblage may span named and pooled source fish compartments, including both pelagic and bottom-associated taxa. Use all represented bony-fish pools; exclude the dasyatid, invertebrate, producer and detrital pools. Source composition is a proxy, not regional unidentified-catch composition. Source catch allocation rejected because at least one candidate Y is a printed dash (unknown), not zero. Complete Table6.5 biomass proportions are used under a source assemblage/catchability proxy and fixed time/area transfer assumption; model-estimated parenthesized biomass remains labelled estimated. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_REPORTING.
Trachurus trachurus	3.71	359,796.90	20,502,881.57	Caranx hippos (100%)	Very low	The catch label is a carangid, but its inclusion is not documented in the listed Caranx/Decapterus/Hemicaranx/Lichia pool. That pool is the closest represented jack/scad analogue; the dedicated Chloroscombrus and Trachinotus pools are less directly related. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, TAXONOMY, CATCH.
Katsuwonus pelamis	4.43	65,034.56	19,449,198.43	Sphyraena guanchancho (100%)	Very low	The source barracuda pool is a represented predatory water-column fish assemblage. This unlisted pelagic/deep-water fish has no matching source compartment; body form, offshore/deep habitat and trophic function may differ. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_BARRACUDA.
Sardina pilchardus	3.06	1,489,902.95	19,007,082.95	Sardinella maderensis (100%)	Very low	This small pelagic is not listed in the source other-clupeid pool. Sardinella is the closest schooling small-pelagic analogue; different taxonomy and, for deep-water taxa, habitat are explicitly transferred. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_PILCHARD.
Scomber colias	3.91	209,117.99	18,886,387.37	Sphyraena guanchancho (100%)	Very low	The source barracuda pool is a represented predatory water-column fish assemblage. This unlisted pelagic/deep-water fish has no matching source compartment; body form, offshore/deep habitat and trophic function may differ. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_BARRACUDA.

Similiparma lurida	3.50	198.86	6,987.32 Pomadasys jubelini (100%)	Very low	The source Pomadasys/Plectorhinchus pool represents coastal bottom-associated fishes. Provider classification identifies this taxon as Small reef assoc. fish (<30 cm); it is unlisted and transferred only as a weak coastal demersal/reef fish analogue. Deep-water, reef and body-form/trophic differences remain. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_DEMERSAL.
Pontinus kuhlii	4.05	52.89	6,593.21 Pomadasys jubelini (100%)	Very low	The source Pomadasys/Plectorhinchus pool represents coastal bottom-associated fishes. Provider classification identifies this taxon as Medium bathydemersals (30 - 89 cm); it is unlisted and transferred only as a weak coastal demersal/reef fish analogue. Deep-water, reef and body-form/trophic differences remain. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_DEMERSAL.
Gymnothorax polygonius	4.50	18.60	6,535.64 Pomadasys jubelini (100%)	Very low	The source Pomadasys/Plectorhinchus pool represents coastal bottom-associated fishes. Provider classification identifies this taxon as Large demersals (>=90 cm); it is unlisted and transferred only as a weak coastal demersal/reef fish analogue. Deep-water, reef and body-form/trophic differences remain. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_DEMERSAL.
Mora moro	3.75	103.12	6,443.47 Sphyraena guanchancho (100%)	Very low	The source barracuda pool is a represented predatory water-column fish assemblage. This unlisted pelagic/deep-water fish has no matching source compartment; body form, offshore/deep habitat and trophic function may differ. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_BARRACUDA.
Pseudupeneus prayensis	3.24	325.86	6,292.07 Pomadasys jubelini (100%)	Very low	The source Pomadasys/Plectorhinchus pool represents coastal bottom-associated fishes. Provider classification identifies this taxon as Medium demersals (30 - 89 cm); it is unlisted and transferred only as a weak coastal demersal/reef fish analogue. Deep-water, reef and body-form/trophic differences remain. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_DEMERSAL.

Trichiurus	?	0.00	0.00 Trichiurus lepturus (100%)	Very low	The dedicated Trichiurus lepturus compartment is the closest represented elongate predatory/benthopelagic fish analogue. This broader label or other scabbard/snake-mackerel taxon is not source membership; deep habitat and taxonomy may differ. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, TAXONOMY, CATCH.
Tylosurus	?	0.00	0.00 Hemiramphus brasiliensis (100%)	Medium	The source historical Belonidae pool lists the orthographic forms Tylosorus crocodilus and Tylosorus acus rafale alongside halfbeaks/needlefishes. The authoritative Tylosurus genus is an assumed extension of that represented needlefish set; the source spelling and complete genus membership remain uncertain. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, TAXONOMY, CATCH.
Tylosurus crocodilus	?	0.00	0.00 Hemiramphus brasiliensis (100%)	Low	Table3.3 Saloum explicitly lists Tylosorus crocodilus in the historical Belonidae pool represented by Hemiramphus brasiliensis. WoRMS verifies Tylosurus crocodilus as a belonid but does not recognize the printed genus spelling Tylosorus; the matching epithet and related needlefish context support this tentative orthographic bridge. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, TAXONOMY, CATCH.
Uranoscopus scaber	?	0.00	0.00 Pomadasys jubelini (100%)	Very low	The source Pomadasys/Plectorhinchus pool represents coastal bottom-associated fishes. Provider classification identifies this taxon as Medium demersals (30 - 89 cm); it is unlisted and transferred only as a weak coastal demersal/reef fish analogue. Deep-water, reef and body-form/trophic differences remain. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_GROUPS, THESIS_MEMBERS, CATCH, FAO_DEMERSAL.
Veneridae	?	0.00	0.00 Macrobenthos (100%)	High	Thesis section4.2.2.2.b explicitly places bivalves and gastropods in macrobenthos; WoRMS verifies the catch label's class. No within-pool species split is required. One supported or disclosed analogue group; 100% by definition, not an observed composition estimate. Sources: THESIS_MACROBENTHOS, THESIS_GROUPS, TAXONOMY, CATCH.

Sources and applicability

Thesis source membership and proxy weights are distinct from whole LME applicability and diagnostic availability.

Source	What this evidence supports	Open evidence	Record date
Villanueva (2004) thesis Table6.5, PDF138 / printed114	Exact source37 group identities, pooled principal flags, literal B and Y, estimates and dashes. Y is estimated from B and F; no direct geographic caught-mass allocation.	Open THESIS_GROUPS	
Villanueva thesis Table3.3, PDF82–83 / printed58–59	Actual Sine-Saloum pool membership and representative/other marks. Repeated Psettodes and single-species-caption conflicts remain disclosed. Table3.3 is group composition, distinct from the multi-ecosystem species list.	Open THESIS_MEMBERS	
Villanueva thesis section3.3.4.1, PDF77 / printed53	Shrimp principal taxa and Sine-Saloum P.notialis context; model crabs explicitly Portunidae. Other crabs/lobsters require analogues.	Open THESIS_CRUSTACEANS	
Villanueva thesis sections3.3.4.2.a and4.2.2.2.b, PDF78 and87	Macrobenthos explicitly contains bivalves/gastropods, annelids/polychaetes and echinoids; representative Corbula trigona. No cephalopod compartment is documented.	Open THESIS_MACROBENTHOS	
WoRMS authoritative exact-name records, retrieved3 October2026	Raw responses are retained for125 taxon/source names, with123 uniquely resolved records. The exact source spellings Sphyraena guanchancho and Tylosorus crocodilus are unresolved; historical source spellings are not silently replaced.	Open TAXONOMY	2026-10-03
LME_027.xlsx Catch / Catch and Overview / Settings	Exact512 taxon reporting labels, common/functional classifications and2019 landings; classifications support weak habitat analogues, not exhaustive source membership.	Open CATCH	

FAO marine resources statistical tables	Marine fishes nei is Osteichthyes, separated from sharks/rays/chimaeras; reporting scope supports excluding group17 from broad bony-fish candidate sets.	Open FAO_REPORTING	2026-10-03
FAO Sharks of the World species catalogue, Part2	Primary taxonomy and habitat context for unrepresented shark analogues; no shark source membership or observed allocation is implied.	Open FAO_SHARKS	2026-10-03
FAO Sphyraena sphyraena species identification	Barracuda as a predatory pelagic/coastal fish analogue; source pool transfer still includes different species/body forms and offshore/deep habitats.	Open FAO_BARRACUDA	2026-10-03
FAO living marine resources guide: Dasyatis pastinaca	Demersal soft-bottom stingray feeding on benthic fishes/crustaceans/molluscs; supports only the disclosed benthic analogue.	Open FAO_DASYATIS	2026-10-03
FAO Mugil cephalus species profile	Detritus, microalgae and shallow/estuarine feeding context supports the explicitly weak herbivore analogue.	Open FAO_MULLET	2026-10-03
FAO CECAF central Gulf of Guinea demersal/shrimp resources	Source-represented croaker/grunt/threadfin/sole assemblage occurs in estuarine and coastal bottom communities; supports an ecological analogue only, with whole-LME transfer uncertainty.	Open FAO_DEMERSAL	2026-10-03