

Regional model validation record

Field	Entry
Region	Pacific, Eastern Central high seas, HS_077. Regional workbook
Catch source	Sea Around Us regional catch, 1950–2019; reference: 2019 landings. Eastern Central Pacific high-seas reporting unit; historical local download retained. Catch inputs Regional catch
Selected article	Olson, R.J.; Watters, G.M. (2003). A model of the pelagic ecosystem in the eastern tropical Pacific Ocean. IATTC Bulletin 22(3), pp. 135–218. DOI: not recorded. Source PDF Official IATTC bulletin Paper folder
Other known articles	No additional article is recorded for HS_077 in the reviewed local inventory; this is not an exhaustive literature search.
Selected model	077HS_1_Eastern_tropical_Pacific_(1993-1997); final ETP7, 1993–1997. 39 source ecological groups plus one computational import. Pelagic domain: 20°N–20°S, 150°W to the approximate American shelf break. Selected JSON Model folder Accepted loaded state
Other models from the same article	ETP1: initial 34-group, explicitly unbalanced model for the same broad period/domain. ETP2–ETP6: intermediate revisions; separate complete static tables unavailable. ETP5 sensitivities and Ecosim runs are scenarios. ETP7 is the final published parameterization. Source model inventory
Selection rationale	Article: recorded as the selected pilot article; comparative criterion beyond that designation: ?. Model: ? — the recorded diet-consistency concern does not explain choosing among model versions.
Model extraction	Nine material published diet-sum errors are retained in canonical inputs. The accepted historical runtime normalizes ten diets at 0.001 tolerance, including Albacore 1.001; infers missing BA; supplies missing-catch zeros and detritus/import defaults. Source BA and detritus routing remain undocumented. Censored whale EE and fleet catches are distinct from exact zero. No source values were restored or rebalanced. Reconstruction report Source and runtime review
SPPR calculation	[MANUAL — Researcher: calculation choices and reasons] [Excluded groups, scope and relevant values; other adjustments]

Field	Entry
GE diagnostics	No negative SPPR entries across the basal-source columns. rho_living=0.53880579; b=0.00000000; Detritus(39) SPPR=1.68916596.
TE diagnostics	Negative sources: diet_import(40) → Toothed whales(4); Small producers(38) → Toothed whales(4); Large phytoplankton(37) → Toothed whales(4). rho_living=0.62720534; Detritus(39) SPPR=0.03354576.
Geographic fit	Approximate A: 47% of HS_077 is covered by the study. B: 46–48% of the study is covered by HS_077. Boundary estimate
Temporal fit	1993–1997 model averages; catch series 1950–2019. The 2019 reference is 22–26 years later. Fixed ecosystem coefficients and assumed composition weights do not reconstruct annual ecosystem changes.
Other	TE fails PP-budget closure (relative gap 1.75287544) and has near-zero efficiencies in five mammal/bird groups. GE and With Egestion direct OK apply to the accepted transformed runtime; source/ecological admission remains unestablished. Exact matrix/diagnostic review
Open issues and next action	[Researcher: open issues and next actions]
Review and reproducibility	Researcher name: [name] review date: [date]

Taxon mapping and coverage

Reference: 2019 landings; all catch labels, including zero catch and unidentified categories; no group filter.

39 source ecological groups (+1 computational import); 26 catch labels.

Simple-chain PPR totals 226,007,970.44 t C in 2019.

Method: independent simple trophic chain using saved classic taxon coefficients.

[Taxon mapping appendix](#) and [descriptive Sources sheet](#)

Overall confidence	Taxa	Catch percentage	PPR percentage
High	4	53.8378%	53.0702%
Medium	18	44.0842%	46.4103%
Low	0	0.0000%	0.0000%
Very low	4	2.0780%	0.5194%
Unresolved	0	0.0000%	0.0000%

The summary uses the weaker required membership or allocation confidence. The following tables show each component separately.

Group assignment rules

Plain-language rule	Confidence	PPR percentage
Explicit final source membership in the group or stage union.	High	98.7237%
Unknown finfish or shark/ray composition uses represented-pool proxies.	Very low	0.5194%
Verified current and historical names identify the source taxon.	High	0.4922%
Related pelagic billfish or mako extends listed source species.	Medium	0.2071%
Family spans a supported complete set of named and mixed fish groups.	Medium	0.0575%
Southern bluefin uses Pacific bluefin despite habitat/residency mismatch.	Very low	0.0000%

Membership evidence: [final definitions, synonyms and complete candidates](#)

Allocation weight rules

Plain-language rule	Confidence	PPR percentage
One group receives the entire catch taxon; no split uncertainty.	High	53.0702%
Historical source total removals provide assumed composition, transferred across areas, years and catch bases.	Medium	46.8406%
Unreported candidate catches reject the catch proxy; complete model biomass assumes regional composition/catchability.	Medium	0.0892%

Each percentage is the simple-chain PPR associated with taxa using that rule. Weights and assumptions: [source catch, biomass and stage evidence](#)


Unresolved: none. Very low: Elasmobranchii and the two unidentified marine/pelagic finfish labels use assumed represented-pool composition; Thunnus maccoyii uses a Pacific bluefin analogue. These four labels represent 2.0780% of landings and 0.5194% of simple-chain PPR. Exact candidates and limitations are in the appendix.

Geographic evidence screenshots

Target region R and reconstructed study area S share geographic coordinates and scale. Approximate shelf-break sensitivity and the published study-area denominator support the estimates.

Region area


Eastern Central Pacific high seas HS_077



HS_077: repaired Sea Around Us high-seas region 77 boundary; Natural Earth 1:50m land exclusion. [Target boundary](#) | [Image](#)

Study area in the article

Reviewer reconstruction of source PDF7 bounds



Olson and Watters (2003), printed p. 137 / PDF7: 20°N–20°S, 150°W to American shelf break; 32.8 million km². Coast-following reconstruction; the original article supplies no study map. [Source passage](#) | [Bounds page](#) | [Reconstructed image](#)

Original geographic study figure: unavailable. Article Figures 1–4 show sensitivity and dynamic trajectories rather than the study boundary.