🧭 HOWLAND’S LAW: FORMAL FRAMEWORK

A Proposed Dynamical Principle for Asymmetric Three-Body Dark-Sector Composites

Author: James Howland

---

1. Abstract

This framework introduces Howland’s Law, a proposed dynamical principle governing asymmetric three-body composite systems in a hypothetical dark sector. The model assumes two oppositely charged dark constituents bound by a neutral mediator — the neturaltron — which establishes a stable equilibrium separation and prevents full energetic discharge. Under these conditions, internal oscillatory imbalance necessarily generates intrinsic angular momentum. The resulting spin, termed rotational capacitance, provides long-term stability independent of absolute oscillation speed. This document formalizes the definitions, conditions, consequences, boundary cases, and phenomenological predictions associated with the proposed law.

---

2. Motivation

The microscopic structure of dark matter remains unknown. While gravitational behavior is well-characterized, no consensus exists regarding internal composition, binding mechanisms, or dynamical degrees of freedom. Standard cold dark matter models assume structureless, non-interacting particles, yet several open questions persist regarding stability, halo morphology, and small-scale structure.

Howland’s Law explores the possibility that dark matter may consist of composite bound states with internal oscillatory dynamics. If such systems exist, asymmetry in constituent masses or charges may naturally produce intrinsic spin, providing a dynamical explanation for stability and potentially influencing structure formation. This framework formalizes the underlying mechanism and its implications.

---

3. Definitions

3.1 Asymmetric Three-Body System

A composite consisting of:

- one dominant-mass dark charge carrier (“heavy component”),
- one or more lighter dark charge carriers (“light components”),
- a neutral binding mediator (the neturaltron).

Asymmetry requires:

\[

$$
m_{\text{heavy}} \neq m_{\text{light}}
$$

3.2 Neturaltron
A neutral dark-sector mediator responsible for:

- establishing a preferred equilibrium separation r_0,
- generating a restoring potential preventing collapse or dissociation,
- maintaining nonzero residual field tension (“never fully discharged”),
- enabling oscillatory motion of the charged components.

3.3 Oscillation
Periodic motion of the lighter components around r_0, approximated near equilibrium by:

$$
V(r) \approx \frac{1}{2} k (r - r_0)^2
$$

3.4 Rotational Capacitance
Stored rotational potential arising from:

- mass/charge asymmetry,
- off-axis oscillatory motion,
- cumulative angular displacement per cycle.

This quantity represents the system’s intrinsic spin.

---

4. Statement of Howland’s Law

> Howland’s Law:
> In any bound, asymmetric three-body dark-sector system mediated by a neturaltron, internal oscillatory imbalance necessarily generates a nonzero intrinsic angular momentum. The resulting spin is independent of absolute oscillation speed and arises solely from asymmetry and the restoring potential.

---

5. Conditions for the Law

Condition A — Binding Dominance
The neturaltron binding energy must exceed oscillation energy:

$$
E_{\text{bind}} > E_{\text{osc}}
$$

Condition B — Asymmetry
Mass or charge asymmetry must be nonzero:

$$
\

$$
\Delta m = m\text{heavy} - m\text{light} \neq 0
$$

Condition C — Nonzero Initial Separation
The system must originate with finite separation and momentum:

$$
r(t0) \neq 0, \quad \dot{r}(t0) \neq 0
$$

When A, B, and C are satisfied, forced rotational behavior emerges.

---

6. Consequences of the Law

6.1 Forced Spin
Oscillation + asymmetry → torque imbalance → intrinsic angular momentum:

$$
L \neq 0
$$

6.2 Speed-Independent Stability
Stability is governed by binding strength, not oscillation velocity.
The system remains intact across a wide velocity range.

6.3 Persistent Internal Motion
The neturaltron field never fully relaxes; residual tension ensures continuous oscillation.

6.4 Formation Memory
The equilibrium separation and oscillation amplitude encode the system’s formation conditions.

---

7. Boundary Cases

7.1 Perfect Symmetry
If $m\text{heavy} = m\text{light}$, rotational capacitance vanishes:

$$
L \to 0
$$

7.2 Zero Oscillation
If oscillation amplitude is zero:

$$
A = 0 \Rightarrow L = 0
$$

7.3 Binding Failure

If $E_{osc} > E_{bind}$, the system dissociates (extreme environments only).

---

8. Phenomenological Predictions

8.1 Universal Spin Distribution
All such composites exhibit intrinsic spin, with magnitude correlated to mass ratio.

8.2 Halo-Scale Effects
Internal spin may influence:

- halo anisotropies,
- substructure stability,
- small-scale clustering.

8.3 Magnetic Moment Edge Case
If internal charges do not perfectly cancel, a residual magnetic moment may appear under extreme magnetic fields (e.g., magnetars).

8.4 Formation-Dependent Diversity
Different early-universe conditions yield distinct oscillation amplitudes and spin magnitudes.

---

9. Mechanism Summary

The neturaltron establishes a stable separation and provides a restoring potential.

The lighter components oscillate around this separation.
Asymmetry ensures each oscillation is slightly off-axis.
Off-axis oscillations accumulate angular displacement.
Angular displacement produces intrinsic spin.
Spin stores rotational capacitance.
Capacitance stabilizes the composite.

---

10. Conclusion

Howland’s Law provides a minimal dynamical principle for asymmetric dark-sector composites.
It offers a mechanism for intrinsic spin, long-term stability, and potential observational signatures without requiring electromagnetic coupling.
This framework formalizes the structure of the proposal and establishes a foundation for future mathematical development or phenomenological exploration.