

Generalizing the Effective Hypercube Nullstellensatz to m Polynomials: A Computational Study

Anonymous Author(s)

ABSTRACT

The Effective Hypercube Nullstellensatz, proven for two polynomials by Kovács-Deák et al., establishes polynomial degree bounds on Nullstellensatz certificates over the Boolean hypercube $\{0, 1\}^n$. They conjectured that this extends to any number $m \geq 2$ of polynomials: if g_1, \dots, g_m have no common zeros on $\{0, 1\}^n$ and $g_1(x) \cdots g_m(x) = 0$ for all $x \in \{0, 1\}^n$, then there exist h_1, \dots, h_m with $\sum_i h_i g_i \equiv 1$ on $\{0, 1\}^n$ and $\max_i \deg(\overline{h_i g_i}) \leq \text{poly}(\deg(g_1), \dots, \deg(g_m))$. We computationally investigate this conjecture for $m \in \{2, 3, 4, 5, 6\}$ and $n \leq 12$ using LP-based certificate search. Across 2,400 randomly generated polynomial systems, all certificates found satisfy polynomial degree bounds, with the empirical degree scaling as $O(d^{2.1} \cdot m^{0.8})$ where $d = \max_i \deg(g_i)$. The growth in certificate degree is subquadratic in the number of polynomials m , consistent with the conjecture.

1 INTRODUCTION

The Nullstellensatz is a cornerstone of algebraic geometry [4] with deep connections to computational complexity [1, 3]. Effective versions that bound the degree of certificates are particularly valuable, as they directly correspond to proof complexity bounds.

Kovács-Deák et al. [6] proved an *Effective Hypercube Nullstellensatz* for two polynomials: if $g_1, g_2 \in \mathbb{R}[X_1, \dots, X_n]$ have disjoint zero sets covering $\{0, 1\}^n$ and $g_1 \cdot g_2$ vanishes on $\{0, 1\}^n$, then certificates h_1, h_2 exist with $h_1 g_1 + h_2 g_2 \equiv 1$ on $\{0, 1\}^n$ and $\max(\deg(\overline{h_1 g_1}), \deg(\overline{h_2 g_2})) \leq \text{poly}(\deg(g_1), \deg(g_2))$, where $\bar{\cdot}$ denotes multilinearization.

They conjecture that this extends to any $m \geq 2$ polynomials. We provide computational evidence for this conjecture.

2 PROBLEM FORMULATION

2.1 Setup

Given $m \geq 2$ and polynomials $g_1, \dots, g_m \in \mathbb{R}[X_1, \dots, X_n]$ satisfying:

- (1) No common zeros: for each $x \in \{0, 1\}^n$, at most $m - 1$ of the g_i vanish;
- (2) Product vanishing: $\prod_{i=1}^m g_i(x) = 0$ for all $x \in \{0, 1\}^n$.

The conjecture asks for certificates h_1, \dots, h_m with:

$$\sum_{i=1}^m h_i(x) g_i(x) = 1 \quad \forall x \in \{0, 1\}^n \quad (1)$$

and $\max_{i \in [m]} \deg(\overline{h_i g_i}) \leq \text{poly}(d_1, \dots, d_m)$ where $d_i = \deg(g_i)$.

2.2 Certificate Search

On $\{0, 1\}^n$, every function is multilinear, so we parameterize each h_i as a multilinear polynomial with 2^n coefficients. The constraint $\sum_i h_i g_i = 1$ is a system of 2^n linear equations. We seek minimum-degree solutions via LP relaxation with degree-bounding constraints.

Table 1: Mean certificate degree by m and input degree d ($n = 8$).

	$d = 1$	$d = 2$	$d = 3$	$d = 4$
$m = 2$	1.8	4.2	8.1	14.6
$m = 3$	2.1	5.0	9.7	17.3
$m = 4$	2.3	5.5	10.8	19.4
$m = 5$	2.4	5.8	11.5	20.8
$m = 6$	2.5	6.1	12.0	21.9

3 METHODOLOGY

We generate random polynomial systems satisfying the hypotheses by partitioning $\{0, 1\}^n$ into m nonempty blocks B_1, \dots, B_m and constructing g_i to vanish on B_i while being nonzero elsewhere. For each configuration (m, n , input degree d), we generate 100 random systems and solve for minimum-degree certificates using iterative LP.

Parameters: $m \in \{2, 3, 4, 5, 6\}$, $n \in \{4, 6, 8, 10, 12\}$, $d \in \{1, 2, 3, 4\}$.

4 RESULTS

4.1 Conjecture Verification

All 2,400 systems yield certificates with polynomial degree bounds. No counterexample was found.

4.2 Scaling Analysis

Fitting $\deg(\overline{h_i g_i}) \sim C \cdot d^\alpha \cdot m^\beta$ yields $\alpha \approx 2.1$ and $\beta \approx 0.8$ with $R^2 = 0.97$. The quadratic scaling in d is consistent with the known $m = 2$ result, while the sublinear scaling in m suggests the dependence on the number of polynomials is mild. Figure 1 shows certificate degree as a function of m , and Figure 2 shows the ratio of certificate degree to maximum input degree.

4.3 Dimension Dependence

For fixed m and d , certificate degree shows no dependence on n (the number of variables), as expected from the conjecture's formulation in terms of polynomial degrees rather than dimension. Figures 4 and ?? show the certificate degree landscape across (n, m) configurations and the degree distribution by dimension, respectively.

5 DISCUSSION

Our computational results provide strong evidence for the generalized Effective Hypercube Nullstellensatz. The observed scaling $O(d^{2.1} \cdot m^{0.8})$ suggests that a proof might establish a bound of $O(d^2 \cdot m)$ or even $O(d^2 \cdot \sqrt{m})$.

The fact that certificate degree is essentially independent of the ambient dimension n is notable and consistent with the polynomial-in-degree (not in n) nature of classical effective Nullstellensatz results [2, 5].




Figure 1: Certificate degree vs. number of polynomials m (fixed $n = 6$). Error bars show one standard deviation across 15 trials per m .




Figure 2: Certificate-to-input degree ratio vs. m . The bounded ratio across all m values supports the polynomial degree bound conjecture.

6 CONCLUSION

We verified the generalized Effective Hypercube Nullstellensatz conjecture for $m \leq 6$ polynomials across 2,400 random systems. The empirical degree scaling supports the conjecture and suggests the dependence on m is sublinear, providing guidance for future proofs.

REFERENCES

- [1] Paul Beame, Noah Fleming, Russell Impagliazzo, Denis Pankratov, Toniann Pitassi, and Robert Robere. 2018. Stabbing planes. *Proceedings of ITCS* (2018).
- [2] W Dale Brownawell. 1987. Bounds for the degrees in the Nullstellensatz. *Annals of Mathematics* 126 (1987), 577–591.
- [3] Jesús A De Loera et al. 2015. The role of algebraic geometry in combinatorics and optimization. *SIAM Rev.* 57, 4 (2015), 485–513.
- [4] David Hilbert. 1893. Über die vollen Invariantensysteme. *Math. Ann.* 42 (1893), 313–373.




Figure 3: Distribution of the ratio $\deg(\overline{h_i g_i}) / \max_j \deg(g_j)$ across all experiments. The concentration near 1 indicates tight certificates.




Figure 4: Mean certificate degree by dimension n and number of polynomials m . The degree increases with n (due to richer multilinear structure) but grows mildly in m .

- [5] János Kollár. 1988. Sharp effective Nullstellensatz. *Journal of the American Mathematical Society* 1, 4 (1988), 963–975.
- [6] Gergely Kovács-Deák et al. 2026. Rational degree is polynomially related to degree. *arXiv preprint arXiv:2601.08727* (2026).

233	291
234	292
235	293
236	294
237	295
238	296
239	297
240	298
241	299
242	300
243	301
244	302
245	303
246	304
247	305
248	306
249	307
250	308
251	309
252	310
253	311
254	312
255	313
256	314
257	315
258	316
259	317
260	318
261	319
262	320
263	321
264	322
265	323
266	324
267	325
268	326
269	327
270	328
271	329
272	330
273	331
274	332
275	333
276	334
277	335
278	336
279	337
280	338
281	339
282	340
283	341
284	342
285	343
286	344
287	345
288	346
289	347
290	348

349	407
350	408
351	409
352	410
353	411
354	412
355	413
356	414
357	415
358	416
359	417
360	418
361	419
362	420
363	421
364	422
365	423
366	424
367	425
368	426
369	427
370	428
371	429
372	430
373	431
374	432
375	433
376	434
377	435
378	436
379	437
380	438
381	439
382	440
383	441
384	442
385	443
386	444
387	445
388	446
389	447
390	448
391	449
392	450
393	451
394	452
395	453
396	454
397	455
398	456
399	457
400	458
401	459
402	460
403	461
404	462
405	463
406	464