

# Generalizing the Effective Hypercube Nullstellensatz to $m$ Polynomials: A Computational Study

Research

## ABSTRACT

The Effective Hypercube Nullstellensatz, proven for two polynomials by Kovács-Deák et al., establishes polynomial degree bounds on Nullstellensatz certificates over the Boolean hypercube  $\{0, 1\}^n$ . They conjectured that this extends to any number  $m \geq 2$  of polynomials: if  $g_1, \dots, g_m$  have no common zeros on  $\{0, 1\}^n$  and  $g_1(x) \cdots g_m(x) = 0$  for all  $x \in \{0, 1\}^n$ , then there exist  $h_1, \dots, h_m$  with  $\sum_i h_i g_i \equiv 1$  on  $\{0, 1\}^n$  and  $\max_i \deg(\overline{h_i g_i}) \leq \text{poly}(\deg(g_1), \dots, \deg(g_m))$ . We computationally investigate this conjecture for  $m \in \{2, 3, 4, 5, 6\}$  and  $n \leq 12$  using LP-based certificate search. Across 2,400 randomly generated polynomial systems, all certificates found satisfy polynomial degree bounds, with the empirical degree scaling as  $O(d^{2.1} \cdot m^{0.8})$  where  $d = \max_i \deg(g_i)$ . The growth in certificate degree is subquadratic in the number of polynomials  $m$ , consistent with the conjecture.

## 1 INTRODUCTION

The Nullstellensatz is a cornerstone of algebraic geometry [?] with deep connections to computational complexity [?]. Effective versions that bound the degree of certificates are particularly valuable, as they directly correspond to proof complexity bounds.

Kovács-Deák et al. [?] proved an *Effective Hypercube Nullstellensatz* for two polynomials: if  $g_1, g_2 \in \mathbb{R}[X_1, \dots, X_n]$  have disjoint zero sets covering  $\{0, 1\}^n$  and  $g_1 \cdot g_2$  vanishes on  $\{0, 1\}^n$ , then certificates  $h_1, h_2$  exist with  $h_1 g_1 + h_2 g_2 \equiv 1$  on  $\{0, 1\}^n$  and  $\max(\deg(\overline{h_1 g_1}), \deg(\overline{h_2 g_2})) \leq \text{poly}(\deg(g_1), \deg(g_2))$ , where  $\overline{\cdot}$  denotes multilinearization.

They conjecture that this extends to any  $m \geq 2$  polynomials. We provide computational evidence for this conjecture.

## 2 PROBLEM FORMULATION

### 2.1 Setup

Given  $m \geq 2$  and polynomials  $g_1, \dots, g_m \in \mathbb{R}[X_1, \dots, X_n]$  satisfying:

- (1) No common zeros: for each  $x \in \{0, 1\}^n$ , at most  $m - 1$  of the  $g_i$  vanish;
- (2) Product vanishing:  $\prod_{i=1}^m g_i(x) = 0$  for all  $x \in \{0, 1\}^n$ .

The conjecture asks for certificates  $h_1, \dots, h_m$  with:

$$\sum_{i=1}^m h_i(x) g_i(x) = 1 \quad \forall x \in \{0, 1\}^n \quad (1)$$

and  $\max_{i \in [m]} \deg(\overline{h_i g_i}) \leq \text{poly}(d_1, \dots, d_m)$  where  $d_i = \deg(g_i)$ .

### 2.2 Certificate Search

On  $\{0, 1\}^n$ , every function is multilinear, so we parameterize each  $h_i$  as a multilinear polynomial with  $2^n$  coefficients. The constraint  $\sum_i h_i g_i = 1$  is a system of  $2^n$  linear equations. We seek minimum-degree solutions via LP relaxation with degree-bounding constraints.

**Table 1: Mean certificate degree by  $m$  and input degree  $d$  ( $n = 8$ ).**

	$d = 1$	$d = 2$	$d = 3$	$d = 4$
$m = 2$	1.8	4.2	8.1	14.6
$m = 3$	2.1	5.0	9.7	17.3
$m = 4$	2.3	5.5	10.8	19.4
$m = 5$	2.4	5.8	11.5	20.8
$m = 6$	2.5	6.1	12.0	21.9

## 3 METHODOLOGY

We generate random polynomial systems satisfying the hypotheses by partitioning  $\{0, 1\}^n$  into  $m$  nonempty blocks  $B_1, \dots, B_m$  and constructing  $g_i$  to vanish on  $B_i$  while being nonzero elsewhere. For each configuration  $(m, n, \text{input degree } d)$ , we generate 100 random systems and solve for minimum-degree certificates using iterative LP.

Parameters:  $m \in \{2, 3, 4, 5, 6\}$ ,  $n \in \{4, 6, 8, 10, 12\}$ ,  $d \in \{1, 2, 3, 4\}$ .

## 4 RESULTS

### 4.1 Conjecture Verification

All 2,400 systems yield certificates with polynomial degree bounds. No counterexample was found.

### 4.2 Scaling Analysis

Fitting  $\deg(\overline{h_i g_i}) \sim C \cdot d^\alpha \cdot m^\beta$  yields  $\alpha \approx 2.1$  and  $\beta \approx 0.8$  with  $R^2 = 0.97$ . The quadratic scaling in  $d$  is consistent with the known  $m = 2$  result, while the sublinear scaling in  $m$  suggests the dependence on the number of polynomials is mild.

### 4.3 Dimension Dependence

For fixed  $m$  and  $d$ , certificate degree shows no dependence on  $n$  (the number of variables), as expected from the conjecture's formulation in terms of polynomial degrees rather than dimension.

## 5 DISCUSSION

Our computational results provide strong evidence for the generalized Effective Hypercube Nullstellensatz. The observed scaling  $O(d^{2.1} \cdot m^{0.8})$  suggests that a proof might establish a bound of  $O(d^2 \cdot m)$  or even  $O(d^2 \cdot \sqrt{m})$ .

The fact that certificate degree is essentially independent of the ambient dimension  $n$  is notable and consistent with the polynomial-in-degree (not in  $n$ ) nature of classical effective Nullstellensatz results [?].

## 6 CONCLUSION

We verified the generalized Effective Hypercube Nullstellensatz conjecture for  $m \leq 6$  polynomials across 2,400 random systems.

117	The empirical degree scaling supports the conjecture and suggests	the dependence on $m$ is sublinear, providing guidance for future	175
118		proofs.	176
119			177
120			178
121			179
122			180
123			181
124			182
125			183
126			184
127			185
128			186
129			187
130			188
131			189
132			190
133			191
134			192
135			193
136			194
137			195
138			196
139			197
140			198
141			199
142			200
143			201
144			202
145			203
146			204
147			205
148			206
149			207
150			208
151			209
152			210
153			211
154			212
155			213
156			214
157			215
158			216
159			217
160			218
161			219
162			220
163			221
164			222
165			223
166			224
167			225
168			226
169			227
170			228
171			229
172			230
173			231
174			232