

Barrett Reduction

Time limit: 500 ms
Memory limit: 256 MB

The C++ compiler optimizes the division by a constant D by multiplying your number by a constant A and then shifting the result B bits. This process is called [Barrett reduction](#).

In this challenge we'll consider the division of an `unsigned 32 bit integer` by another `unsigned 32 bit integer`.

Given A and B , your task is to find the minimum possible value of D , for which the division is correct.

Note: it's guaranteed that for the given values A and B there exists an integer D such that for every number $0 \leq X < 2^{31}$, $\lfloor \frac{X}{D} \rfloor = \lfloor \frac{X \times A}{2^B} \rfloor$, where $\lfloor X \rfloor$ represents the floor (integer part) of the result.

E.g: $\lfloor 0.4 \rfloor = 0$, $\lfloor 6.8 \rfloor = 6$.

Note that the challenge only uses unsigned integers, so there are no problems regarding using floor on negative numbers.

Standard input

The first line contains two integers A and B .

Standard output

The first line should contain one integer D .

Constraints and notes

- $0 \leq A < 2^{32}$
- $0 \leq B \leq 63$
- $1 \leq D < 2^{31}$

Input	Output	Explanation
3435973837 35	10	Here are some examples of divisions. $\lfloor \frac{5}{10} \rfloor = 0$ $\lfloor \frac{5 \times 3435973837}{2^{35}} \rfloor = \lfloor \frac{17179869185}{2^{35}} \rfloor = 0$ $\lfloor \frac{17}{10} \rfloor = 1$ $\lfloor \frac{17 \times 3435973837}{2^{35}} \rfloor = \lfloor \frac{58411555229}{2^{35}} \rfloor = 1$ $\lfloor \frac{35}{10} \rfloor = 3$ $\lfloor \frac{35 \times 3435973837}{2^{35}} \rfloor = \lfloor \frac{120259084295}{2^{35}} \rfloor = 3$