HDR DATASET NOTE

Metastatic melanoma on immune checkpoint blockade

HDR-AU-0001 | version 1.0.0 | 2026-11-11

DOI	Pending, minted at publication
CREATORS	Imad El Badisy
COVERAGE	Australia
PERIOD	2013 to 2017
UNIT	Patient
LICENCE	ODbL 1.0
LANDING PAGE	https://elbadisyimad.com/hdr/datasets/au-melanoma-checkpoint-gide/

Description

Patients with stage IV melanoma treated with anti-PD-1 alone (pembrolizumab or nivolumab) or combined with ipilimumab, at Melanoma Institute Australia and other centres: 75 patients with treatment, best response, tumour neoantigen counts and overall survival.

A small immunotherapy cohort that links tumour features to survival, useful for teaching survival analysis in oncology.

GOOD FOR

- Teaching survival analysis in immuno-oncology
- Exploring neoantigen counts as prognostic markers

NOT GOOD FOR

- Comparing the two regimens, which were not randomised
- Using best response as a baseline predictor

Origin

ORGANISATION	Melanoma Institute Australia and The University of Sydney
PUBLICATION	*Gide TN et al. (2019). Distinct immune cell populations define response to anti-PD-1 monotherapy and anti-PD-1/anti-CTLA-4 combined therapy. Cancer Cell 35(2): 238-255*
COUNTRY	Australia
ADDRESS	https://www.cbioportal.org/study/summary?id=mel_iatlas_gide_2019
ACCESSED	2026-11-11
REUSE TERMS	Released on cBioPortal (study mel_iatlas_gide_2019) under the ODC Open Database License (ODbL) 1.0, which requires attribution and share-alike.
HOW OBTAINED	Taken from the BioBench R package (dataset melgide), which took them from the cBioPortal study mel_iatlas_gide_2019 (iAtlas harmonised).

Files

File	Content	Rows	Cols	Size
`melanoma_gide.csv`	One row per patient	75	11	7 kB
`au-melanoma-checkpoint-gide.xlsx`	All tables in one workbook, with About and Variables sheets			15 kB

SHA-256 checksums of every file are listed in `manifest.csv` on the landing page.

Variables

melanoma_gide.csv One row per patient

Variable	Type	Description
time	number	Months from start of treatment to death or end of follow-up
status	integer	Status at end of follow-up (1 dead, 0 censored)
age	number	Age (years)
sex	string	Sex (Female, Male)
treatment	string	Regimen (anti-PD-1, ipilimumab + anti-PD-1)
drug	string	Drugs received (Pembrolizumab, Nivolumab, Ipilimumab Pembrolizumab, Ipilimumab Nivolumab)
response	string	Best response by RECIST (Complete Response, Partial Response, Stable Disease, Progressive Disease)
cta_self_neoantigen	integer	Number of cancer-testis antigen and self neoantigens in pre-treatment tumour RNA
splice_neoantigen	integer	Number of splice-variant neoantigens in pre-treatment tumour RNA
fusion_neoantigen	integer	Number of gene-fusion neoantigens in pre-treatment tumour RNA
erv_neoantigen	integer	Number of endogenous retrovirus neoantigens in pre-treatment tumour RNA

Known issues

Read these before analysis.

1. **Demonstration entry.** Taken from the BioBench R package to show how HDR presents a dataset. It has not gone through HDR review and will be removed before launch.
2. **response is measured after treatment starts.** Using it to predict survival introduces immortal time bias.
3. **Small sample** (75 patients, 30 deaths).
4. **Neoantigen counts are missing** for one patient.

How to cite

Imad El Badisy (2026). Metastatic melanoma on immune checkpoint blockade (version 1.0.0) [Data set]. HDR: Health Data for Research. https://elbadisyimad.com/hdr/datasets/au-melanoma-checkpoint-gide/

Please also cite the origin: Melanoma Institute Australia and The University of Sydney, Gide TN et al. (2019). Distinct immune cell populations define response to anti-PD-1 monotherapy and anti-PD-1/anti-CTLA-4 combined therapy. Cancer Cell 35(2): 238-255.

Versions

Version	Date	Changes
1.0.0	2026-11-11	Demonstration release

This note was generated by HDR from the deposit metadata on 2026-11-11. HDR datasets reproduce their origin as faithfully as possible; they are not official statistics. When a value matters for a decision, check it against the origin named above.