

MINIREVIEW

Hyperthermophilic enzymes – stability, activity and implementation strategies for high temperature applications

Larry D. Unsworth^{1,2}, John van der Oost³ and Sotirios Koutsopoulos⁴

1 Department of Chemical and Materials Engineering, University of Alberta, Edmonton, Canada

2 National Research Council – National Institute for Nanotechnology, University of Alberta, Edmonton, Canada

3 Laboratory of Microbiology, Wageningen University, the Netherlands

4 Center for Biomedical Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA

Keywords

adsorption; covalent bonding; encapsulation; genomic and proteomic considerations; hyperthermostable enzymes; ion pairs; protein immobilization; structural features

Correspondence

S. Koutsopoulos, Center for Biomedical Engineering, Massachusetts Institute of Technology, NE47-307, 500 Technology Square, Cambridge, MA 02139-4307, USA
Fax: +1 617 258 5239
Tel: +1 617 324 7612
E-mail: sotiris@mit.edu

Current theories agree that there appears to be no unique feature responsible for the remarkable heat stability properties of hyperthermostable proteins. A concerted action of structural, dynamic and other physicochemical attributes are utilized to ensure the delicate balance between stability and functionality of proteins at high temperatures. We have thoroughly screened the literature for hyperthermostable enzymes with optimal temperatures exceeding 100 °C that can potentially be employed in multiple biotechnological and industrial applications and to substitute traditionally used, high-cost engineered mesophilic/thermophilic enzymes that operate at lower temperatures. Furthermore, we discuss general methods of enzyme immobilization and suggest specific strategies to improve thermal stability, activity and durability of hyperthermophilic enzymes.

(Received 28 February 2007, accepted 11 May 2007)

doi:10.1111/j.1742-4658.2007.05954.x

Introduction

In general, it is agreed that living organisms can be grouped into four main categories as defined by the temperature range that they grow in: psychrophiles, mesophiles, thermophiles and hyperthermophiles [1]. The origin of extremophilic organisms has long been debated. Based on the analysis of 16S and 18S rRNA gene sequence data, it was shown that, in the evolutionary history of the three domains of living organisms, bacterial and archaeal hyperthermophiles are closest to the root of the phylogenetic tree of life [2]. Therefore, it has been postulated that hyperthermophiles actually precede mesophilic microorganisms [3].

Intuitively, this is in agreement with current theories about the environmental conditions on the surface of Earth when life emerged. According to this theory, all biomolecules evolved to be functional and stable at high temperatures, and adapted to low temperature environments. However, another theory suggests that hyperthermophiles arose from mesophiles via adaptation to high temperature environments. This hypothesis is based on the supposition that ancestral RNA could not be stable at elevated temperatures [4,5].

The first hyperthermophilic organisms from the *Sulfolobus* species was discovered in 1972 in hot acidic springs in Yellowstone Park [6]. Subsequently, over 50 hyperthermophiles have been discovered in

Abbreviations

ADH, alcohol dehydrogenase; G-C, guanine-cytosine.

environments of extreme temperatures: near or above 100 °C. Examples of environments that, until recently, were considered as being inhospitable to life include volcanic areas rich in sulfur and 'toxic' metals and hydrothermal vents in the deep sea (approximately 4 km below sea level) of extremely high pressure [7]. Recently discovered hyperthermophiles have been observed to grow at temperatures as high as 121 °C [8]. Interestingly, hyperthermophilic microorganisms do not grow below temperatures of 50 °C and, in some cases, do not grow below 80–90 °C [7]. Yet, they can survive at ambient temperatures, in the same way that we can preserve mesophilic organisms in the fridge for prolonged times. Hyperthermozymes, in particular, are essentially inactive at moderate temperatures and gain activity as temperatures increase [9].

Hyperthermozyme function at elevated temperatures is a unique attribute that may enable their use in a plethora of biotechnological and biocatalytic applications, where the opportunities are relevant to (a) how we might employ hyperthermostable enzymes for applications where extreme temperatures are required and (b) how we can engineer enzymes in general to maintain their functionality over a broad range of temperatures. In this minireview, we aim to highlight some of the unique characteristics of hyperthermophilic proteins, at the genome, transcriptome and proteome level, which allow for functionality at high temperatures. Moreover, strategies will be discussed with respect to optimizing the thermostability and activity of free as well as immobilized enzymes. The end goal is to provide a system that is able to operate under temperatures higher than those currently employed in systems based on mesophilic and thermophilic biocatalysts.

Hyperthermostability: genomic and proteomic considerations

The survival of hyperthermophiles necessitates a cellular machinery that operates at extreme temperatures. Thus, all aspects of the complex biomolecular systems have to be functional at high temperatures (i.e. individual proteins, genetic coding material, transcription/translation systems, etc.). By comparing differences between mesophilic, thermophilic and hyperthermophilic biomolecules, it is anticipated that a clearer understanding of the major factors that allow for enzymatic activity at higher temperatures will be provided.

Genome-transcriptome level considerations

Although thermal denaturation of dsDNA is known to be affected by its nucleotide composition [10,11]

and that an increase in guanine-cytosine (G-C) content could result in an increase in DNA thermostability, it has been shown that no correlation exists between G-C content and the optimal growth temperature (T_{opt}) of bacterial organisms [10]. Others suggest that, when specific families of prokaryotes (i.e. bacteria and archaea) are analyzed, there may be significant increases in G-C content that coincide with an increase in T_{opt} [12]. However, it has also been observed that for some cases, a decrease in the frequency of SSS and SSG codons occurs with an increase in T_{opt} , which obscures the uniform increase in G-C content [13].

Interestingly, at the level of RNA, there is a growing body of work suggesting that a correlation does exist between G-C content and T_{opt} [14]. A survey of the small subunit rRNA sequences from archaeal, bacterial and eukaryotic lineages (mesophiles, thermophiles and hyperthermophiles) revealed that there is a significant correlation of the G-C content of the paired stem regions (Watson-Crick base pairing) of the 16S rRNA genes, with the actual length of the stem, and with their T_{opt} [15].

In spite of attempts to correlate the G-C content of hyperthermophilic genomes with their T_{opt} , it should be noted that experiments performed *in vitro* and statistical genomic analyses may not accurately represent the situation *in vivo*. It is generally accepted that the DNA and RNA of hyperthermophilic microorganisms are also stabilized through a combination of mechanisms, including increased intracellular electrolyte concentrations, binding of positively charged proteins and histones and spatially confined atomic fluctuations due to macromolecular crowding [16,17]. In addition, supercoiling plays an important role in stability of chromosomal DNA; all hyperthermophilic bacteria and archaea have the enzyme reverse gyrase, which affects DNA topology and appears to be essential for growth at extreme temperatures [18].

Proteome level considerations

It is generally acknowledged that, although hyperthermophilic proteins may have similar functions as their mesophilic counterparts, there may be intrinsic differences that allow them to maintain structural stability and activity at elevated temperatures. In general, protein stability at extreme temperatures above 90 °C is a complex issue that has been attributed to many factors: (a) amino acid composition (including a decrease in thermolabile residues such as Asn and Cys); (b) hydrophobic interactions; (c) aromatic interactions, ion pairs and increased salt bridge networks;

(d) oligomerization and intersubunit interactions; (e) packing and reduction of solvent-exposed surface area; (f) metal binding; (g) substrate stabilization; (h) a decrease in number and size of surface loops; and (i) modifications in the α -helix and β -sheet content [19–26].

Apart from the above mentioned intrinsic factors, extrinsic factors also have been demonstrated to contribute to protein stability in the context of a biological cell. This mainly concerns the so-called compatible solutes, a wide range of small stabilizing molecules (including sugar-derivatives such as trehalose, mannosyl-glycerate and di-myo-inositol-phosphate) [27]. Another factor usually forgotten when discussing hyperthermophilic proteins is their stability at intracellular conditions. Protein stability studies are generally conducted in dilute protein solutions *in vitro*. Such studies are likely to provide meaningful results when secreted, extracellular proteins are considered. However, these conditions may not represent the real situation found inside the cell: macromolecular crowding and naturally occurring small molecules such as metabolites and sugars are expected to play a significant role in protein stability [28,29].

Recent work has shown that the denaturation temperature (T_d) of the globular protein, CutA1, from the hyperthermophile *Pyrococcus horikoshii* OT3 approaches 150 °C [30]. Upon comparing the crystal structures of CutA1 from *Escherichia coli*, *Thermus thermophilus* and *P. horikoshii* OT3 (T_{opt} of 37, 75 and 95 °C, respectively), it was observed that there was a drastic increase in the number of intrasubunit ion pairs (1, 12 and 30, respectively) as T_{opt} increased. Moreover, this increase in intrasubunit ion pairs was directly related to the relative decrease in neutral amino acids and a significant increase in polar amino acids (i.e. Asp, Glu, Lys, Arg and Tyr). It is thought that the increased presence of ion pairs confers thermal stability due to the significantly reduced desolvation penalty for ion pair formation at increased temperatures [31].

Work by Szilagy and Zavodszky [32] categorized thermophilic proteins based on the T_{opt} of the microorganism. They compared the crystal structures of proteins from moderate thermophilic microorganisms (T_{opt} = 45–80 °C) and extreme thermophilic microorganisms (T_{opt} \approx 100 °C). It was observed that the number of ion pairs increased with increasing growth temperature, whereas other parameters, such as hydrogen bonds and the polarity of buried surfaces, do not directly correlate with T_{opt} . Furthermore, the authors concluded that proteins from moderate and extreme

thermophilic organisms are stabilized via different mechanisms. However, although these trends are consistent with previous studies, it should be noted that not all proteins from hyperthermophiles are hyperthermostable. There are proteins from hyperthermophilic organisms that denature at temperatures between 70 and 80 °C and, conversely, proteins from thermophilic organisms that exhibit melting temperatures of approximately 100 °C.

Upon comparing citrate synthases from the hyperthermophilic *Pyrococcus furiosus* (T_{opt} = 100 °C), the thermophilic *Thermoplasma acidophilum* (T_{opt} = 55 °C), the mesophilic mammal (pig; T_{opt} = 37 °C), and the psychrophilic bacterium (Antarctic strain DS2-3R; T_{opt} = 4 °C), it was observed that subunit contacts are crucial for enhancing the thermostability of these homodimeric enzymes [33]. Specifically, it was shown, using three site-directed mutants of *P. furiosus* and *T. acidophilum* citrate synthases, that ionic interactions are essential to their thermal stability. Indeed, ionic interactions, including ionic networks, are thought to be crucial among enzymes with activities around 100 °C [33]. Finally, it was also shown that thermostability does not guarantee thermoactivity. This final point is of particular interest because it highlights the delicate balance between thermostability and thermoactivity that must be considered when employing hyperthermozymes for biotechnological and biocatalytic applications.

Protein molecules are not fixed structures, as depicted in crystallographic representations. Rather, they exhibit a dynamic nature as described by their conformational flexibility, which in turn depends on the fluctuations of the protein atoms. Earlier work [9], which was later confirmed for other homologues proteins [34], suggested that the flexibility of a hyperthermostable protein is lower than that of thermophilic and mesophilic proteins at room temperature and increases with temperature, so as to allow for enzymatic activity near 100 °C. It is only upon achieving these high temperatures that sufficient molecular flexibility (via atomic motions) exists to facilitate the necessary conformational changes required for enzymatic activity (e.g. binding, releasing the substrate, etc.) [9].

Opportunities for biotechnological applications

Perhaps the quintessential example of a successful biotechnological application of thermozymes is the use of Taq polymerase, isolated from *Thermus aquaticus* [35], for PCR [36]. The groundbreaking discovery that proteins from hyperthermophilic microorganisms could be

expressed in mesophiles (e.g. *E. coli*) without losing their conformation, heat stability or activity not only lead to further characterization, but also initiated research on applying them to biocatalysis and biotechnology fields. Obviously, the ability of hyperthermostable proteins to be functional at elevated temperatures presents a number of potential opportunities: (a) the enzymatic processing of many natural polymers is significantly limited by their solubility, this barrier could be overcome by increasing the operating temperatures; (b) the viscosity of the medium increases as temperature is raised; (c) diffusion limitations of the reactants and of the products are minimized; (d) favorable thermodynamics (i.e. for endothermic reactions) would result in increased yields when the reaction is performed at high temperatures; (e) the reactions kinetics are faster at high temperatures; (f) enzymatic processing at temperatures near or above 100 °C minimizes the risk of bacterial contamination in food and drug biosynthesis applications; (g) enzyme immobilization may increase heat stability and therefore, improve biocatalyst performance; and (h) protein engineering by rational design and/or random mutagenesis of hyperthermostable enzymes may result in even more thermostable enzymes.

Several enzymes have already replaced many traditional synthetic chemistry processes. To date, the majority of industrially used enzymes are from bacteria and fungi; the result of 'natural evolution'. In some cases, their properties have been improved through: (a) rational design using combinatorial approaches (i.e. 'computational evolution') [37] and (b) random approaches using high-throughput systems (i.e. 'laboratory evolution') [38–40].

The profit motivation for substituting traditional enzymes with hyperthermostable counterparts is enormous, given that the global enzyme market currently exceeds €4 billion per year. The challenge is obvious: rather than investing more effort in generating mutant mesophilic proteins that operate at high temperatures, a more straightforward approach may be to search the existing protein database for the appropriate hyperthermophilic enzyme that normally functions at higher temperatures. Utilizing this approach would obviously avoid the expensive and laborious enzyme engineering process, and revolutionize industrial and biotechnological processes. Obviously, this approach relies on the availability of hyperthermophile orthologs: enzymes with improved stability, and with similar substrate specificity, enantioselectivity and catalytic activity.

Some hyperthermostable proteins, with optimal operation temperatures at or above 100 °C, are summarized in Table 1. Novel hyperthermostable enzymes,

of known or unknown functions, are constantly being discovered, presenting a huge potential for being employed in a number of applications, including starch processing, cellulose degradation and ethanol production, pulp bleaching, leather and textile processing, chemical synthesis, food processing, and the production of detergents, cosmetics, pharmaceuticals, etc. [41–50].




















Thermal stability and enzymatic activity upon immobilization

Successful implementation of hyperthermozymes to many applications depends on their ability to retain activity upon exposure to the harsh conditions required for most enzymatic reactions: non-natural solvents, high temperature and pressure. In addition to these constraints, many processes require the enzyme to be removable from the reaction medium, reusable or at least recyclable, while not contaminating the product stream by its presence. Enzyme immobilization on the surface of a carrier may address many of the issues listed above. Methods commonly employed for this purpose are covalent bonding [51,52], entrapment [53–55] and physical adsorption [56–58]. Adsorption is considered as the dominant mechanism of interaction of a protein with a surface and, in principle, is the initial event that precedes immobilization through covalent bonding or encapsulation. In general, the immobilized enzyme acquires an increased stability at high temperatures [59–61]. However, the key to successfully utilizing enzymes for biotechnological applications is to ensure that upon immobilization the enzyme remains functional.

Protein adsorption mechanisms and events

The interaction of proteins with surfaces often leads to their adsorption (i.e. excess accumulation of protein at the interface compared to the bulk). Physical adsorption is a mild method of immobilization. Protein adsorption events are largely directed by interfacial phenomena in the vicinal region between the surface and the adsorbing species within the bulk contacting medium [57,62,63]. These interfacial phenomena are mainly driven by electrostatic and hydrophobic interactions. Electrostatic interactions can be repulsive or attractive, depending on the net charges of the surface and of the protein. Hydrophobic interactions are thermodynamically favorable because they increase the system entropy by reducing the extent of unfavorable interactions between polar solvent molecules and hydrophobic moieties (i.e. the hydrophobic patches of

Table 1. Hyperthermostable enzymes with commercial interest and optimal activity over 100 °C in aqueous media.

Enzyme	Microorganism	Microorganism <i>T</i> _{opt.} (°C)	Protein <i>T</i> _{opt.} (°C)	Optimal pH	Molecular mass (kDa)	Reference
α -Amylase (α -glucosidic bonds)	  <i>Pyrococcus furiosus</i>	100	106	6.5–7.5	129 (α_2)	[83]
	 <i>Pyrococcus furiosus</i>	100	100	4.5	54	[84]
	 <i>Pyrococcus woesei</i>	100	100	5.5	68	[85]
	<i>Staphylothermus marinus</i>	90	100	5.0	–	[86]
	<i>Methanococcus jannaschii</i>	85	120	5.0–8.0	–	[87]
Pullulanase type II (α -1,6 glycosidic bonds)	<i>Pyrococcus woesei</i>	100	100	6.0	90	[88]
	<i>Pyrococcus furiosus</i>	100	102	6.0	89	[83]
	 <i>Pyrodictium abyssi</i>	98	105	9.0	–	[89]
Pullulan hydrolase III (α -1,6 and α -1,4 glycosidic bonds)	 <i>Thermococcus aggregans</i>	85	100	6.5	83	[90]
Phospho-glucose/mannose isomerase	<i>Pyrobaculum aerophilum</i>	100	102	7.4	65 (α_2)	[91]
Glucose isomerase	<i>Thermotoga maritima</i>	80	105	6.5–7.5	180 (α_4)	[92]
β -Mannosidase	 <i>Pyrococcus furiosus</i>	100	105	7.4	220 (α_4)	[93]
α -Glucosidase	<i>Thermococcus strain AN1</i>	80	130	–	63	[45]
	 <i>Thermococcus hydrothermalis</i>	80	120	5.5	57	[94]
	 <i>Pyrococcus woesei</i>	100	100	5.0–5.5	90	[95]
	 <i>Sulfolobus solfataricus</i>	88	> 120	4.5	80	[96]
	 <i>Pyrococcus furiosus</i>	100	115	5.0–6.0	135	[97]
β -Glucosidase	 <i>Pyrococcus furiosus</i>	100	105	–	232 (α_4)	[98]
	 <i>Pyrococcus horikoshii</i>	95	> 100	6.0	35	[99]
	 <i>Thermotoga neapolitana</i>	80	103	7.0–7.5	61	[100]
α -Galactosidase	 <i>Pyrococcus furiosus</i>	100	100	10.0	155	[101]
Threonine (alcohol) dehydrogenase	 <i>Pyrococcus furiosus</i>	100	100	6.1–8.8	32	[102]
Alcohol dehydrogenase	 <i>Pyrococcus furiosus</i>	100	100	6.2–6.6	59	[103]
Carboxypeptidase	 <i>Pyrococcus furiosus</i>	100	100	7.0–7.5	330 (α_6)	[104]
Aminopeptidase	 <i>Pyrococcus horikoshii</i>	95	100	6.0–7.0	40	[105]
	<i>Thermococcus strain NA1</i>	80	> 100	8.0	38	[106]
	<i>Pyrococcus furiosus</i>	100	> 100	–	93	[107]
Glukokinase	<i>Pyrococcus furiosus</i>	100	110	–	114	[108]
Sucrose α -glucohydrolase	<i>Desulfurococcus mucosus</i>	88	105	–	52	[109]
Serine protease	<i>Thermoc. kodakaraensis</i> KOD1	95	> 100	7.0	45	[110]
Thiol protease	<i>Pyrococcus furiosus</i>	100	100	6.3	124 (α_6)	[111]
Metalloprotease	<i>Pyrococcus furiosus</i>	100	104	6.0–7.0	30	[112]
β -1,4-endoglucanase	<i>Pyrobaculum aerophilum</i>	100	> 100	6.0	205 (α_4)	[113]
Pyruvate kinase	<i>Aeropyrum pernix</i>	93	> 100	6.1	207 (α_4)	[113]
	<i>Thermotoga maritima</i>	80	> 100	5.9	190 (α_4)	[113]
	<i>Pyrococcus furiosus</i>	100	125	7.4	180 (α_4)	[114]
Methylthioadenosine phosphorylase	<i>Sulfolobus solfataricus</i>	87	120	7.4	160 (α_6)	[115]
	<i>Thermoc. kodakaraensis</i> KOD1	95	> 100	5.0	312 (α_{10})	[116]
Fructose 1,6-biphosphate aldolase	<i>Sulfolobus-solfataricus</i>	87	100	–	133 (α_4)	[117]
Glucokinase	<i>Aeropyrum pernix</i>	93	> 100	6.2	36	[118]
ADP-dependent glucokinase	<i>Pyrococcus furiosus</i>	100	> 100	7.5	98 (α_2)	[119]
	<i>Thermococcus litoralis</i>	85	> 100	7.5	52	[119]
Glucanotransferase	<i>Thermococcus strain B1001</i>	85	110	5.0–5.5	83	[120]
4- α -glucanotransferase	<i>Pyrococcus furiosus</i> KOD1	100	100	6.0–8.0	77	[121]
Esterase	<i>Pyrococcus furiosus</i>	100	100	7.6	–	[122]
Metalloproteinase	<i>Aeropyrum pernix</i> K1	90	100	5.0–9.0	52	[123]
Aminoacylase	<i>Pyrococcus furiosus</i>	100	100	6.5	170 (α_4)	[124]

the protein and the hydrophobic surface of the sorbent).

The difficulty faced when discussing protein adsorption mechanisms arises from the fact that proteins are highly spatially organized, with various substructures

that have differing stabilities, hydrophilicities and charges at given environmental conditions, such as temperature, concentration, ionic strength and pH. Thus, the diverse chemical and physical properties of proteins and surfaces provide multiple interaction

pathways that facilitate adsorption. It is this innate nature of proteins and surfaces that makes it difficult to predict the mechanism of protein adsorption, thus making it difficult to control the process and consistently generate a surface filled with stable and functional enzymes [57].

A common problem associated with the adsorption of enzymes is the conformational changes observed upon adsorption. Such a structural modification may ultimately lower or even diminish the catalytic efficacy of adsorbed enzymes; as discussed below, activation of enzyme activity may occur in rare cases. This excludes any discussion on enzymes that only become active upon adsorption. In general, however, protein immobilization strategies aim to minimize surface-induced conformational changes of adsorbed proteins.

The effect of adsorption on protein structure, thermostability and enzymatic activity was recently highlighted in a series of studies involving hyperthermostable glucanase from *P. furiosus* [60,61,64]. The conformation of the enzyme in the adsorbed state was determined using spectroscopically 'invisible' particles. It was found that thermal stability and enzymatic activity were dependent on the resulting structure of the adsorbed protein and that this structure was affected by the sorbent hydrophilicity. The denaturation temperatures of the free enzyme in solution and adsorbed to hydrophilic or hydrophobic surfaces were 109, 116 and 133 °C, respectively [61]. Compared to solution free enzyme, adsorption to hydrophobic sorbents led to slightly distorted secondary and tertiary structures [65]. In all cases, the specific enzymatic activity of the enzyme did not change upon adsorption.

Several examples of adsorption-induced activation of enzymes exist and the thermostable lipases are of particular interest because they have the potential for being employed in a myriad of biotech applications [66]. In aqueous media, lipases are usually found in a conformation where a 'flap' blocks the active center [67] and only upon adsorption to colloidal drops of oil is this conformation perturbed enough to allow for enzymatic activity [68]. Work with the lipase QL from *Alcaligenes* sp. showed that physical adsorption on a hydrophobic surface led to: (a) a 135% increase in enzymatic activity, relative to the free enzyme; (b) a 20 °C increase of the optimal temperature for enzymatic activity; and (c) surface regeneration [69], unlike immobilization through chemical grafting.

Therefore, when designing an efficient means of introducing hyperthermozymes to the reaction mixture, it is evident that both the enzyme's and the sorbent's physical and chemical properties must be considered. A general observation is that the majority of proteins

tend to adsorb relatively well on hydrophobic surfaces. However, when interacting with hydrophobic surfaces, enzymes generally appear more susceptible to conformational perturbations as compared to adsorption on hydrophilic surfaces [56,57]. Moreover, conditions such as pH and ionic strength can affect the adsorbed amount of the enzyme. For example, it has been observed that changes in pH may lead not only to increased protein adsorption, but also to higher specific activity than the free enzyme [70]. Furthermore, adsorption-induced conformational changes are less when adsorption occurs at pH values near the protein's pI and that this is responsible for an increase in activity [71].

In physical adsorption, proteins become immobilized on the surface of the sorbent through multiple contact points resulting from the interaction between the sorbent and charged and/or hydrophobic amino acid side chains. Depending on the adsorbing conditions, as well as the protein and surface properties, these interactions, which individually are marginally stable, may result in irreversible immobilization of the protein at the interface when considered in total. Also, depending on the solution conditions (e.g. pH, ionic strength, the presence of a detergent), physically adsorbed enzymes may be displaced from the surface of the carrier [72].

Covalent bonding

It is generally accepted that some of the main benefits associated with covalent immobilization include: (a) increased thermal stability; (b) an ability to scale up to reactor applications; (c) ease of interaction with solution compared to encapsulated enzymes; and (d) decreased probability of the enzyme being displaced from the surface and contaminating the reaction solution. Strategies for the covalent immobilization of enzymes have been reviewed elsewhere [51,73]; this minireview rather focuses on correlating protein stability and activity upon bonding, particularly highlighting mild, multipoint attachment techniques [52,74,75].

Optimizing the multipoint covalent immobilization of thermophilic esterases from *Bacillus stearothermophilus* to agarose gels, yielded: (a) 30 000 and 600-fold increases in half-life compared to free and single-point attached enzymes, respectively; (b) retention of 65% of residual activity (cf. soluble) upon bonding; and (c) retention of 70% activity (cf. immobilized) after 1 week of exposure to organic solvents [75]. The case for optimizing the surface-enzyme interaction to retain activity is further highlighted by work conducted on

modified epoxy supports, where it was shown that some surfaces preserved 75–100% of the activity (cf. free enzyme), whereas other combinations lead to full inactivation of the enzyme [74]. Moreover, epoxy modification of the gel surface leads to the precise control of the covalent bonds formed with the enzyme [52].

Despite the various successful cases realized in covalently attaching enzymes to surfaces, the means of attachment can lead to enzyme inactivation. It has been shown that unreacted functional groups can further react with bonded enzymes that are active, resulting in their inactivation after long periods of incubation at high operating temperatures [76]. Thus, a major immobilization criterion involves neutralizing these reactive groups to prevent the surface from adversely affecting the half life of the enzyme.

Encapsulation

Enzyme encapsulation has the potential to provide a microenvironment that increases thermal stability and facilitates enzymatic activity at high temperatures. Although treated separately, encapsulation includes both adsorption and covalent bonding strategies with the difference that, in this case, the enzyme is confined at least on two dimensions by the encapsulating material. This section focuses on correlating protein stability and activity using traditional and novel encapsulation schemes that employ a variety of materials: silica based materials (e.g. sol-gel matrices, mesoporous silica) [28,53,77], aluminosilicates [55], polymers [54,78] and organoclays [79,80].

Sol-gels are commonly used for protein encapsulation. It has been shown that, upon silica entrapment, the mesophilic α -lactalbumin exhibited a 25–32 °C increase in thermal stability and did not fully denature at 95 °C, even after prolonged treatment [53]. However, this same system did not stabilize apomyoglobin [53]. Immobilization of horse heart cytochrome *c* by encapsulation into mesoporous silica led to improved stability and lifetimes of several months; heating to 100 °C for 24 h resulted in a residual activity of 61–74%, compared to the untreated free enzyme [55]. Polyacrylamide gels have also been used as an encapsulating material for various proteins, resulting in an increase in melting temperature [78]. Furthermore, it was observed that coencapsulation of yeast alcohol dehydrogenase (ADH) with a hyperthermophilic chaperone (group II) from *Thermococcus* strain KS-1 resulted in a significant increase in residual activity: ADH-only and ADH-chaperone yielded residual activities of 15% and 78%, respectively, after 5 days [81].

Intercalation of proteins between layered materials such as protein-organoclay lamellar composites may serve as an effective support providing increased protein stability [82]. The intercalation of glucose oxidase into functionalized phyllosilicate clay yielded systems where activity at denaturing pH values (i.e. between 6 and 9) was maintained at 90% of the free enzyme [80]; a trait ascribed mainly to increased electrostatic interactions between enzyme and surface.

Encapsulation provides a platform for protecting enzymes from thermal inactivation during prolonged exposure to elevated temperatures, provided that adequate interactions occur between the surface and the enzyme. The successful implementation of encapsulated hyperthermozymes obviously requires that the matrix materials are also able to withstand high temperatures.

Strategies for enhancing thermal stability and activity of hyperthermozymes

Crucial for the development and optimization of high-temperature biocatalysis systems is the need to gain further understanding of structural differences between hyperthermozymes and their mesophilic and thermophilic homologs, as well as the effect of immobilization on their structural rearrangement and resulting activity at high temperatures.

Through examining proteomic level differences between hyperthermophilic proteins and their thermo/mesophilic counterparts, it is evident that Nature has employed multiple mechanisms to ensure high temperature activity. However, it appears that the resounding message for increasing the thermal stability of proteins revolves around three central tenets: (a) substitute polar for neutral amino acids so as to further increase the number of ion pair interactions; (b) delete surface loops to decrease molecular flexibility; and (c) minimize cavity volumes to increase packing density.

Because the adsorption configuration and conformational features at interfaces cannot yet be accurately predicted for enzymes, it is difficult to design a platform that works for any given enzymatic system and to find remedies to treat decreased activities of adsorbed enzymes. The delicate balance between thermostability and thermoactivity must be maintained when employing hyperthermozymes for biotechnological and biocatalytic applications. However, several studies on a range of enzymes indicate that successful immobilization strategies can lead to increased thermal stability, operation over a wide pH range, protection from non-natural solvents and higher specific

activities over prolonged operational lifetimes. It is important to consider that protein structural and chemical characteristics need to be correlated to the physical chemical properties of the carrier. As a general guideline: (a) hydrophilic surfaces may be preferred over hydrophobic surfaces; (b) electrostatic effects should be reduced by immobilizing at a solution pH near the pI; (c) surface concentration of enzymes should be maximized to inhibit denaturation events; (iv) there is the need to ensure carrier durability at the optimal, hyperthermozyme operating temperature; and (v) multipoint attachment strategies should be utilized, both to prevent protein leaching and to increase heat stability.

The integration of this information, combined with previous strategies used to enhance the thermostability of mesophilic and thermophilic proteins, should provide an efficient route for the development of catalytic systems based on hyperthermozymes. Research efforts should be focused on facilitating the transfer from meso/thermophilic to hyperthermophilic based catalytic systems.

Future focus

In the genomic era, new hyperthermophilic enzymes with novel properties will be discovered via thorough comparative genomic–proteomic analysis combined with high-throughput structural and functional characterization. The genomes of several hyperthermophilic microorganisms have been sequenced, whereas others are forthcoming (<http://www.genomesonline.org/>). Hyperthermophiles are hosts for a high number of genes, many of which encode proteins of unknown function. A wide range of thermostable and biologically novel enzymes for an array of potential applications is expected to become available simply by searching the ever expanding (meta-)genome sequence databases.

The characterization of these novel proteins has great potential for the chemical and pharmaceutical industries ('White Biotechnology'), as they are applied to the synthesis of chemical compounds that are currently difficult to synthesize using traditional synthetic methods. In addition, these natural enzymes will provide the basis for further protein engineering via the described computational and/or laboratory combinatorial approaches, undoubtedly ushering in a new stage of high temperature enzymatics.

References

- 1 Stetter KO (1996) Hyperthermophilic prokaryotes. *FEMS Microbiol Rev* **18**, 149–158.

- 2 Woese CR, Kandler O & Wheelis ML (1990) Towards a natural system of organisms: proposal for the domains archaea, bacteria, and eucarya. *Proc Natl Acad Sci USA* **87**, 4576–4579.
- 3 Woese CR (1998) The universal ancestor. *Proc Natl Acad Sci USA* **95**, 6854–6859.
- 4 Miller SL & Lazcano A (1995) The origin of life – did it occur at high temperatures? *J Mol Evol* **41**, 689–692.
- 5 Forterre P (1996) A hot topic: the origin of hyperthermophiles. *Cell* **85**, 789–792.
- 6 Brock TD, Brock KM, Belly RT & Weiss RL (1972) *Sulfolobus*: a new genus of sulfur-oxidizing bacteria living in low pH and high temperature. *Arch Mikrobiol* **84**, 54–68.
- 7 Stetter KO (2006) History of discovery of the first hyperthermophiles. *Extremophiles* **10**, 357–362.
- 8 Kashefi K & Lovley DR (2003) Extending the upper temperature limit for life. *Science* **301**, 934.
- 9 Koutsopoulos S, van der Oost J & Norde W (2005) Temperature dependant structural and functional features of a hyperthermostable enzyme using elastic neutron scattering. *Proteins* **61**, 377–384.
- 10 Hickey DA & Singer GAC (2004) Genomic and proteomic adaptations to growth at high temperature. *Genome Biol* **5**, 117–121.
- 11 Russell AP & Holleman DS (1974) The thermal denaturation of DNA: average length and composition of denatured areas. *Nucleic Acids Res* **1**, 959–978.
- 12 Musto H, Naya H, Zavala A, Romero H, Alvarez-Valin F & Bernardi G (2004) Correlations between genomic GC levels and optimal growth temperatures in prokaryotes. *FEBS Lett* **573**, 73–77.
- 13 Basak S & Ghosh TC (2005) On the origin of genomic adaptation at high temperature for prokaryotic organisms. *Biochem Biophys Res Commun* **330**, 629–632.
- 14 Galtier N & Lobry JR (1997) Relationships between genomic GC content, RNA secondary structures and optimal growth temperature in prokaryotes. *J Mol Evol* **44**, 632–636.
- 15 Wang H-C, Xia X & Hickey D (2006) Thermal adaptations of the small subunit ribosomal RNA gene: a comparative study. *J Mol Evol* **63**, 120–125.
- 16 Grogan DW (1998) Hyperthermophiles and the problem of DNA instability. *Mol Microbiol* **28**, 1043–1049.
- 17 Daniel RM & Cowan DA (2000) Biomolecular stability and life at high temperatures. *Cell Mol Life Sci* **57**, 250–264.
- 18 Atomi H, Matsumi R & Imanaka T (2004) Reverse gyrase is not a prerequisite for hyperthermophilic life. *J Bacteriol* **186**, 4829–4833.
- 19 Vielle C & Zeikus GJ (2001) Hyperthermophilic enzymes: sources, uses and molecular mechanisms for thermostability. *Microbiol Mol Biol R* **65**, 1–43.
- 20 Chan MK, Mukund S, Kletzin A, Adams MW & Rees DC (1995) Structure of a hyperthermophilic

- tungstopterin enzyme, aldehyde ferredoxin oxidoreductase. *Science* **267**, 1463–1469.
- 21 Salminen T, Teplyakov A, Kankare J, Cooperman BS, Lahti R & Goldman A (1996) An unusual route to thermostability disclosed by the comparison of *Thermus thermophilus* and *Escherichia coli* inorganic pyrophosphatases. *Protein Sci* **5**, 1014–1025.
- 22 Lee D-W, Hong Y-H, Choe E-A, Lee S-J, Kim S-B, Lee H-S, Oh J-W, Shin H-H & Pyun Y-R (2005) A thermodynamic study of mesophilic, thermophilic and hyperthermophilic 1-arabinose isomerases: The effects of divalent metal ions on protein stability at elevated temperatures. *FEBS Lett* **579**, 1261–1266.
- 23 Britton KL, Baker PJ, Borges KM, Engel PC, Pasquo A, Rice DW, Robb FT, Scandurra R, Stillman TJ & Yip KSP (1995) Insights into thermal stability from a comparison of the glutamate-dehydrogenases from *Pyrococcus furiosus* and *Thermococcus litoralis*. *Eur J Biochem* **229**, 688–695.
- 24 Tanaka Y, Tsumoto K, Yasutake Y, Umetsu M, Yao M, Fukada H, Tanaka I & Kumagai I (2004) How oligomerization contributes to the thermostability of an Archaeon protein: protein L-isoaspartyl-O-methyltransferase from *Sulfolobus tokodaii*. *J Biol Chem* **279**, 32957–32967.
- 25 Karlstrom M, Steen IH, Madern D, Fedoy A-E, Birke-land N-K & Ladenstein R (2006) The crystal structure of a hyperthermostable subfamily II isocitrate dehydrogenase from *Thermotoga maritima*. *FEBS J* **273**, 2851–2868.
- 26 Russell RJM, Ferguson JMC, Haugh DW, Danson MJ & Taylor GL (1997) The crystal structure of citrate synthase from the hyperthermophilic bacterium *Pyrococcus furiosus* at 1.9 Å resolution. *Biochemistry* **36**, 9983–9994.
- 27 Ramos A, Raven NDH, Sharp RJ, Bartolucci S, Rossi M, Cannio R, Lebbink J, van der Oost J, deVos WM & Santos H (1997) Stabilization of enzymes against thermal stress and freeze-drying by mannosylglycerate. *Appl Env Microbiol* **63**, 4020–4025.
- 28 Eggers DK & Valentine JS (2001) Crowding and hydration effects on protein conformation: a study with sol-gel encapsulated proteins. *J Mol Biol* **314**, 911–922.
- 29 Ellis RJ (2001) Macromolecular crowding: an important but neglected aspect of the intracellular environment. *Curr Opin Struct Biol* **11**, 114–119.
- 30 Tanaka T, Sawano M, Ogasahara K, Sakaguchi Y, Bagautdinov B, Katoh E, Kuroishi C, Shinkai A, Yokoyama S & Yutani K (2006) Hyper-thermostability of CutA1 protein, with a denaturation temperature of nearly 150°C. *FEBS Lett* **580**, 4224–4230.
- 31 Elcock AH (1998) The stability of salt bridges at high temperatures: implications for hyperthermophilic proteins. *J Mol Biol* **284**, 489–502.
- 32 Szilagyi A & Zavodszky P (2000) Structural differences between mesophilic, moderately thermophilic and extremely thermophilic protein subunits: results of a comprehensive survey. *Struct Fold Des* **8**, 493–504.
- 33 Arnott MA, Michael RA, Thompson CR, Hough DW & Danson MJ (2000) Thermostability and thermoactivity of citrate synthases from the thermophilic and hyperthermophilic Archaea, *Thermoplasma acidophilum* and *Pyrococcus furiosus*. *J Mol Biol* **304**, 657–668.
- 34 Tehei M, Madern D, Franzetti B & Zaccari G (2005) Neutron scattering reveals the dynamic basis of protein adaptation to extreme temperature. *J Biol Chem* **280**, 40974–40979.
- 35 Brock TD & Freeze H (1969) *Thermus aquaticus* gen. n. & sp. N., a nonsporulating extreme thermophile. *J Bacteriol* **98**, 289–297.
- 36 Saiki RK, Gelfand DH, Stoffel S, Scharf SJ, Higuchi R, Horn GT, Mullis KB & Erlich HA (1988) Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* **239**, 487–491.
- 37 Looger LL, Dwyer MA, Smith JJ & Hellinga HW (2003) Computational design of receptor and sensor proteins with novel functions. *Nature* **423**, 185–190.
- 38 Cherry JR, Lamsa MH, Schneider P, Vind J, Svendsen A, Jones A & Pedersen AH (1999) Directed evolution of a fungal peroxidase. *Nat Biotechnol* **17**, 379–384.
- 39 Lehmann M, Pasamontes L, Lassen SF & Wyss M (2000) The consensus concept for thermostability engineering of proteins. *Biochim Biophys Acta – Prot Struct Mol Enzymol* **1543**, 408–415.
- 40 Crameri A, Whitehorn EA, Tate E & Stemmer WPC (1996) Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat Biotechnol* **14**, 315–319.
- 41 Bouzas TD, Barros-Velazquez J & Villa TG (2006) Industrial applications of hyperthermophilic enzymes: a review. *Prot Pept Lett* **13**, 445–451.
- 42 Bauer MW, Driskill LE & Kelly RM (1998) Glycosyl hydrolases from hyperthermophilic microorganisms. *Curr Opin Biotechnol* **9**, 141–145.
- 43 Haki GD & Rakshit SK (2003) Developments in industrially important thermostable enzymes: a review. *Biores Technol* **89**, 17–34.
- 44 Kim MS, Park JT, Kim YW, Lee HS, Nyawira R, Shin HS, Park CS, Yoo SH, Kim YR, Moon TW *et al.* (2004) Properties of a novel thermostable glucoamylase from the hyperthermophilic archaeon *Sulfolobus solfataricus* in relation to starch processing. *Appl Environ Microbiol* **70**, 3933–3940.
- 45 Piller K, Daniel PM & Petach HH (1996) Properties and stabilization of an extracellular alpha-glucosidase from the extremely thermophilic archaeobacteria *Thermococcus* strain AN1: enzyme activity at 130°C. *Biochim Biophys Acta – Prot Struct Molec Enzymol* **1292**, 197–205.

- 46 Eichler J (2001) Biotechnological uses of archaeal extremozymes. *Biotechnol Adv* **19**, 261–278.
- 47 Cannio R, Di Prizito N, Rossi M & Morana A (2004) A xylan-degrading strain of *Sulfolobus solfataricus*: isolation and characterization of the xylanase activity. *Extremophiles* **8**, 117–124.
- 48 Takagi M, Nishioka M, Kakihara H, Kitabayashi M, Inoue H, Kawakami B, Oka M & Imanaka T (1997) Characterization of DNA polymerase from *Pyrococcus* sp. strain KOD1 and its application to PCR. *Appl Environ Microbiol* **63**, 4504–4510.
- 49 Lundberg KS, Shoemaker DD, Adams MWW, Short JM, Sorge JA & Mathur EJ (1991) High-fidelity amplification using a thermostable DNA-polymerase isolated from *Pyrococcus-furiosus*. *Gene* **108**, 1–6.
- 50 Matsumoto K, Kazuno Y, Higashimura N, Ohshima T & Sakuraba H (2006) *Process for Production of Chiral Hydroxyaldehydes*. Eur Patent Application No. 1 734 129. A1.
- 51 Rao SV, Anderson KW & Bachas LG (1998) Oriented immobilization of proteins. *Microchim Acta* **128**, 127–143.
- 52 Grazu V, Abian O, Mateo C, Batista-Viera F, Fernandez-Lafuente R & Guisan JM (2005) Stabilization of enzymes by multipoint immobilization of thiolated proteins on new epoxy-thiol supports. *Biotechnol Bioengin* **90**, 597–605.
- 53 Eggers DK & Valentine JS (2000) Molecular confinement influences protein structure and enhances thermal protein stability. *Protein Sci* **10**, 250–261.
- 54 Yan M, Ge J, Liu Z & Ouyang P (2006) Encapsulation of single enzyme in nanogel with enhanced biocatalytic activity and stability. *J Am Chem Soc* **128**, 11008–11009.
- 55 Lee C-H, Lang J, Yen C-W, Shih P-C, Lin T-S & Mou C-Y (2005) Enhancing stability and oxidation activity of cytochrome c by immobilization in the nanochannels of mesoporous aluminosilicates. *J Phys Chem B* **109**, 12277–12286.
- 56 Norde W & Zounggrana T (1998) Surface-induced changes in the structure and activity of enzymes physically immobilized at solid/liquid interfaces. *Biotechnol Appl Biochem* **28**, 133–143.
- 57 Norde W (2003) Driving forces for protein adsorption at solid surfaces. In *Biopolymers at Interfaces*, 2nd edn (Malmsten M, ed.). Marcel Dekker, Inc., New York, NY.
- 58 Ladero M, Ruiz G, Pessela BCC, Vian A, Santos A & Garcia-Ochoa F (2006) Thermal and pH inactivation of an immobilized thermostable beta-galactosidase from *Thermus* Sp. Strain T2: comparison to the free enzyme. *Biochem Eng J* **31**, 14–24.
- 59 Simpson HD, Haufler UR & Daniel RM (1991) An extremely thermostable xylanase from the thermophilic eubacterium *Thermotoga*. *Biochem J* **277**, 413–417.
- 60 Koutsopoulos S, van der Oost J & Norde W (2004) Adsorption of an endoglucanase from the hyperthermophilic *Pyrococcus furiosus* on hydrophobic (polystyrene) and hydrophilic (silica) surfaces increases heat stability. *Langmuir* **20**, 6401–6406.
- 61 Koutsopoulos S, van der Oost J & Norde W (2005) Structural features of an hyperthermostable endo-b-1,3-glucanase in solution and adsorbed on 'invisible' particles. *Biophys J* **88**, 467–474.
- 62 Unsworth LD, Sheardown H & Brash JL (2005) Protein resistance of surfaces prepared by sorption of end-thiolated poly(ethylene glycol) to gold: effect of surface chain density. *Langmuir* **21**, 1036–1041.
- 63 van Oss CJ (2003) *Interfacial Forces in Aqueous Media*. Marcel Dekker, Inc, New York, NY.
- 64 Koutsopoulos S, Tjeerdsma A-M, Lieshout JFT, van der Oost J & Norde W (2005) In situ structure and activity studies of an enzyme adsorbed on spectroscopically undetectable particles. *Biomacromolecules* **6**, 1176–1184.
- 65 Koutsopoulos S, van der Oost J & Norde W (2005) Conformational studies of a hyperthermostable enzyme. *FEBS J* **272**, 5484–5496.
- 66 Svendsen A, Clausen IG, Patkar SA, Kim B & Thellersen M (1997) Protein engineering of microbial lipases industrial interest. *Methods Enzymol* **284**, 317–340.
- 67 Brady L, Brzozowski AM, Derwenda ZS, Dodson E, Dodson G, Tolley S, Turkenburg JP, Christiansen L, Huge-Jensen B, Norskov L *et al.* (1990) A serine protease triad forms the catalytic center of a triacylglycerol lipase. *Nature* **343**, 767–770.
- 68 Palomo JM, Munoz G, Fernandex-Lorente G, Mateo C, Fernandex-Lafuente R & Guisan JM (2002) Interfacial adsorption of lipases on a very hydrophobic support (octadecyl-Sepabeads): immobilization, hyperactivation and stabilization of the open form of lipases. *J Mol Catal B Enzym* **19–20**, 279–286.
- 69 Wilson L, Palomo JM, Fernandez-Lorente G, Illanes A, Guisan JM & Fernandez-Lafuente R (2006) Effect of lipase–lipase interactions in the activity, stability and specificity of a lipase from *Alcaligenes* sp. *Enzyme Microb Tech* **39**, 259–264.
- 70 Barrias CC, Martins MCL, Miranda MCS & Barbosa MA (2005) Adsorption of a therapeutic enzyme to self-assembled monolayers: effect of surface chemistry and solution pH on the amount and activity of adsorbed enzyme. *Biomaterials* **26**, 2695–2704.
- 71 Haynes CA & Norde W (1994) Globular proteins at solid/liquid interfaces. *Colloids Surf B Biointerfaces* **2**, 517–566.
- 72 Giacomelli CE & Norde W (2001) The adsorption–desorption cycle. Reversibility of the BSA-silica system. *J Coll Interf Sci* **233**, 234–240.
- 73 Pinheiro HM, Kennedy JF & Cabral JMS (1996) Immobilized enzymes and cells. In *Interfacial Phenomena and*

- Bioproducts*. (Brash JL & Wojciechowski PW, eds), pp. 311–350. Marcel Dekker, New York, NY.
- 74 Mateo C, Fernandez-Lorente G, Abian O, Fernandez-Lafuente R & Guisan JM (2000) Multifunctional epoxy supports: a new tool to improve the covalent immobilization of proteins. the promotion of physical adsorptions of proteins on the supports before their covalent linkage. *Biomacromolecules* **1**, 739–745.
- 75 Fernandez-Lafuente R, Cowan DA & Wood ANP (1995) Hyperstabilization of a thermophilic esterase by multipoint covalent attachment. *Enzyme Microbial Technol* **17**, 366–372.
- 76 Saito T, Yoshida Y, Kawashima K, Lin KH, Inagaki H, Maeda S & Kobayashi T (1997) Influence of aldehyde groups on the thermostability of an immobilized enzyme on an inorganic support. *Mater Sci Eng C – Biomimetic Supramol Syst* **C5**, 149–152.
- 77 Pierre AC (2004) The sol-gel encapsulation of enzymes. *Biocatal Biotransform* **22**, 145–170.
- 78 Bolis D, Politou AS, Kelly G, Pastore A & Temussi PA (2003) Protein stability in nanocages: a novel approach for influencing protein stability by molecular confinement. *J Mol Biol* **336**, 203–212.
- 79 Patil AJ, Muthusamy E & Mann S (2004) Synthesis and self-assembly of organoclay-wrapped biomolecules. *Angew Chem Int Ed* **43**, 4928–4933.
- 80 Patil AJ, Muthusamy E & Mann S (2005) Fabrication of functional protein–organoclay lamellar nanocomposites by biomolecule-induced assembly of exfoliated aminopropyl-functionalized magnesium phyllosilicates. *J Mater Chem* **15**, 3838–3843.
- 81 Kohda J, Kawanishi H, Suehara K-I, Nakano Y & Yano T (2006) Stabilization of free and immobilized enzymes using hyperthermophilic chaperonin. *J Biosci Bioeng* **101**, 131–136.
- 82 Coradin T, Coupe A & Livage J (2003) Intercalation of biomolecules in the MnPS₃ layered phase. *J Mater Chem* **13**, 705–707.
- 83 Brown SH, Costantino HR & Kelly RM (1990) Characterization of amylolytic enzyme-activities associated with the hyperthermophilic archaeobacterium *pyrococcus-furiosus*. *Appl Environ Microbiol* **56**, 1985–1991.
- 84 Jørgensen S, Vorgias CE & Antranikian G (1997) Cloning, sequencing, characterization, and expression of an extracellular alpha-amylase from the hyperthermophilic archaeon *Pyrococcus furiosus* in *Escherichia coli* and *Bacillus subtilis*. *J Biol Chem* **272**, 16335–16342.
- 85 Koch R, Spreinat K, Lemke K & Antranikian G (1991) Purification and properties of a hyperthermoactive α -amylase from the archaeobacterium *Pyrococcus woesei*. *Arch Microbiol* **155**, 572–578.
- 86 Bragger JM, Daniel RM, Coolbear T & Morgan HW (1989) Very stable enzymes from extremely thermophilic archaeobacteria and eubacteria. *Appl Microbiol Biotechnol* **31**, 556–561.
- 87 Kim JW, Flowers LO, Whiteley M & Peeples TL (2001) Biochemical confirmation and characterization of the family-57-like alpha-amylase of *Methanococcus jannaschii*. *Folia Microbiol* **46**, 467–473.
- 88 Rüdiger A, Jørgensen PL & Antranikian G (1995) Isolation and characterization of a heat stable pullulanase from the hyperthermophilic archaeon *Pyrococcus woesei* after cloning and expression of its gene in *Escherichia coli*. *Appl Environ Microbiol* **61**, 567–575.
- 89 Andrade CMMC, Aguiar WB & Antranikian G (2001) Physiological aspects involved in production of xylanolytic enzymes by deep-sea hyperthermophilic archaeon *Pyrodictum abyssi*. *Appl Biochem Biotechnol* **91**, 655–669.
- 90 Niehaus F, Peters A, Groudieva T & Antranikian G (2000) Cloning, expression and biochemical characterization of a unique thermostable pullulan-hydrolysing enzyme from the hyperthermophilic archaeon *Thermococcus aggregans*. *FEMS Microbiol Lett* **190**, 223–239.
- 91 Hansen T, Urbanke C & Schönheit P (2004) Bifunctional phosphoglucose/phosphomannose isomerase from the hyperthermophilic archaeon *Pyrobaculum aerophilum*. *Extremophiles* **8**, 507–512.
- 92 Brown SH, Sjöholm C & Kelly RM (1993) Purification and characterization of a highly thermostable glucose-isomerase produced by the extremely thermophilic eubacterium, *Thermotoga-maritima*. *Biotechnol Bioeng* **41**, 878–886.
- 93 Bauer MW, Bylina EJ, Swanson RV & Kelly RM (1997) Comparison of a β -glucosidase and a β -mannosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Biol Chem* **271**, 23749–23755.
- 94 Galichet A & Belarbi A (1999) Cloning of an α -glucosidase gene from *Thermococcus hydrothermalis* by functional complementation of a *Saccharomyces cerevisiae* mal11 mutant strain. *FEBS Lett* **458**, 188–192.
- 95 L  veque E, Janecek S & Belarbi HB (2000) Thermophilic archaeal amylolytic enzymes. *Enz Microbiol Technol* **26**, 3–14.
- 96 Rolfmeier M, Haseltine C, Bini E, Clark A & Blum P (1998) Molecular characterization of the α -glucosidase gene (malA) from the hyperthermophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **180**, 1287–1295.
- 97 Brown SH & Kelly RM (1993) Characterization of amylolytic enzymes, having both α -1,4 and α -1,6 hydrolytic activity, from the thermophilic archaea *Pyrococcus furiosus* and *Thermococcus litoralis*. *Appl Environ Microbiol* **59**, 2614–2621.
- 98 Kengen SWM, Luesink EJ, Stams AJM & Zehnder AJB (1993) Purification and characterization of an extremely thermostable β -glucosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Eur J Biochem* **213**, 305–312.
- 99 Matsui I, Sakai Y, Matsui E, Kikuchi H, Kawarabayashi Y & Honda K (2000) Novel substrate specificity of

- a membrane-bound beta-glycosidase from the hyperthermophilic archaeon *Pyrococcus horikoshii*. *FEBS Lett* **467**, 195–200.
- 100 Duffaud GD, McCutchen CM, Leduc P, Parker KN & Kelly RM (1997) Purification and characterization of extremely thermostable beta-mannanase, beta-mannosidase, and alpha-galactosidase from the hyperthermophilic eubacterium *Thermotoga neapolitana* 5068. *Appl Environ Microbiol* **63**, 169–177.
- 101 Machielsen R & van der Oost J (2006) Production and characterization of a thermostable L-threonine dehydrogenase from the hyperthermophilic archaeon. *Pyrococcus Furiosus* *FEBS J* **273**, 2722–2729.
- 102 Machielsen R, Uria AR, Kengen SWM & van der Oost J (2006) Production and characterization of a thermostable alcohol dehydrogenase that belongs to the aldoketo reductase superfamily. *Appl Environ Microbiol* **72**, 233–238.
- 103 Cheng TC, Ramakrishnan V & Chan SI (1999) Purification and characterization of cobalt-activated carboxypeptidase from the hyperthermophilic Archaeon *Pyrococcus furiosus*. *Protein Sci* **8**, 2474–2486.
- 104 Mori K & Ishikawa K (2005) New deblocking aminopeptidases from *Pyrococcus horikoshii*. *Biosci Biotech Bioch* **69**, 1854–1860.
- 105 Lee HS, Kim YJ, Bae SS, Jeon JH, Lim JK, Jeong BC, Kang SG & Lee JH (2006) Cloning, expression, and characterization of aminopeptidase P from the hyperthermophilic archaeon *Thermococcus* sp strain NA1. *Appl Environ Microbiol* **72**, 1886–1890.
- 106 Story SV, Shah C, Jenney FE & Adams MWW (2005) Characterization of a novel zinc-containing, lysine-specific aminopeptidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Bacteriol* **187**, 2077–2083.
- 107 Badr HR, Sims KA & Adams MWW (1994) Purification and characterization of a sucrose α -glucohydrolase (invertase) from the hyperthermophilic archaeon *Pyrococcus furiosus*. *System Appl Microbiol* **17**, 1–6.
- 108 Kengen SWM, Tuininga JE, de Bok FAM, Stams AJM & de Vos WM (1995) Purification and characterization of a novel ADP-dependent glucokinase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Biol Chem* **270**, 30453–30457.
- 109 Cowan DA, Smolenski KA, Daniel RM & Morgan HW (1987) An extremely thermostable extracellular proteinase from a strain of the archaeobacterium *Desulfurococcus* growing at 88°C. *Biochem J* **247**, 121–133.
- 110 Morikawa M, Izawa Y, Rashid N, Hoaki T & Imanaka T (1994) Purification and characterization of a thermostable thiol protease from a newly isolated hyperthermophilic *Pyrococcus* sp. *Appl Environ Microbiol* **60**, 4559–4566.
- 111 Halio SB, Bauer MW, Mukund S, Adams MWW & Kelly RM (1997) Purification and characterization of two functional forms of intracellular protease Pfpl from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Appl Environ Microbiol* **63**, 289–295.
- 112 Gueguen Y, Voorhorst WGB, van der Oost J & de Vos WM (1997) Molecular and biochemical characterization of an endo-b-1,3-glucanase of the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Biol Chem* **272**, 31258–31264.
- 113 Johnsen U, Hansen T & Schonheit P (2003) Comparative analysis of pyruvate kinases from the hyperthermophilic archaea *Archaeoglobus fulgidus*, *Aeropyrum pernix*, and *Pyrobaculum aerophilum* and the hyperthermophilic bacterium *Thermotoga maritima* – unusual regulatory properties in hyperthermophilic archaea. *J Biol Chem* **278**, 25417–25427.
- 114 Cacciapuoti G, Bertoldo C, Brio A, Zappia V & Porcelli M (2003) Purification and characterization of 5'-methylthioadenosine phosphorylase from the hyperthermophilic archaeon *Pyrococcus furiosus* – substrate specificity and primary structure. *Extremophiles* **7**, 159–168.
- 115 Cacciapuoti G, Porcelli M, Bertoldo C, Derosa M & Zappia V (1994) Purification and characterization of extremely thermophilic and thermostable 5'-methylthioadenosine phosphorylase from the archaeon *Sulfolobus solfataricus* – purine nucleoside phosphorylase-activity and evidence for intersubunit disulfide bonds. *J Biol Chem* **269**, 24762–24769.
- 116 Imanaka H, Fukui T, Atomi H & Imanaka T (2002) Gene cloning and characterization of fructose-1,6-bisphosphate aldolase from the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1. *J Biosci Bioeng* **94**, 237–243.
- 117 Buchanan CL, Connaris H, Danson MJ, Reeve CD & Hough DW (1999) An extremely thermostable aldolase from *Sulfolobus solfataricus* with specificity for non-phosphorylated substrates. *Biochem J* **343**, 563–570.
- 118 Hansen T, Reichstein B, Schmid R & Schonheit P (2002) The first archaeal ATP-dependent glucokinase, from the hyperthermophilic crenarchaeon *Aeropyrum pernix*, represents a monomeric, extremely thermophilic ROK glucokinase with broad hexose specificity. *J Bacteriol* **184**, 5955–5965.
- 119 Koga S, Yoshioka I, Sakuraba H, Takahashi M, Sakasegawa S, Shimizu S & Ohshima T (2000) Biochemical characterization, cloning, and sequencing of ADP-dependent (AMP-forming) glucokinase from two hyperthermophilic archaea, *Pyrococcus furiosus* and *Thermococcus litoralis*. *J Biochem* **128**, 1079–1085.
- 120 Tachibana Y, Kuramura A, Shirasaka N, Suzuki Y, Yamamoto T, Fujiwara S, Takagi M & Imanaka T (1999) Purification and characterization of an extremely thermostable cyclomaltodextrin glucanotransferase from a newly isolated hyperthermophilic archaeon, a *Thermococcus* sp. *Appl Environ Microbiol* **65**, 1991–1997.

- 121 Tachibana Y, Fujiwara S, Takagi M & Imanaka T (1997) Cloning and expression of the 4- α -glucanotransferase gene from the hyperthermophilic archaeon *Pyrococcus* sp KOD1, and characterization of the enzyme. *J Ferment Bioeng* **83**, 540–548.
- 122 Ikeda M & Clark DS (1998) Molecular cloning of extremely thermostable esterase gene from hyperthermophilic archaeon *Pyrococcus furiosus* in *Escherichia coli*. *Biotechnol Bioeng* **57**, 624–629.
- 123 Sako Y, Croocker PC & Ishida Y (1997) An extremely heat-stable extracellular proteinase (aeropyrolysin) from the hyperthermophilic archaeon *Aeropyrum pernix* K1. *FEBS Lett* **415**, 329–334.
- 124 Story SV, Grunden AM & Adams MWW (2001) Characterization of an aminoacylase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Bacteriol* **183**, 4259–4268.