

Ex situ Gap Analysis

Report generate from the GAMMA application
date: 2025-07-07

Summary of results for Taxus chinensis

The gap analysis was conducted using a total of 202 records. Of these 3 were germplasm records and 199 were reference records.

The relationship between these observation types is recorded by a sampling representativeness score.


A buffer size of 50 was used to generate the results for the ecological and geographic representativeness score.

The average for these three scores is used to calculate a final exsitu conservation score.

Sampling Representativeness Score: 1.49.
Geographic representativeness score: 2.86
Ecological representativeness score: 4.55.
Final conservation score: 2.97.

Definitions of ex situ gap analysis scores are below.

Map



Gap Analysis Scores




Table of gap analysis Data

The following table contains all the records that were used to generate the gap analysis results.

Show 10 entries

Search:

Accession Number	Taxon Name	Current Germplasm Type	Collection Date	source	Locality	Collector	Latitude	Longitude
id00674	Taxus chinensis	G	2011	upload	[REDACTED]	HUMAN_OBSERVATION	[REDACTED]	[REDACTED]
id00673	Taxus chinensis	G	2008	upload	[REDACTED]	HUMAN_OBSERVATION	[REDACTED]	[REDACTED]
id00676	Taxus chinensis	G		upload	[REDACTED]	HUMAN_OBSERVATION	[REDACTED]	[REDACTED]
id04551	Taxus chinensis	H	2018	upload	[REDACTED]	PRESERVED_SPECIMEN	[REDACTED]	[REDACTED]
id02247	Taxus chinensis	H	2017	upload	[REDACTED]	PRESERVED_SPECIMEN	[REDACTED]	[REDACTED]
id02245	Taxus chinensis	H	2013	upload	CN PRESENT	HUMAN_OBSERVATION	[REDACTED]	[REDACTED]
id01948	Taxus chinensis	H	2011	upload	[REDACTED]	PRESERVED_SPECIMEN	[REDACTED]	[REDACTED]
id01879	Taxus chinensis	H	2010	upload	[REDACTED]	PRESERVED_SPECIMEN	[REDACTED]	[REDACTED]
id06187	Taxus chinensis	H	2010	upload	[REDACTED]	PRESERVED_SPECIMEN	[REDACTED]	[REDACTED]
id04790	Taxus chinensis	H	2008	upload	[REDACTED]	PRESERVED_SPECIMEN	[REDACTED]	[REDACTED]

Showing 1 to 10 of 202 entries

Previous12345...[REDACTED]

Definitions of occurrence data categories

Germplasm Records (G) : Occurrences in which a living sample (via plant or seed) is present in an (*ex situ*), conservation system (i.e., botanical garden, seed bank, genebank, etc.).

Reference Records (H) : Occurrences that have a supporting herbarium or other reference record.

Sampling Representativeness Score (SRS)

Ex situ: The Sampling Representativeness Score *ex situ* (SRS *ex situ*) calculates the ratio of germplasm accessions (G) available in *ex situ* repositories to reference/voucher (H) records for each taxon.

Geographic Representativeness Score (GRS)

Ex situ: The Geographic Representativeness Score *ex situ* (GRS *ex situ*) uses a user defined km-radius buffer created around each G collection coordinate point to estimate geographic areas already well collected within the distribution of each taxon, also created using buffers around H reference points. This is calculated as the proportion of the distribution covered by the G buffers.

Ecological Representativeness Score (ERS)

Ex situ: The Ecological Representativeness Score *ex situ* (ERS *ex situ*) calculates the proportion of terrestrial ecoregions represented within the G buffered areas out of the total number of ecoregions occupied by the potential distribution.

Final Conservation Score (FCS)

Ex situ: The Final Conservation Score *ex situ* (FCS *ex situ*) was derived by calculating the average of the three *ex situ* conservation metrics.

Prioritization using FSC

In considering the analysis of multiple species, FSC may be used to aid prioritize species action with Urgent Priority (UP) for further conservation action assigned when FCS < 25, High Priority (HP) assigned when 25 ≤ FCS < 50, Medium Priority (MP) when 50 ≤ FCS < 75, and Low Priority (LP) when FCS ≥75.