

# 1 Abstract

This document is the study guide for Nathaniel Barlow's Complex Variables examination on 13 December 2018. It was written by Katie Volz using materials given in class.

The examination consists of twelve topics, each of which are detailed in this study guide. Each topic includes a bulleted list of required knowledge followed by an example problem as given in class.

In this document and in this course, the imaginary number is referred to as  $i$ .

## 2 Topic 1: Sketching a Curve in the Complex Plane

### 2.1 Required Knowledge

- The imaginary number, referred in this course as  $i$ , is defined as  $i = \sqrt{-1}$ . Thus,  $i^2 = -1$ .
- The domain of complex numbers is defined as any number which can be expressed as  $a + bi$ , where both  $a$  and  $b$  are real.
- For  $z = a + bi$  ( $a \in \mathbb{R}, b \in \mathbb{R}$ ),  $a = \operatorname{Re}[z]$  ( $a$  is the real part of  $z$ ) and  $b = \operatorname{Im}[z]$  ( $b$  is the imaginary part of  $z$ ).

### 2.2 Problem

*Note: This problem uses knowledge from Topic 2, as it is difficult to construct a meaningful problem with just the information given.*

Sketch the graph of  $z \in \mathbb{C}$  in the complex plane, where:  $|z - 2i| = \operatorname{Im}[z]$

### 2.3 Solution

Let  $a = \operatorname{Re}[z], b = \operatorname{Im}[z]$ .

$$|z - 2i| = b \tag{1}$$

$$|a + ib - 2i| = b \tag{2}$$

$$|a + i(b - 2)| = b \tag{3}$$

$$\sqrt{a^2 + (b - 2)^2} = b \tag{4}$$

$$a^2 + (b - 2)^2 = b^2 \tag{5}$$


$$a^2 + b^2 - 4b + 4 = b^2 \tag{6}$$

$$a^2 + 4 = 4b \tag{7}$$

$$4b = a^2 + 4 \tag{8}$$

$$b = \frac{a^2}{4} + 1 \tag{9}$$

This is the equation of a parabola. The graph is below.



## 3 Topic 2: Complex Arithmetic

### 3.1 Required Knowledge

- $|z|$ , or the modulus of  $z$ , is the distance from zero. Thus  $|a + bi| = \sqrt{a^2 + b^2}$ .
- $\arg(z)$  is an angle that the complex number forms with the positive real ray. Thus,  $\arg(z)$  can result in an infinite number of angles (each offset by  $2\pi$ ).
- $\text{Arg}(z)$  is the smallest possible angle that the complex number forms with the positive real ray.
- In the case that a  $z$  is negative and real,  $\text{Arg}(z)$  is considered to be  $\pi$ . Thus, for any  $z \in \mathbb{C}$ ,  $-\pi < \text{Arg}(z) \leq \pi$ .
- Euler's Identity states that for any  $z \in \mathbb{C}$ ,  $z = re^{i\theta}$  where  $\theta = \arg(z)$ ,  $r = |z|$ . This form is called polar form.
- $|e^z| = e^{\text{Re}[z]}$  and  $\arg(e^z) = \text{Im}[z] + 2k\pi (k \in \mathbb{R})$
- Cosine Identities:  $\cos(z) = \text{Re}[e^{iz}] = \frac{e^{iz} + e^{-iz}}{2}$
- Sine Identities:  $\sin(z) = \text{Im}[e^{iz}] = \frac{e^{iz} - e^{-iz}}{2i}$


### 3.2 Problem

Let  $z = 2\pi - 2\pi i$ .

- Plot  $z$  in the complex plane.
- Compute  $|z|$ .
- Write  $z$  in polar form.
- Compute  $e^z$ .
- Compute  $\arg(e^z)$ .
- Evaluate  $\cos(z)$ .

### 3.3 Solution (a)

The graph is included below.



### 3.4 Solution (b)

$$|z| = \sqrt{(\text{Re}[z])^2 + (\text{Im}[z])^2} \quad (10)$$

$$|z| = \sqrt{(2\pi)^2 + (2\pi)^2} \quad (11)$$

$$= \sqrt{4\pi^2 + 4\pi^2} \quad (12)$$

$$= \sqrt{8\pi^2} \quad (13)$$

$$= c\sqrt{2}\pi \quad (14)$$

### 3.5 Solution (c)

Use the unit circle provided on the examination to solve this problem.

$$\arg(z) = \frac{-\pi}{4} \quad (15)$$

### 3.6 Solution (d)

Use the solutions from (b) and (c) along with Euler's Identity.

$$z = re^{i\theta} \quad (16)$$

$$z = |z|e^{i\arg(z)} \quad (17)$$

$$z = 2\sqrt{2}\pi e^{\frac{-i\pi}{4}} \quad (18)$$

As the problem only asked for one solution, we could have used any solution for  $\arg(z)$ .

### 3.7 Solution (e)

$$|e^z| = e^{\text{Re}[z]} = e^{2\pi} \quad (19)$$

### 3.8 Solution (f)

$$\arg(e^z) = \text{Im}[z] + 2k\pi, \quad k \in \mathbb{Z} \quad (20)$$

### 3.9 Solution (g)

$$\cos(z) = \frac{e^{iz} + e^{-iz}}{2} \quad (21)$$

$$= \frac{e^{i(2\pi-2\pi i)} + e^{-i(2\pi-2\pi i)}}{2} \quad (22)$$

$$= \frac{e^{2\pi i} e^{2\pi} + e^{-2\pi i} e^{-2\pi}}{2} \quad (23)$$

$$= \frac{e^{2\pi} + e^{-2\pi}}{2} \quad (24)$$

## 4 Topic 3: Raising a Complex Number to an Integral Power

### 4.1 Required Knowledge

- In order to raise a complex number to an integral power, convert the complex number to polar form and distribute the exponent to both the coefficient and the power on  $e$ .
- For raising to an integral power, any argument for the polar form may be chosen (typically this is the principal argument).
- In order to rationalize the denominator of a complex fraction, multiply the top and bottom of the fraction by the complex conjugate of the denominator ( $a - bi$ )

### 4.2 Problem

Evaluate the following and put the result in the form  $a + bi$ :

$$z = \left(\frac{-i}{1+i}\right)^{13} \quad (25)$$

### 4.3 Solution

$$z = \left(\frac{-i}{1+i} * \frac{1-i}{1-i}\right)^{13} \quad (26)$$

$$= \left(\frac{-1-i}{2}\right)^{13} \quad (27)$$

$$= \left(-\frac{1}{2} - \frac{1}{2}i\right)^{13} \quad (28)$$

$$= \left(2^{-\frac{1}{2}} e^{\frac{-3\pi}{4}i}\right)^{13} \quad (29)$$

$$= 2^{-\frac{13}{2}} e^{\frac{-39\pi}{4}i} \quad (30)$$

$$= 2^{-8} \sqrt{2} e^{\frac{-\pi}{4}i} \quad (31)$$

$$= \frac{\sqrt{2}}{128} \left(\cos\left(\frac{\pi}{4}\right) + i \sin\left(\frac{\pi}{4}\right)\right) \quad (32)$$

$$= \frac{\sqrt{2}}{128} \left( \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) \quad (33)$$

$$= \frac{1}{128} (1 + i) \quad (34)$$

$$= \frac{1}{128} + \frac{1}{128}i \quad (35)$$

## 5 Topic 4: Raising a Complex Number to a Rational Power

### 5.1 Required Knowledge

- In order to raise a complex number to a rational power, follow the same steps as an integral power.
- However, the general form for  $\arg(z)$  must be used, adding  $2k\pi i$  to the principal argument, and solve for any  $k$ .
- The number of roots will be equal to the denominator of the exponent when it is fully reduced (unless the radius is zero, in which case there is only one distinct root).
- In order to find all roots, enumerate  $k$  (starting at any point) until the required number of roots are found.

### 5.2 Problem

Solve for all roots of  $z$  and put the result in the form  $a + bi$ :

$$z^3 = -125 \quad (36)$$

### 5.3 Solution

Let  $k \in \mathbb{Z}$ .

$$z^3 = 125e^{i(\pi + 2k\pi)} \quad (37)$$

$$z = 125^{\frac{1}{3}} e^{i\left(\frac{\pi + 2k}{3}\right)} \quad (38)$$

$$z = 5e^{i\left(\frac{\pi + 2k}{3}\right)} \quad (39)$$

There are three roots, which we will call  $z_0$ ,  $z_1$ ,  $z_2$ .  $z_0$  will be found when  $k = 0$ ,  $z_1$  when  $k = 1$ , and  $z_2$  when  $k = 2$ .

$$z_0 = 5e^{i\frac{\pi}{3}} \quad (40)$$

$$z_0 = 5\left(\cos\left(\frac{\pi}{3}\right) + i\sin\left(\frac{\pi}{3}\right)\right) \quad (41)$$

$$z_0 = 5\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) \quad (42)$$

$$z_0 = \frac{5\sqrt{3}}{2} + \frac{5}{2}i \quad (43)$$

$$z_1 = 5e^{i\pi} \quad (44)$$

$$z_1 = 5(\cos(\pi) + i\sin(\pi)) \quad (45)$$

$$z_1 = 5(-1 + 0i) \quad (46)$$

$$z_1 = -5 \quad (47)$$

$$z_2 = 5e^{i\frac{4\pi}{3}} \quad (48)$$

$$z_2 = 5(\cos(\frac{4\pi}{3}) + i\sin(\frac{4\pi}{3})) \quad (49)$$

$$z_2 = 5(-\frac{\sqrt{3}}{2} + \frac{1}{2}i) \quad (50)$$

$$z_2 = -\frac{5\sqrt{3}}{2} + \frac{5}{2}i \quad (51)$$

$$z \in \{\frac{5\sqrt{3}}{2} + \frac{5}{2}i, -5, -\frac{5\sqrt{3}}{2} + \frac{5}{2}i\} \quad (52)$$

## 6 Topic 5: Differentiability of a Complex Function

*Note: Originally, Topic 5 was "Finding the Limits of a Complex Function", however, this has been removed from the examination.*

### 6.1 Required Knowledge

- The CauchyRiemann equations for a function  $z = u(x, y) + iv(x, y)$  are  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$  and  $\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$
- A complex function is differentiable in the complex plane if and only if the CauchyRiemann equations hold for it.
- If the CauchyRiemann equations hold, the complex derivative of a function  $f(z)$  is  $f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i\frac{\partial u}{\partial y}$ .

### 6.2 Problem

Determine whether the complex derivative of  $f(x, y) = x + ix^2$  exists. If it exists, find the derivative.

### 6.3 Solution

First, solve for the partial derivatives.

$$\frac{\partial u}{\partial x} = 1 \quad (53)$$

$$\frac{\partial v}{\partial x} = 2x \quad (54)$$

$$\frac{\partial u}{\partial y} = 0 \quad (55)$$

$$\frac{\partial v}{\partial y} = 0 \quad (56)$$

Check that the CauchyRiemann equations hold:

$$\frac{\partial u}{\partial x} \stackrel{?}{=} \frac{\partial v}{\partial y} \text{ \& } \frac{\partial v}{\partial x} \stackrel{?}{=} -\frac{\partial u}{\partial y} \quad (57)$$

$$1 \stackrel{?}{=} 0 \text{ \& } 2x \stackrel{?}{=} -0 \quad (58)$$

These equations hold nowhere. Thus, the function is not differential anywhere. However, we can attempt to calculate the derivative using the derivative equations:

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y} \quad (59)$$

$$= 1 + 2xi = 0 \quad (60)$$

The two equations do not agree. This is a consequence of the CauchyRiemann equations not holding.

## 7 Topic 6: Finding roots using complex logarithms

### 7.1 Required Knowledge

- In this course, the real logarithm function is specified as  $\log_e(x)$  to prevent a self-reference when defining the complex logarithm function.
- The complex logarithm function  $\ln(z)$  is defined as  $\log_e(|z|) + i \arg(z)$ .

### 7.2 Problem

Solve for all roots of  $z$ , where:

$$e^{iz} = 3 + 3i \quad (61)$$

### 7.3 Solution

Let  $k \in \mathbb{Z}$

$$\ln(e^{iz}) = \ln(3 + 3i) \quad (62)$$

$$iz = \ln(3 + 3i) \quad (63)$$

$$iz = \log_e(|3 + 3i|) + i \arg(3 + 3i) \quad (64)$$

$$iz = \log_e(3\sqrt{2}) + i\left(\frac{\pi}{4} + 2k\pi\right) \quad (65)$$

$$z = -i \log_e(3\sqrt{2}) + \frac{\pi}{4} + 2k\pi \quad (66)$$

$$z = \frac{\pi}{4} + 2k\pi - i(\log_e 3 + \frac{1}{2} \log_e 2) \quad (67)$$

## 8 Topic 7: Using the parametric circle formula to solve a complex integral

### 8.1 Required Knowledge

- For an integral  $\int_C f(z)dz$ , the integral of  $f(z)$  along the circular arc  $C$ , replace  $z$  with  $z_0 + re^{i\theta}$  ( $z_0$  being the centre), and  $C$  with  $\theta$ , from 0 to the final angle.

## 8.2 Problem

Solve the following integral:

$$\int_C \frac{1}{\operatorname{Re}[z]} dz \quad (68)$$

Where  $C$  is the circular arc centred at the origin with a radius of 2 from  $\theta = 0$  to  $\theta = \frac{\pi}{4}$ .

## 8.3 Solution

Replace  $z$  with its polar form,  $z = z_0 + re^{i\theta}$ . In this case,  $z_0 = 0$  as it is centred at the origin, and  $r = 2$ , thus  $z = 2e^{i\theta}$ , and  $dz = 2ie^{i\theta}d\theta$ .  $\theta$  is the path of integration, from 0 to  $\frac{\pi}{4}$ .

$$\int_0^{\frac{\pi}{4}} \frac{1}{\operatorname{Re}[2e^{i\theta}]} 2ie^{i\theta} d\theta \quad (69)$$

$$\int_0^{\frac{\pi}{4}} \frac{1}{2 \operatorname{Re}[e^{i\theta}]} 2ie^{i\theta} d\theta \quad (70)$$

$$\int_0^{\frac{\pi}{4}} \frac{1}{\operatorname{Re}[e^{i\theta}]} ie^{i\theta} d\theta \quad (71)$$

$$\int_0^{\frac{\pi}{4}} \frac{ie^{i\theta}}{\operatorname{Re}[e^{i\theta}]} d\theta \quad (72)$$

$$\int_0^{\frac{\pi}{4}} \frac{i \cos(\theta) - \sin(\theta)}{\cos(\theta)} d\theta \quad (73)$$

$$\int_0^{\frac{\pi}{4}} (i - \tan(\theta)) d\theta \quad (74)$$

$$[i\theta + \ln(\cos(\theta))]_0^{\frac{\pi}{4}} \quad (75)$$

$$\frac{\pi}{4}i - 0i + \ln(\cos(\frac{\pi}{4})) - \ln(\cos(0)) \quad (76)$$

$$\frac{\pi}{4}i + \ln(2^{-\frac{1}{2}}) - \ln(1) \quad (77)$$

$$\frac{\pi}{4}i - \frac{1}{2} \ln(2) - 0 \quad (78)$$

$$-\frac{1}{2} \ln(2) + \frac{\pi}{4}i \quad (79)$$

## 9 Topic 8: Taylor's Theorem

### 9.1 Required Knowledge

- The Taylor series of a function  $f(z)$  about  $z_0$  is the series  $\sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n$ .
- The radius of convergence of the Taylor series about  $z_0$  is the distance to the nearest singularity.
- Singularities can be found with any of the following terms:

- $\ln(0)$
- $[0]^{\frac{m}{n}}$ , where  $m$  and  $n$  are relatively prime integers and  $n > 1$
- $[0]^{-m}$ , where  $m$  is a positive integer




## 9.2 Problem

Find the radius of convergence of the following function about  $z_0 = 1 + i$ :

$$f(z) = \frac{\ln(z - 3)}{z} \quad (80)$$

## 9.3 Solution

Singularities occur at  $z = 0$  and  $z = 3$ , which are plotted below.



The radius of convergence is the minimum distance to either of these singularities, which are  $r_0$  and  $r_3$ .

$$r_0 = \sqrt{(1 - 0)^2 + (1 - 0)^2} \quad (81)$$

$$r_0 = \sqrt{1 + 1} \quad (82)$$

$$r_0 = \sqrt{2} \quad (83)$$

$$r_3 = \sqrt{(1 - 3)^2 + (1 - 0)^2} \quad (84)$$

$$r_3 = \sqrt{(-2)^2 + 1^2} \quad (85)$$

$$r_3 = \sqrt{4 + 1} \quad (86)$$

$$r_3 = \sqrt{5} \quad (87)$$

The lesser of these two radii is  $\sqrt{2}$ , thus the radius of convergence is  $\sqrt{2}$ .

# 10 Topic 9: Laurent Series and Singularity Classification

## 10.1 Required Knowledge

- The Laurent series of a function is the Taylor series evaluated at a singularity.
- The principal part of the Laurent series is the part where the exponent on  $(z - z_0)$  is negative.
- The analytic part of the Laurent series is the part where the exponent on  $(z - z_0)$  is zero or positive.

- If the principal part is an infinite series, then there is a singularity at  $z_0$  is an essential singularity.
- If the principal part is a finite non-zero series, then there is a pole at  $z_0$ . The order of the pole is the number of non-zero terms in the series.
- If the principal part is zero, then the singularity at  $z_0$  is removable.

## 10.2 Problem

Identify the types of singularities of each of the following:

- $\frac{\cos(z)}{z^4}$
- $\frac{e^{-1/z^2}}{z^2}$
- $\frac{1-e^z}{z}$

## 10.3 Solution

For the first, there is a singularity at  $z = 0$ . Replace  $\cos(z)$  by its Taylor series at  $z_0 = 0$ . The Taylor series for  $\cos(z)$  will be given on the examination:

$$\cos(z) = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \frac{z^6}{6!} \dots \quad (88)$$

$$\frac{\cos(z)}{z^4} = \frac{1}{z^4} - \frac{1}{z^2 2!} + \frac{1}{4!} - \frac{z^2}{6!} \dots \quad (89)$$

The first two terms are the principal part, thus this is a pole of order 2.

For the second, let  $w = \frac{-1}{z^2}$  and perform the Taylor expansion at  $w = 0$ .

$$e^w = 1 + w + \frac{w^2}{2!} + \frac{w^3}{3!} + \frac{w^4}{4!} \dots \quad (90)$$

$$e^{\frac{-1}{z^2}} = 1 + \frac{-1}{z^2} + \frac{(\frac{-1}{z^2})^2}{2!} + \frac{(\frac{-1}{z^2})^3}{3!} + \frac{(\frac{-1}{z^2})^4}{4!} \dots \quad (91)$$

$$\frac{e^{\frac{-1}{z^2}}}{z^2} = \frac{1}{z^2} - \frac{1}{z^4} + \frac{1}{2!z^6} - \frac{1}{3!z^8} + \frac{1}{4!z^{10}} \dots \quad (92)$$

The principal part is an infinite series, thus this is an essential singularity.

For the third, replace  $e^z$  with its Taylor series expanded at  $z_0 = 0$ .

$$e^z = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} \dots \quad (93)$$

$$1 - e^z = z + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} \dots \quad (94)$$

$$\frac{1 - e^z}{z} = 1 + \frac{z}{2!} + \frac{z^2}{3!} + \frac{z^3}{4!} \dots \quad (95)$$

As the principal part is zero, this is a removable singularity.

# 11 Topic 10: Circular Integrals in the Complex Plane

## 11.1 Required Knowledge

- Cauchy's Residue Theorem states that if a function  $f(z)$  is analytic everywhere, then the integral over a closed path is equal to  $2\pi i$  times the sum of residues enclosed by the path.
- The residue of a function  $f(z)$  at  $z_0$  is the term with  $\frac{1}{z}$  in the Laurent series.
- For a simple pole at  $z_0$ ,  $\text{Res}(f, z_0) = \lim_{z \rightarrow z_0} (z - z_0)f(z)$ .
- For a pole of order  $n$  at  $z_0$ ,  $\text{Res}(f, z_0) = \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z - z_0)^n f(z)$ .

## 11.2 Problem

Solve the following integral:

$$\oint_C (z^2 + 3 + \frac{(z^2 + 1)e^z}{(z + i)(z - 1)^3}) dz \quad (96)$$

## 11.3 Solution

Separate into two separate integrals. The left integral is zero, as it is a closed integral with no singularities enclosed. Factor the right side, eliminating a term, and identify the singularities.

$$\oint_C (z^2 + 3) dz + \oint_C \frac{(z^2 + 1)e^z}{(z + i)(z - 1)^3} dz \quad (97)$$

$$\oint_C \frac{(z + i)(z - i)e^z}{(z + i)(z - 1)^3} dz \quad (98)$$

$$\oint_C \frac{(z - i)e^z}{(z - 1)^3} dz \quad (99)$$

There is a pole of order 3 at  $z_0 = 1$ . Thus, the integral equals:

$$\text{Res}\left(\frac{(z - i)e^z}{(z - 1)^3}, z = 1\right) \quad (100)$$

$$\frac{1}{(3 - 1)!} \lim_{z \rightarrow 1} \frac{d^{3-1}}{dz^{3-1}} [(z - 1)^3 \frac{(z - i)e^z}{(z - 1)^3}] \quad (101)$$

$$\frac{1}{2!} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} [(z - i)e^z] \quad (102)$$

$$\frac{1}{2} \lim_{z \rightarrow 1} \frac{d}{dz} [e^z + (z - i)e^z] \quad (103)$$

$$\frac{1}{2} \lim_{z \rightarrow 1} \frac{d}{dz} [(z - i + 1)e^z] \quad (104)$$

$$\frac{1}{2} \lim_{z \rightarrow 1} [e^z + (z - i + 1)e^z] \quad (105)$$

$$\frac{1}{2} \lim_{z \rightarrow 1} [(z - i + 2)e^z] \quad (106)$$

$$\frac{1}{2} [(1 - i + 2)e^1] \quad (107)$$

$$\frac{e(3 - i)}{2} \quad (108)$$

$$\frac{3e}{2} + i\frac{e}{2} \quad (109)$$

## 12 Topic 11: Infinite Integrals on the Real Line

### 12.1 Required Knowledge

- An infinite integral on the real line from  $-\infty$  to  $\infty$  is equal to the contour integral bounded by the real line and in an infinite arc into the positive imaginary plane.
- By Cauchy's Residue Theorem, this integral is equal to  $2\pi i$  times the sum of all residues for singularities enclosed by this contour (where the imaginary part is positive).
- If  $f(z)$  is an even function (that is, for any  $z$ ,  $f(z) = f(-z)$ ),  $\int_{-\infty}^0 f(z)dz = \int_0^{\infty} f(z)dz$ , and  $2 \int_{-\infty}^0 f(z)dz = 2 \int_0^{\infty} f(z)dz = \int_{-\infty}^{\infty} f(z)dz$

### 12.2 Problem

Solve the following integral:

$$\int_{-\infty}^{\infty} \frac{x^2}{(x^2 + 2x + 2)(x^2 + 1)} dx \quad (110)$$

### 12.3 Solution

$$\int_{-\infty}^{\infty} \frac{x^2}{(x^2 + 2x + 2)(x^2 + 1)} dx \quad (111)$$

Let  $C_r$  be the counterclockwise path bounded by the real axis and a semicircle extending into the positive imaginary quadrants with a radius of  $r$ .

$$\lim_{r \rightarrow \infty} \oint_{C_r} \frac{z^2}{(z^2 + 2z + 2)(z^2 + 1)} dz \quad (112)$$

$$\lim_{r \rightarrow \infty} \oint_{C_r} \frac{z^2}{(z - (-1 - i))(z - (-1 + i))(z + i)(z - i)} dz \quad (113)$$

There are simple poles at  $z = -1 - i$ ,  $z = -1 + i$ ,  $z = -i$ , and  $z = i$ . Of these, only  $z = -1 + i$  and  $z = i$ , so these are the only residues we need to consider.

Let  $f(z) = \frac{z^2}{(z - (-1 - i))(z - (-1 + i))(z + i)(z - i)}$

$$\text{Res}[f(z), z = i] = \lim_{z \rightarrow i} \frac{z^2(z - i)}{(z - (-1 - i))(z - (-1 + i))(z + i)(z - i)} \quad (114)$$

$$\text{Res}[f(z), z = i] = \lim_{z \rightarrow i} \frac{z^2}{(z - (-1 - i))(z - (-1 + i))(z + i)} \quad (115)$$

$$\text{Res}[f(z), z = i] = \frac{i^2}{(i - (-1 - i))(i - (-1 + i))(i + i)} \quad (116)$$

$$\text{Res}[f(z), z = i] = \frac{-1}{(i - 1 + i)(i + 1 - i)(2i)} \quad (117)$$

$$\text{Res}[f(z), z = i] = \frac{-1}{(2i + 1)(1)(2i)} \quad (118)$$

$$\text{Res}[f(z), z = i] = \frac{-1}{-4 + 2i} \quad (119)$$

$$\text{Res}[f(z), z = i] = \frac{4 + 2i}{20} \quad (120)$$

$$\text{Res}[f(z), z = i] = \frac{2 + i}{10} \quad (121)$$

$$\text{Res}[f(z), z = i] = \frac{1}{5} + \frac{1}{10}i \quad (122)$$

$$\text{Res}[f(z), z = -1 + i] = \lim_{z \rightarrow -1+i} \frac{z^2(z - (-1 + i))}{(z - (-1 - i))(z - (-1 + i))(z + i)(z - i)} \quad (123)$$

$$\text{Res}[f(z), z = -1 + i] = \lim_{z \rightarrow -1+i} \frac{z^2}{(z - (-1 - i))(z + i)(z - i)} \quad (124)$$

$$\text{Res}[f(z), z = -1 + i] = \frac{(-1 + i)^2}{(-1 + i - (-1 - i))((-1 + i) + i)((-1 + i) - i)} \quad (125)$$

$$\text{Res}[f(z), z = -1 + i] = \frac{-2i}{(2i)(-1)(-1 + 2i)} \quad (126)$$

$$\text{Res}[f(z), z = -1 + i] = \frac{2i}{(2i)(-1 + 2i)} \quad (127)$$

$$\text{Res}[f(z), z = -1 + i] = \frac{1}{-1 + 2i} \quad (128)$$

$$\text{Res}[f(z), z = -1 + i] = \frac{-1 - 2i}{5} \quad (129)$$

$$\text{Res}[f(z), z = -1 + i] = -\frac{1}{5} - \frac{2}{5}i \quad (130)$$

The integral is equal to the sum of the residues times  $2\pi i$ .

$$2\pi \sum \text{Res} = 2\pi i \left( \frac{1}{5} + \frac{1}{10}i - \frac{1}{5} - \frac{2}{5}i \right) \quad (131)$$

$$2\pi \sum \text{Res} = 2\pi i \left( \frac{-3}{10}i \right) \quad (132)$$

$$2\pi \sum \text{Res} = \frac{3\pi}{5} \quad (133)$$

## 13 Topic 12: Fourier Integrals

### 13.1 Required Knowledge

- In infinite integral which satisfies Jordan's Lemma (all integrals in this class) and contains a  $\cos(z)$  term or  $\sin(z)$  term can be solved by replacing the trigonometric function with  $e^{iz}$  and taking the real or imaginary part of the resultant integral.

### 13.2 Problem

Solve the following integral:

$$\int_0^\infty \frac{\cos(3x)}{(x^2 + 1)^2} dx \quad (134)$$

### 13.3 Solution

Because this is an even function, the evaluation of the integral from 0 to  $\infty$  is half the evaluation of the integral from  $-\infty$  to  $\infty$

$$\frac{1}{2} \int_{-\infty}^{\infty} \frac{\cos(3x)}{(x^2 + 1)^2} dx \quad (135)$$

We assume that the integrand satisfies Jordan's Lemma, as we do with all integrands in this course.

$$= \frac{1}{2} \operatorname{Re} \left[ \int_{-\infty}^{\infty} \frac{e^{3ix}}{(x^2 + 1)^2} dx \right] \quad (136)$$

Let  $C_r$  be the counterclockwise path bounded by the real axis and a semicircle extending into the positive imaginary quadrants with a radius of  $r$ .

$$= \frac{1}{2} \operatorname{Re} [2\pi i \int_{C_r} \frac{e^{3iz}}{(z - i)^2(z + i)^2} dz] \quad (137)$$

As an aside, we can show that we can replace  $\operatorname{Re}[wi]$  with  $-\operatorname{Im}[w]$ :

$$\operatorname{Re}[wi] = \operatorname{Re}[i(\operatorname{Re}[w] + i \operatorname{Im}[w])] = \operatorname{Re}[i \operatorname{Re}[w] - \operatorname{Im}[w]] = \operatorname{Re}[-\operatorname{Im}[w]] = -\operatorname{Im}[w] \quad (138)$$

$$= -\pi \operatorname{Im} \left[ \int_{C_r} \frac{e^{3iz}}{(z - i)^2(z + i)^2} dz \right] \quad (139)$$

This function has poles of order 2 at  $z = i$  and  $z = -i$ . However, we only consider  $z = i$ .

$$= -\pi \operatorname{Im} [\operatorname{Res} \left[ \frac{e^{3iz}}{(z - i)^2(z + i)^2}, z = i \right]] \quad (140)$$

$$= -\pi \operatorname{Im} \left[ \lim_{z \rightarrow i} \left[ \frac{1}{(2 - 1)!} \frac{d^{(2-1)}}{dz^{(2-1)}} \left[ \frac{(z - i)^2 e^{3iz}}{(z - i)^2(z + i)^2} \right] \right] \right] \quad (141)$$

$$= -\pi \operatorname{Im} \left[ \lim_{z \rightarrow i} \left[ \frac{1}{(1)!} \frac{d^{(1)}}{dz^{(1)}} \left[ \frac{e^{3iz}}{(z + i)^2} \right] \right] \right] \quad (142)$$

$$= -\pi \operatorname{Im} \left[ \lim_{z \rightarrow i} \left[ \frac{d}{dz} \left[ \frac{e^{3iz}}{(z + i)^2} \right] \right] \right] \quad (143)$$

$$= -\pi \operatorname{Im} \left[ \lim_{z \rightarrow i} \left[ \frac{3ie^{3iz}(z + i)^2 - 2(2 + i)e^{3iz}}{(z + i)^4} \right] \right] \quad (144)$$

$$= -\pi \operatorname{Im} \left[ \frac{3ie^{3i^2}(i + i)^2 - 2(2 + i)e^{3i^2}}{(i + i)^4} \right] \quad (145)$$

$$= -\pi \operatorname{Im} \left[ \frac{3ie^{-3}(i + i)^2 - 2(2 + i)e^{-3}}{(2i)^4} \right] \quad (146)$$

$$= -\pi \operatorname{Im} \left[ \frac{-16ie^{-3}}{16} \right] \quad (147)$$

$$= -\pi \operatorname{Im} \left[ \frac{-i}{e^3} \right] \quad (148)$$

$$= -\pi \frac{-1}{e^3} \quad (149)$$

$$= \frac{\pi}{e^3} \quad (150)$$