

Forum Review

Detection of Reactive Oxygen and Nitrogen Species by EPR Spin Trapping

FREDERICK A. VILLAMENA and JAY L. ZWEIER

ABSTRACT


Electron paramagnetic resonance spin trapping has become an indispensable tool for the specific detection of reactive oxygen free radicals in biological systems. In this review we describe some of the advantages as well as some experimental considerations of this technique and how it can be applied to biological systems to measure oxidative stress. *Antioxid. Redox Signal.* 6, 619–629.

INTRODUCTION

THE HEIGHTENED INTEREST in reactive oxygen (ROS) or nitrogen (RNS) species results from the knowledge accumulated over the past decade regarding their role as critical mediators in various physiological oxidative processes and disease mechanisms (4, 26, 101). Of these species, the oxygen free radicals (OFRs), superoxide and hydroxyl radical, as well as their secondary radicals are of particular biomedical importance. Figure 1 shows how superoxide radical $O_2^{\cdot-}$ production can be mediated by various sources in biological systems, and then transformed to other highly reactive species that lead to cellular injury. Although low concentrations of these species play an important role as modulators of cellular function, signaling, and immune response, high levels of these species can be detrimental to cellular function and viability. Over the last 15 years, it has also been appreciated that the free radical gas nitric oxide (NO) is a critical regulator of a broad range of biological functions. Studies over this time have lead to the understanding of the critical functional role of NO in neurotransmission, immune response, and the regulation of vascular tone. Its ability to modulate $O_2^{\cdot-}$ production has also led to the understanding that NO in high levels is toxic and can result in stroke, hypertension, vascular dysfunction, atherosclerosis, or inactivation of mitochondrial respiratory processes (26). Figure 2 shows the reactivity of NO with various radicals and how this leads to the formation of RNS.

Radicals can be generated from a variety of biochemical and enzymatic pathways. Common pathways of superoxide generation include the enzymes xanthine oxidase, NADPH oxidase, and the mitochondrial electron transport chain (101). These pathways can also result in protein oxidation and radical generation. For example, mitochondrial cytochrome *c* oxidase has been shown to react with H_2O_2 resulting in the formation of protein-centered radicals (8), and organic peroxides in the presence of NADPH or NADH produce carbonyloxyl ($CO_2^{\cdot-}$) and hydroxyl radicals ($\cdot OH$) (25). NO is synthesized by inducible NO synthase (NOS) (or NOS2) from L-arginine (L-Arg) and O_2 . Xia *et al.* (94) demonstrated a shift of inducible NOS enzyme function to $O_2^{\cdot-}$ generation following depletion of cytosolic L-Arg.

Superoxide radical production in cells has been quantitatively measured from polymorphonuclear leukocytes after activation with a variety of stimulators, including phorbol esters or opsonized zymosan (68). Endothelial cells subjected to anoxia and reoxygenation generate $O_2^{\cdot-}$ -derived $\cdot OH$ (102). Oxygen radical generation has also been measured from mitochondria both in isolated form and in cells (27). In organs such as the heart, superoxide ($O_2^{\cdot-}$)-derived $\cdot OH$, $\cdot R$, and $ROO\cdot$ radicals are generated following ischemia and reperfusion (103). More recently, superoxide has been measured in vessels that have shown to be formed by a vascular oxidase and to have an important role in regulating vascular tension (74, 75).



SCHEME 1. Spin trapping mechanisms.

glucamine dithiocarbamate (MGD) or Fe^{2+} -*N*-(dithiocarboxy)sarcosine (DTCS), to form a radical adduct that is detectable by EPR. A covalent bond is formed from the reaction of nitrones and nitroso with reactive radicals, while a coordination bond is formed between NO and Fe^{2+} complexes (see Scheme 1).

Characteristic EPR spectral profiles can be observed for different spin adducts, and therefore transient radicals that are otherwise undetectable under normal conditions can now be observed. Figure 3 shows typical spectra of some of these spin adducts. The ability of these spectra to discern the type of radicals being trapped can provide a wealth of information in terms of mechanism, kinetics, and dose dependence of radical production in biological systems. However, this straight-




FIG. 3. Typical spin-adduct EPR of $\cdot\text{OH}$ (a and e), $\text{O}_2^{\cdot-}$ (b and f), $\cdot\text{SG}$ (c and g) (peaks marked with "x" are dimers), and $\text{CH}_3\cdot\text{CHOH}$ (d and h) adducts of DMPO and DEPMPO in phosphate buffer at pH 7.4 from various radical generating systems.

forward identification of radicals being trapped requires some caution as will be discussed in the subsequent section.

DETECTION OF OXYGEN RADICALS

Classes of spin traps

Several excellent reviews have appeared over the past 2 decades on EPR spin trapping methods of oxygen radical detection (31, 32, 64, 65). Nitrones have been the molecule of choice due to their specificity and ability to quantify transient OFRs. Two classes of spin traps are commonly used: the linear nitrones, PBN and α -(4-pyridyl-1-oxide)-*N*-*tert*-butyl nitron (4-PyOBN); and the pyrroline-based cyclic nitrones, DMPO, 5-diethoxyphosphoryl-5-methyl-1-pyrroline *N*-oxide (DEPMPO), and 5-ethoxycarbonyl-5-methyl-1-pyrroline *N*-oxide (EMPO). Nitroso compounds are also another class of spin traps, but are limited by their instability, their tendency to form dimers in solution and subsequent disproportionation to give EPR-active species, and short-half-lived radical adducts, as well as their high toxicity, particularly in the case of nitrosobenzenes (Scheme 2).

Isotopically labeled spin traps such as those of ^2H and ^{15}N -labeled DMPO (58) can exhibit a significant increase in detection sensitivity due to reduction in the number of hyperfine splittings. Replacement of ^1H with ^2H eliminates the doublet proton splittings, while substitution of ^{14}N with ^{15}N results in doublet rather than triplet splitting. It has also been reported for the spin trap EMPO that the stability of the $\text{O}_2^{\cdot-}$ adduct is improved with ^{15}N -substitution and deuteration (98).

Spin trap toxicity

Cellular toxicity of nitrones such as DMPO and PBN is low, hence making them suitable for application in biological

systems. Cytotoxicity studies on bovine aortic endothelial cells have been performed for some of these nitrones and have shown DMPO to have the least toxicity with a 50% inhibitory concentration of approximately 140 mM compared with PBN and 4-PyOBN with 50% inhibitory concentrations of 9 mM and 5 mM, respectively (29) (see Table 2). Evidence from gross pathology and histopathological examinations showed that the spin traps DMPO (232 mg/100 g by weight) and PBN (100 mg/100 g by weight) injected intraperitoneally gave no cellular damage, while 4-PyOBN was found to be lethal at 100 and 200 mg/100 g by weight (71).


Detection in lipid phase

One of the challenges in spin trapping is the ability to detect radical formation in the lipid phase that would allow trapping of lipid-derived radicals from lipid peroxidation processes. Limitations due to the relatively hydrophilic character of some of the commonly used nitrones can be overcome by derivatization of DMPO (56, 81), DEPMPO (76, 95), or EMPO (77) with bulky or long alkyl chains enabling them to have improved lipophilicity. The partition coefficient K_p , as a measure of spin trap lipophilicity, is defined as the ratio of the spin trap concentration (based on its UV absorbance) in 1-octanol to that in water. Table 1 shows the partition coefficients of the commonly used cyclic and linear nitrones, with PBN being the most lipophilic, giving it an obvious disadvantage of having poor solubility in water. The 1-octyloxy derivative of EMPO has a high K_p value of 42 (77), and the stability of alkoxy adducts is increased compared with other analogues with shorter alkyl chain lengths.


High stability of the superoxide adduct of EMPO was observed compared with its derivatives with higher alkyl chain length. Results show that the 1-octyloxy EMPO derivative is more suitable for detection of lipid alkoxy radicals (77), while DEPMPO is not suitable for trapping applications in the lipid phase because of its strong hydrophilic character (76). The bicyclic nitron Trazon (70) (Scheme 3) gave characteristic EPR spectra for various alkoxy and lipid-derived radicals (78), and spin trapping in a liposomal environment has been carried out using lipid-soluble spin traps such as 2-methyl-nitrosopropane (MNP) (17, 72).

Competitive trapping

Endogenous cellular superoxide radical scavengers such as superoxide dismutase (SOD), ferricytochrome *c*, epinephrine, L-ascorbic acid, hydroxylamine, or pyrogallol can com-



SCHEME 2. Various types of nitrones and nitroso spin traps.



SCHEME 3. Structure of Trazon.

TABLE 1. RADICAL GENERATING SYSTEMS IN AQUEOUS MEDIA

Radical	Generating system	Reference
$\cdot\text{OH}$	$\text{Fe}^{2+}/\text{H}_2\text{O}_2$; $\text{H}_2\text{O}_2/h\nu$	18
$\text{O}_2^{\cdot-}$, $\cdot\text{OOH}$	Hypoxanthine/xanthine oxidase; riboflavin/light	18, 67, 86
$\cdot\text{CH}_3$	$\text{Fe}^{2+}/\text{H}_2\text{O}_2/\text{dimethyl sulfoxide}$	55
$\text{CO}_2^{\cdot-}$	$\text{Fe}^{2+}/\text{H}_2\text{O}_2/\text{NaHCO}_3$	55
$\text{CH}_3\cdot\text{CHCOH}$	$\text{Fe}^{2+}/\text{H}_2\text{O}_2/\text{ethanol}$	59
$\text{RO}\cdot$	$\text{ROH}/\text{Fe}^{3+}$; $\text{ROH}/\text{Pb}(\text{OAc})_4$	12, 95
$\text{GS}\cdot$, $\text{RS}\cdot$	GSSG/UV ; $\text{RSSR}/h\nu$	95
$\text{SO}_3^{\cdot-}$	$\text{SO}_3^{2-}/\text{Cr}^{4+}$; $\text{Fe}^{2+}/\text{H}_2\text{O}_2/\text{SO}_3^{2-}$	45, 55
$\text{H}\cdot$	$(n\text{-Bu})_3\text{SnH}/h\nu$	95
$\cdot\text{Ph}$, $\text{X}\cdot\text{Ph}$	$\text{Ph}(\text{O})\text{COSnCy}_3/h\nu$	95

pete with the nitron spin traps. However, knowledge of the second order rate constants k_s of these scavengers with $\text{O}_2^{\cdot-}$ can help greatly in setting up experimental design, *e.g.*, type or concentration of trap to be employed. Mitsuta *et al.* (52) reported important kinetic data involving reaction of $\text{O}_2^{\cdot-}$ with SODs and SOD-mimic molecules. Data show that SODs have k_s values on the order of $1.6\text{--}3.0 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ at pH 7.8, while reactivity of other SOD-mimics with $\text{O}_2^{\cdot-}$ are relatively lower compared with the SODs, with k_s values ranging only from 0.86 to $43 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, and for hydroxylamine hydrochloride, which has almost the same rate of reaction as DMPO, with k_s of about $1.6 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$. For *in vitro* studies, $10\text{--}50 \text{ mM}$ concentrations of the spin trap are competitive enough and usually give reasonable peak intensity.

Artifacts in spin trapping


Artifacts are commonly encountered in spin trapping experiments. The cyclic nitron DMPO, for example, is known to undergo the Forrester-Hepburn reaction (63), in which a radical adduct is produced from the Fe^{3+} -catalyzed nucleophilic addition of water to nitron and subsequent oxidation to the EPR-active nitroxide according to Scheme 4 (15). While catalase does not affect this adduct formation, the presence of metal chelators exerts inhibition. Formation of the DMPO-OH adduct due to the nucleophilic addition of H_2O in the presence of Fe^{3+} or Cu^{2+} was confirmed using ^{17}O -enriched water (28). Since Fe^{3+} impurities are only present at

micromolar levels under typical experimental conditions, the use of chelators such as diethylenetriaminepentaacetic acid in $100 \mu\text{M}$ concentrations can avoid formation of this artifactual DMPO-OH.


Accurate assignments of spin adducts can be confirmed by independently generating the radical adducts of interest, and computer simulation of the observed EPR spectra (1, 16). Several established procedures have been reported in the literature on how to generate radicals independently and are summarized in Table 1.

Use of scavengers such as SOD and catalase inhibits $\text{O}_2^{\cdot-}$ and $\cdot\text{OH}$ adduct formation, respectively. Secondary radical formation from ethanol, methanol, or dimethyl sulfoxide can be added to confirm the generation of $\cdot\text{OH}$. Formation of C-centered adducts by addition of these secondary radicals, or inhibition of $\cdot\text{OH}$ adduct formation by catalase, can rule out the possibility that the $\cdot\text{OH}$ adduct formation resulted from the decomposition of the $\text{O}_2^{\cdot-}$ adduct (6) *via* a mechanism (3, 65) shown in Scheme 5, or by some nucleophilic addition reaction.

The formation of EPR-detectable nitroxides during spin trapping experiments may not necessarily result from the direct addition of radical to the spin trap. For example, polyunsaturated fatty acid (PUFA) peroxy radicals give rise to PUFA-derived alkoxyl radical adducts rather than primary PUFA peroxy radical adducts, and this initially resulted in



SCHEME 4. Forrester-Hepburn reaction for the formation of hydroxyl adduct via a non-spin trapping mechanism.



SCHEME 5. Hypothetical mechanism for the unimolecular decomposition of the $\text{O}_2^{\cdot-}/\cdot\text{OOH}$ adduct of DMPO.

some confusion regarding the spectral assignments of these adducts in the literature (12). Because of the instability of the primary peroxy radical, conversion to its corresponding alkoxy radical adduct occurs. In the case of DEPMPO, the EPR signals of the $\cdot\text{OH}$ -DEPMPO adduct can also lead to erroneous interpretation, especially in the presence of high-valency metal ions such as Fe^{3+} , Cu^{2+} , or Mn^{4+} . These metal ions can oxidize DEPMPO to form the radical cation species DEPMPO^+ , hence making this species susceptible to hydration to form the $\cdot\text{OH}$ adduct. In addition, this can lead to nucleophilic substitution by anions to give, for example, the $\text{DEPMPO}\text{-SO}_3^-$ radical adduct. Intramolecular rearrangements have been reported in which the ethyl moiety from the phosphoryl group can migrate to the nitronyl carbon to yield a C-centered radical adduct (9).

Kinetic considerations

The apparent rate constant k_{app} values for the second order reaction of various nitrones with $\text{O}_2^{\cdot-}$ and $\cdot\text{OH}$ radicals as well as the half-lives of their respective spin adducts are shown in Table 2. It should be noted, however, that rate constants can vary from one radical generating system to another because of the effect of several factors such as the presence of metal ions, ionic strength, pH, or endogenous radical scavengers. The pH dependence of the rate of $\text{O}_2^{\cdot-}$ trapping by DMPO shows that the rate of trapping is relatively faster in acidic media (18) as determined using the xanthine-xanthine oxidase superoxide generating system. This is due to the formation of hydroperoxyl radical ($\cdot\text{OOH}$), which is far more oxidizing than the unprotonated form, i.e., $E_{\text{red}}^{\circ} = 1.06 \text{ V}$ for $\cdot\text{OOH}$ versus $E_{\text{red}}^{\circ} = 0.94 \text{ V}$ for $\text{O}_2^{\cdot-}$ (5). Furthermore, reaction of DMPO with $\cdot\text{OOH}$ predominates below pH 7.7, while reaction between DMPO with $\text{O}_2^{\cdot-}$ occurs above this pH (18).

Other applications

Spin trapping using kinetic analysis has been employed to deduce the stoichiometry of O_2 reduction by NADPH to gen-

erate $\text{O}_2^{\cdot-}$. It was shown in the NADPH-cytochrome P-450 reductase-paraquat system that the stoichiometry is as follows:



as obtained from the rate extrapolated to infinite [DMPO] (96). The rate of $\text{O}_2^{\cdot-}$ production from 1-methyl-4-phenyl-2,3-dihydropyridinium, a bioactivated intermediate of the parkinsonian-inducing neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, was found to be about $k_1 = 3.97 \times 10^{-6} \text{ s}^{-1}$ from kinetic spin trapping methods using DMPO as a spin trap (97). Spin trapping with DMPO has enabled identification of $\text{O}_2^{\cdot-}$ generation from endothelial NOS (94), neuronal NOS (60), and inducible NOS (93) and characterization of the mechanisms of this process. Recently, DMPO spin trapping has been used to quantitate $\text{O}_2^{\cdot-}$ generation from the vascular NADPH oxidase (74). Simultaneous real-time measurements of free radicals and cardiac contractile function have been performed using *in vivo* spin trapping in isolated beating hearts. *In vivo* EPR techniques have enabled determination of the kinetics of free radical metabolism in normally perfused and globally ischemic hearts at 1–2 GHz EPR frequency (100).

NO DETECTION

NO is generated by the enzyme NOS from the substrates L-Arg, O_2 , and NADPH in the presence of calcium ions, calmodulin, and tetrahydrobiopterin. NO can also be generated from other molecules such as nitroglycerine, nitrites, S-nitrosothiols, diazeniumdiolates, and transition metal nitrosyl complexes. NO production in various *in vivo* and *in vitro* systems has been extensively studied by EPR spin trapping methods. NO has been detected in intact mitochondria, mitochondrial homogenates, and submitochondrial particles with great success (24). A wide variety of literature describes the use of EPR spin trapping methods to detect NO in many ap-

TABLE 2. APPARENT SECOND ORDER RATE CONSTANTS (K_{APP}) FOR THE TRAPPING OF SUPEROXIDE OR HYDROXYL RADICALS, PARTITION COEFFICIENTS (K_p), AND TOXICITY OF COMMONLY USED NITRONE SPIN TRAPS, AND HALF-LIVES ($T_{1/2}$) OF THEIR CORRESPONDING SPIN ADDUCTS

Spin traps	$\text{O}_2^{\cdot-}$ trapping*		$\cdot\text{OH}$ trapping†		Partition coefficient (K_p)‡	Toxicity	
	k_{app} ($\text{M}^{-1} \text{s}^{-1}$)	$t_{1/2}$ (min)	k_{app} ($10^{-9} \text{M}^{-1} \text{s}^{-1}$)	$t_{1/2}$ (min)		Cells [IC_{50} (mM)]	Animals (mg/100 g)§
DMPO	50 ⁽⁸⁶⁾ –60 ⁽¹⁸⁾	1 ⁽⁸⁶⁾	1.9 ⁽⁸⁷⁾ –3.4 ⁽¹⁸⁾	55 ⁽⁸⁷⁾	0.1 ⁽³⁴⁾	138 ⁽²⁹⁾	<200 ⁽⁷¹⁾
EMPO	—	8.6 ^(77, 98)	5.0 ⁽⁸⁷⁾	127 ⁽⁸⁷⁾	0.15 ⁽⁷⁷⁾	—	—
DEPMPO	60 ⁽⁸⁶⁾ –90 ⁽¹⁹⁾	14 ⁽⁸⁶⁾	4.8 ⁽⁸⁷⁾ –7.8 ⁽¹⁹⁾	132 ⁽⁸⁷⁾	0.06 ⁽¹⁾	<25 ⁽⁴⁵⁾	—
DIPPMPO	—	~23 ⁽⁷⁾	4.6 ⁽⁸⁷⁾	158 ⁽⁸⁷⁾	2.1 ⁽⁷⁾	—	—
PBN	N/A	N/A	6.1–8.5 ⁽⁶⁵⁾	0.8 ⁽³³⁾	15 ⁽³⁴⁾	9.4 ⁽²⁹⁾	<100 ⁽⁷¹⁾
4-PyOBN	N/A	N/A	4.0 ⁽⁶⁵⁾	0.2 ⁽³³⁾	0.15 ⁽³⁴⁾	5.4 ⁽²⁹⁾	<100 ⁽⁷¹⁾

References are given as superscripts. A dash indicates data not reported. N/A, not applicable.

*Using a riboflavin-light radical generating system in phosphate buffer, pH 7.0.

†Radical sources: Janzen *et al.* (33) and Villamena *et al.* (87) with UV photolysis of H_2O_2 ; Rosen *et al.* (65) with ionizing radiation- N_2O ; Finkelstein *et al.* (18) and Frejaville *et al.* (19) with $\text{Fe}^{2+}\text{-H}_2\text{O}_2$.

‡In 1-octanol/phosphate buffer system, pH 7.4.

§On Sprague-Dawley rats as lethal doses.

plications, including tumors (46) and models of sepsis such as lipopolysaccharide-induced injury (44), as well as from environmental pollution and cigarette smoke (73). NO formation in whole mice has been measured using L-band EPR spin trapping (20, 40). In rats induced with lipopolysaccharide, spin-trapped NO was observed in the liver, kidney, urinary bladder, and head regions (21).

NO is a relatively inert gas and is highly diffusible in cells and cell membranes because of its moderate solubility in water (approximately 2.0 mM atm^{-1}) at ambient temperature, or 1.4 mM atm^{-1} at 37°C (11) and with higher solubility in non-polar media. Several NO traps such as the complexes of iron with dithiocarbamate ligands (Scheme 6) have been developed that allow NO detection in both polar and non-polar environments. Vanin and co-workers (53, 83, 85) first demonstrated the use of $\text{Fe}(\text{diethyldithiocarbamate})_2$ [$\text{Fe}(\text{DETC})_2$] for NO detection in biological systems. *In vivo* EPR experiments on mice showed that NO trapping by hydrophobic Fe^{2+} -DETC is more efficient than by hydrophilic Fe^{2+} -MGD due to higher stability of this complex in animal tissues (49). In addition to NO trapping forming mononitrosyl iron complexes, dinitrosyl iron complexes can be formed typically from iron sulfur proteins. Dinitrosyl iron complexes have been observed in liver and kidney and show high affinity and membrane permeability in these organs (82).

The redox state of Fe-dithiocarbamates, either of MGD or DETC, plays a critical role in the accurate quantification of NO production. The Fe^{2+} complex can be readily oxidized under aerobic conditions to form Fe^{3+} -dithiocarbamate, which can form NO- Fe^{3+} complexes as observed by using various spectroscopic and electrochemical methods (84). Conversion of the EPR-silent NO- Fe^{3+} -MGD complex to an EPR-detectable NO- Fe^{2+} -MGD can be accomplished with 50% yield by NO itself and by reductants such as ascorbate, hydroquinone, or cysteine with conversion efficiency of up to 99.9%. To limit these redox reactions with oxygen, it has been suggested that under aerobic conditions, the oxidized ferric form of these complexes should be used for cellular or tissue measurements. Moreover, dithiocarbamates are efficient Cu(II) chelators and can inhibit copper enzymes such as SOD (10).

NO can also be generated non-enzymatically (105) in tissues by either direct disproportionation, or reduction of nitrite (NO_2^-) under acidic and highly states, which can occur

in ischemic conditions. Nitrite anions can also form NO at physiological pH, but this only occurs at a very low rate (80) since its rate of NO formation is a function of H^+ and nitrite concentrations. As a caution, it has also been reported that under some conditions Fe^{2+} -MGD can significantly catalyze the reduction of nitrite to NO (78).


Because of the facile reaction of O_2 with NO to form more reactive nitrogen oxide species such as NO_2 , N_2O_3 , and N_2O_4 , preparations of NO stock solutions and of their Fe^{2+} complexes should be made in deoxygenated water. Introduction of these spin trap complexes either *in vitro* or *in vivo* usually involve premixing of the FeSO_4 with excess dithiocarbamate co-ligand. Water-insoluble $\text{Fe}(\text{DETC})_2$ can be introduced as a suspension with serum albumin (79) and has been reported to measure NO in porcine aorta with high sensitivity of 10 pmol/ml . Detection of NO in blood vessels as well as human umbilical endothelial cells has been successfully demonstrated using colloidal Fe^{2+} -DETC prepared by mixing DETC and Fe^{2+} in concentrated Krebs-HEPES solution (38, 39).

Hemoglobin has also been employed as a spin trap and has relatively high specificity and affinity ($K_a = 3 \times 10^{10} \text{ M}^{-1}$) for NO and slow NO dissociation rate ($K_d = 10^3\text{--}10^5 \text{ s}^{-1}$) (42, 43). The highly oxygen-sensitive glucose chars, which are solid-state paramagnetic probes, have also been employed for NO detection with sensitivity of up to $10\text{--}100 \text{ nM}$ levels in anaerobic solutions but are not sensitive enough to measure physiological NO concentrations $<10 \text{ nM}$ (104).

Several non-conventional traps or probes, such as dendrimeric-containing nitronyl nitroxides (66), diazoketones (88), cheletropic *o*-quinodimethane derivatives (2), nitronyl-nitroxides (35, 91), diradicals (48), and nitromethane (62), have also been employed to detect NO, but are limited by their availability, specificity, instability in biological milieu, and short EPR signal half-lives (90).

MISCELLANEOUS SPIN TRAPPING DETECTION METHODS

Both ^{31}P - and ^{19}F -nuclear magnetic resonance (NMR) spectroscopy have been employed in the detection of various radicals through analysis of the decomposition products of their respective adducts with DEPMPO and a fluorinated DMPO analogue (36, 37). In the case of ^{31}P -NMR, O-centered radical, *e.g.*, $\cdot\text{OH}$ and $\text{O}_2^{\cdot-}$, adducts of DEPMPO gave characteristic peaks at 27.05 ppm that disappeared over time to only a peak signal corresponding to DEPMPO. Potapenko *et al.* (57) had proposed a non-radical mechanism for the formation of $\text{SO}_3^{\cdot-}$ -DEPMPO from the initial nucleophilic addition of the SO_3^{2-} anion to DEPMPO to form the corresponding hydroxylamine as a mixture of diastereomers as characterized by ^{31}P -, ^{13}C -, and ^1H -NMR spectroscopy, and its subsequent oxidation leads to the formation of its corresponding nitroxides. Using electrochemical (47) and liquid chromatography with electrochemical detection (54) techniques, spin adducts and their decomposition products have been analyzed. Lipid derived C-centered pentadienyl radicals have been detected and identified *in vitro* and *in vivo* by liquid chromatography/EPR and liquid chromatography/mass spectrometry (61) using



SCHEME 6. Various types of dithiocarbamates used as ligands for NO trapping.

4-PyOBN as a spin trap. Proton magnetic resonance imaging detection of NO production based on (MGD)₂-Fe²⁺-NO complex formation and the effect of these complexes on proton relaxation have been reported in rats (22).

Cyclic hydroxylamines are also employed as OFR probes although not spin traps *per se*, since the nitroxide formation results from the oxidation of the hydroxylamine by OFRs to give an EPR signal, and there is no trapping/bonding of the radical. The specificity of these cyclic hydroxylamines is still not clear at the moment since the effect of other potential oxidizing agents in the biological milieu has not yet been reported. A negatively charged cyclic hydroxylamine, 1-hydro-4-phosphonoxy-2,2,6,6-tetramethylpiperidine, has been used to detect ROS for diagnosis of extracellular inflammation-induced oxidative stress in cultured macrophages, blood, and *in vivo* applications (13).

CONCLUSION

EPR spin trapping is a powerful technique for measuring free radicals in biomedical applications. It enables sensitive and specific detection of free radicals and characterization of the mechanisms of their formation in a broad range of applications. In spite of the limitations associated with spin trapping techniques such as the need to improve efficiency of radical trapping, selectivity and availability of spin traps, the limited stability of spin adducts, and possible formation of artifactual spin adducts, the use of this technique for detection of free radicals has become an indispensable tool for the detection of OFRs and NO in biological systems. There is an ongoing need for improved spin traps and related development efforts. Further progress in spin trap development together with advances in EPR technology, including low-frequency EPR instrumentation, for *in vivo* applications should enable this technique to be extended to address many critical questions regarding the role of free radicals in cellular function and disease in animal models and in humans.

ABBREVIATIONS

L-Arg, L-arginine; DEPMPO, 5-diethoxyphosphoryl-5-methyl-1-pyrroline *N*-oxide; DETC, diethyldithiocarbamate; DMPO, 5,5-dimethyl-1-pyrroline *N*-oxide; DTCS, *N*-(dithiocarboxy)sarcosine; EMPO, 5-ethoxycarbonyl-5-methyl-1-pyrroline *N*-oxide; EPR, electron paramagnetic resonance; MGD, *N*-methyl-D-glucamine dithiocarbamate; NMR, nuclear magnetic resonance; NO, nitric oxide; NOS, nitric oxide synthase; OFR, oxygen free radical; PBN, *N*-tert-butyl- α -phenyl nitroxide; PUFA, polyunsaturated fatty acid; 4-PyOBN, α -(4-pyridyl-1-oxide)-*N*-tert-butyl nitroxide; RNS, reactive nitrogen species; ROS, reactive oxygen species; SOD, superoxide dismutase; UV, ultraviolet.

REFERENCES

- Barbati S, Clement JL, Olive G, Roubaud V, Tuccio B, and Tordo P. ¹³P labeled cyclic nitrones: a new class of spin traps for free radicals in biological milieu. In: *Free Radicals in Biology, Environment*, edited by Minisci F. Dordrecht, The Netherlands: Kluwer Academic Publishers, 1997, pp. 39–47.
- Batz M, Korth H-G, and Sustmann R. A novel method for the detecting nitric oxide by formation of fluorescent products based on cheletropic spin traps. *Angew Chem Int Ed* 36: 1501–1503, 1997.
- Benson SW. Kinetics of pyrolysis of alkyl hydroperoxides and their O-O bond dissociation energies. *J Chem Phys* 40: 1007–1013, 1964.
- Bilenko MV. *Ischemia and Reperfusion of Various Organs: Injury, Mechanisms, Methods of Prevention and Treatment*. Huntington, NY, Nova Science, 2001.
- Buettner GR. The pecking order of free radicals and antioxidants: lipid peroxidation, α -tocopherol and ascorbate. *Arch Biochem Biophys* 300: 535–543, 1993.
- Buettner GR and Oberley LW. Consideration in the spin trapping of superoxide and hydroxyl in aqueous systems using 5,5-dimethyl-1-pyrroline-1-oxide. *Biochem Biophys Res Commun* 83: 69–74, 1978.
- Chalier F and Tordo P. DIPPMPPO, a crystalline analog of the nitroxide DEPMPO, synthesis and spin trapping properties. *J Chem Soc Perkin Trans 2*: 2110–2117, 2002.
- Chen YR, Gunther MR, and Mason RP. An electron spin resonance spin-trapping investigation of the free radicals formed by the reaction of mitochondrial cytochrome c oxidase with H₂O₂. *J Biol Chem* 274: 3308–3314, 1999.
- Clement JL, Gilbert BC, Ho WF, Jackson ND, Newton MS, Silvester S, Timmins GS, Tordo P, and Whitwood AC. Use of a phosphorylated spin trap to discriminate between hydroxyl radical and other oxidising species. *J Chem Soc Perkin Trans 2*: 1715–1717, 1998.
- Cocco D, Calabrese L, Rigo A, Argese E, and Rotilio G. Re-examination of the reaction of diethyldithiocarbamate with the copper of superoxide dismutase. *J Biol Chem* 256: 8983–8986, 1981.
- Dean JA. *Lange's Handbook of Chemistry*, 15th ed. New York: McGraw-Hill, 1998.
- Dikalov S and Mason RP. Spin trapping of polyunsaturated fatty acid-derived peroxy radicals: reassignment to alkoxy radical adducts. *Free Radic Biol Med* 30: 187–197, 2001.
- Dikalov S, Dikalova A, and Mason RP. Noninvasive diagnostic tool for inflammation-induced oxidative stress using electron spin resonance spectroscopy and an extracellular cyclic hydroxylamine. *Arch Biochem Biophys* 402: 218–226, 2002.
- Eaton GR, Eaton SS, and Ohno K. *EPR Imaging and In Vivo EPR*. Boca Raton, FL: CRC Press, 1991.
- Ebersson L. Spin trapping: problems and artifacts. In: *Toxicology of the Human Environment: The Critical Role of Free Radicals*, edited by Rhodes CJ. New York: Taylor and Francis, 2000, pp. 25–41.
- ESR at NIEHS/NIH: the gateway to biomedical ESR research [Online]. National Institute of Environmental Health Science. <http://epr.niehs.nih.gov/>, 2002, April 4.
- Feix JB and Kalyanaram B. Spin trapping of lipid-derived radicals in liposomes. *Biochim Biophys Acta* 992: 230–235, 1989.

1. Barbati S, Clement JL, Olive G, Roubaud V, Tuccio B, and Tordo P. ¹³P labeled cyclic nitrones: a new class of

18. Finkelstein E, Rosen GM, and Rauckman EJ. Spin trapping. Kinetics of the reaction of superoxide and hydroxyl radicals with nitrones. *J Am Chem Soc* 102: 4995, 1980.
19. Frejaville C, Karoui H, Tuccio B, Le Moigne F, Culcasi M, Pietri S, Lauricella R, and Tordo P. 5-(Diethoxyphosphoryl)-5-methyl-1-pyrroline N-oxide: a new efficient phosphorylated nitron for the *in vitro* and *in vivo* spin trapping of oxygen-centered radicals. *J Med Chem* 38: 258–265, 1995.
20. Fujii H and Berliner LJ. Nitric oxide: prospects and perspectives of *in vivo* detection by L-band EPR spectroscopy. *Phys Med Biol* 43: 1949–1956, 1998.
21. Fujii H and Berliner LJ. Ex vivo EPR detection of nitric oxide in brain tissue. *Magn Reson Med* 42: 599–602, 1999.
22. Fujii H, Wan X, Zhong J, Berliner LJ, and Yoshikawa K. In vivo imaging of spin-trapped nitric oxide in rats with septic shock: MRI spin trapping. *Magn Reson Med* 42: 235–239, 1999.
23. Gianni L, Zweier JL, Levy A, and Myers CE. Characterization of the cycle of iron-mediated electron transfer from Adriamycin to molecular oxygen. *J Biol Chem* 260: 6820–6826, 1985.
24. Giulivi C, Poderoso JJ, and Boveris A. Production of nitric oxide by mitochondria. *J Biol Chem* 273: 11038–11043, 1998.
25. Greenley TL, and Davies MJ. Radical production from peroxide and peracid tumor promoters: EPR spin trapping studies. *Biochem Biophys Acta* 1157: 23–31, 1993.
26. Halliwell B and Gutteridge JMC. Oxidative stress: adaptation, damage and death. In: *Free Radicals in Biology and Medicine*. Oxford: Oxford University Press, 1999, pp. 246–350.
27. Han D, Williams E, and Cadenas E. Mitochondrial respiratory chain-dependent generation of superoxide anion and its release into the intermembrane space. *Biochem J* 353: 411–416, 2001.
28. Hanna PH, Chamulitrat W, and Mason RP. When are metal ion-dependent hydroxyl and alkoxyl radical adducts of DMPO artifacts? *Arch Biochem Biophys* 296: 640–644, 1992.
29. Haseloff RF, Mertsch K, Rohde E, Baeger I, Grigor'ev IA, and Blasig IE. Cytotoxicity of spin trapping compounds. *FEBS Lett* 418: 73–75, 1997.
30. Hiramoto K, Ohkawa T, and Kikugawa K. Release of nitric oxide together with carbon-centered radicals from N-nitrosamines by ultraviolet light irradiation. *Free Radic Res* 35: 803–813, 2001.
31. Janzen EG. A critical review of spin trapping in biological systems. *Free Radic Biol Med* 4: 115–154, 1980.
32. Janzen EG and Haire DL. Two decades of spin trapping. *Adv Free Radic Chem* 1: 253–295, 1990.
33. Janzen EG, Kotake Y, and Hinton RD. Stabilities of hydroxyl radical spin adducts of PBN-type spin traps. *Free Radic Biol Med* 12: 169–173, 1992.
34. Janzen EG, West MS, Kotake Y, and DuBose CM. Biological spin trapping methodology. III. Octanol-water partition coefficients of spin-trapping compounds. *J Biochem Biophys Methods* 32: 183–190, 1996.
35. Joseph J, Kalyanaraman B, and Hyde JS. Trapping of nitric oxide by nitronyl nitroxide: an electron spin resonance investigation. *Biochem Biophys Res Commun* 192: 926–934, 1993.
36. Khramtsov VV, Berliner LJ, and Clanton TL. NMR spin trapping: detection of free radical reactions using a phosphorous containing nitron spin trap. *Magn Reson Med* 42: 228–234, 1999.
37. Khramtsov VV, Reznikov VA, Berliner LJ, Litkin AK, Grigor'ev IA, and Clanton TL. NMR spin trapping: detection of free radical reactions with a new fluorinated DMPO analog. *Free Radic Biol Med* 30: 1099–1107, 2001.
38. Kleschyov AL and Munzel T. Advanced spin trapping of vascular nitric oxide using colloid iron dithiocarbamate. *Methods Enzymol* 359: 43–51, 2002.
39. Kleschyov AL, Hanke Mollnau H, Oelze M, Meinertz T, Huang Y, Harrison DG, and Munzel T. Spin trapping of vascular nitric oxide using colloidal Fe(II)-diethyldithiocarbamate. *Biochem Biophys Res Commun* 275: 672–677, 2000.
40. Komarov AM. In vivo detection of nitric oxide distribution in mice. *Mol Cell Biochem* 234/235: 387–392, 2002.
41. Kondo T and Riesz P. Sonochemistry of nitron spin traps in aqueous solutions. Evidence for pyrolysis radicals from spin traps. *Free Radic Biol Med* 7: 259, 1989.
42. Kosaka H and Shiga T. Detection of nitric oxide by electron spin resonance using hemoglobin. In: *Methods in Nitric Oxide Research*, edited by Feelisch M and Stamler JS. New York: John Wiley & Sons, 1996, pp. 373–381.
43. Kosaka H, Watanabe M, and Yoshihara H. Detection of nitric oxide production in lipopolysaccharide-treated rats by ESR using carbon monoxide hemoglobin. *Biochim Biophys Res Commun* 184: 1119–1124, 1992.
44. Lecour S, Chevet D, Maupoil V, Moisan M, Bernard C, Zahnd J, Touchard G, Briot F, and Rochette L. Intrarenal detection of nitric oxide using electron spin resonance spectroscopy in hypertensive lipopolysaccharide-treated rats. *J Cardiovasc Pharmacol* 40: 9–17, 2002.
45. Liu KJ, Miyake M, Panz T, and Swartz H. Evaluation of DEPMPO as a spin trapping agent in biological systems. *Free Radic Biol Med* 26: 714–721, 1999.
46. Loibl S, von Minckwitz G, Weber S, Sinn H-P, Schinik-Kerth VB, Lobysheva I, Nepveu F, Wolf G, Strebhardt K, and Kaufmann M. Expression of endothelial and inducible nitric oxide synthase in benign and malignant lesions of the breast and measurement of nitric oxide using electron paramagnetic resonance spectroscopy. *Cancer* 95: 1191–1198, 2002.
47. Ma Z, Zhao B, and Yuan Z. Application of electrochemical and spin trapping techniques in the investigation of hydroxyl radicals. *Anal Chim Acta* 389: 213–218, 1999.
48. Marx L and Rassat A. Application of a spin-labeled spin trap to the detection of nitric oxide. *Angew Chem Int Ed* 39: 4494–4496, 2000.
49. Mikoyan VD, Kubrina LN, Serezhnikov VA, Stukan RA, and Vanin AF. Complexes of Fe(II) with diethyldithiocarbamate or N-methyl-D-glucamine dithiocarbamate as traps of nitric oxide in animal tissues: comparative investigations. *Biochim Biophys Acta* 1336: 225–234, 1997.
50. Misik V and Riesz P. Free radical formation by ultrasound in organic liquids: a spin trapping and EPR study. *J Phys Chem* 98: 1634–1640, 1994.

51. Misik V, Miyoshi N, and Riesz P. EPR spin-trapping study of the sonolysis of H₂O/D₂O mixtures: probing the temperatures of cavitation regions. *J Phys Chem* 99: 3605–3611, 1995.
52. Mitsuta K, Hiramatsu M, Ohya-Nishiguchi H, Kamada H, and Fujii K. A kinetic analysis of superoxide adduct formation in the presence of typical scavengers. *Bull Chem Soc Jpn* 67: 529–538, 1994.
53. Mordvintcev P, Mulsch A, Busse R, and Vanin AF. On-line detection of nitric oxide formation in liquid aqueous phase by EPR. *Anal Biochem* 199: 142, 1991.
54. Novakov CP, Feierman D, Cederbaum AI, and Stoyanovsky DA. An ESR and HPLC-EC assay for the detection of alkyl radicals. *Chem Res Toxicol* 14: 1239–1246, 2001.
55. Olive G, Mercier A, Le Moigne F, Rockenbauer A, and Tordo P. 2-Ethoxycarbonyl-2-methyl-3,4-dihydro-2H-pyrrole-1-oxide: evaluation of the spin trapping properties. *Free Radic Biol Med* 28: 403–408, 2000.
56. Ouari O, Polidori A, Pucci B, Tordo P, and Chaher F. Synthesis of a glycolipidic amphiphilic nitron as a new spin trap. *J Org Chem* 64: 3554–3556, 1999.
57. Potapenko DI, Clanton TL, Bagryanskaya EG, Gritsan NP, Reznikov VA, and Khramtsov V. Nonradical mechanism of (bi)sulfite reaction with DEPMPO: cautionary note for SO₃^{•-} radical spin trapping. *Free Radic Biol Med* 34: 196–206, 2003.
58. Pou S, Rosen GM, Wu Y, and Rosen GM. Synthesis of deuterium and ¹⁵N-containing pyrroline 1-oxides: spin trapping study. *J Org Chem* 55: 4438–4443, 1990.
59. Pou S, Ramos CL, Gladwell T, Renks E, Centra M, Young D, Cohen MS, and Rosen GM. A kinetic approach to the selection of a sensitive spin trapping system for the detection of hydroxyl radical. *Anal Biochem* 217: 76–83, 1994.
60. Pou S, Keaton L, Surichamorn W, and Rosen GM. Mechanism of superoxide generation by neuronal nitric-oxide synthase. *J Biol Chem* 274: 9573–9580, 1999.
61. Qian SY, Tomer KB, Yue G-H, Guo Q, Kadiiska MB, and Mason RP. Characterization of the initial carbon-centered pentadienyl radical and subsequent radicals in lipid peroxidation: identification via on-line high performance liquid chromatography/electron spin resonance and mass spectrometry. *Free Radic Biol Med* 33: 998–1009, 2002.
62. Reszka K, Chignell CF, and Bilski P. Spin trapping of nitric oxide by aci-nitromethane in aqueous solutions. *J Am Chem Soc* 116: 4119–4120, 1994.
63. Rhodes CJ. *Toxicology of the Human Environment*. London: Taylor and Francis, 2000.
64. Rosen GM and Finkelstein E. Use of spin trap in biological system. *Adv Free Radic Biol Med* 1: 345–375, 1985.
65. Rosen GM, Britigan BE, Halpern HJ, and Pou S. *Free Radicals: Biology and Detection by Spin Trapping*. New York: Oxford University Press, 1999.
66. Rosen GM, Porasuphatana S, Tsai P, Ambulos NP, Galtsev VE, Ichikawa K, and Halpern HJ. Dendrimeric-containing nitronyl nitroxides as spin traps for nitric oxide: synthesis, kinetic, and stability studies. *Macromolecules* 36: 1021–1027, 2003.
67. Roubaud V, Sankarapandi S, Kuppusamy P, Tordo P, and Zweier JL. Quantitative measurement of superoxide generation using the spin trap 5-(diethoxyphosphoryl)-5-methyl-1-pyrroline-N-oxide. *Anal Biochem* 247: 404–411, 1997.
68. Roubaud V, Sankarapandi S, Kuppusamy P, Tordo P, and Zweier JL. Quantitative measurement of superoxide generation and oxygen consumption from leukocytes using electron paramagnetic resonance spectroscopy. *Anal Biochem* 257: 210–217, 1998.
69. Samuni A, Carmichael AJ, Russo A, Mitchell JB, and Riesz P. On the spin trapping and ESR detection of oxygen-derived radicals generated inside cells. *Proc Natl Acad Sci U S A* 83: 7593–7597, 1986.
70. Sankuratri N and Janzen EG. Synthesis and spin trapping chemistry of a novel bicyclic nitron: 1,3,3-trimethyl-6-azabicyclo[3.2.1]oct-6-ene-N-oxide (Trazon). *Tetrahedron Lett* 37: 5313–5316, 1996.
71. Schaefer CF, Janzen EG, West MS, Poyer JL, and Kosanke SD. Blood chemistry changes in the rat induced by high doses of nitronyl free radical spin traps. *Free Radic Biol Med* 21: 427–436, 1996.
72. Sheng P, Feix J, and Kalyanaraman B. Characterization of radical adducts formed during photochemical spin trapping in liposomes. *Photochem Photobiol* 52: 523–531, 1990.
73. Shinagawa K, Tokimoto T, and Shirane K. Spin trapping of nitric oxide in aqueous solutions of cigarette smoke. *Biochem Biophys Res Commun* 253: 99–103, 1998.
74. Souza HP, Liu X, Samouilov A, Kuppusamy P, Laurindo FR, and Zweier J. Quantitation of superoxide generation and substrate utilization by vascular NAD(P)H oxidase. *Am J Physiol* 282: H466–H474, 2002.
75. Spiekermann S, Landmesser U, Dikalov S, Bredt M, Gamez G, Tatge H, Reepschlager N, Hornig B, Drexler H, and Harrison DG. Electron spin resonance characterization of vascular xanthine and NAD(P)H oxidase activity in patients with coronary artery disease: relation to endothelium-dependent vasodilation. *Circulation* 107: 1383–1389, 2003.
76. Stolze K, Udilova N, and Nohl H. Spin trapping of lipid radicals with DEPMPO-derived spin traps: detection of superoxide, alkyl and alkoxy radicals in aqueous and lipid phase. *Free Radic Biol Med* 29: 1005–1014, 2000.
77. Stolze K, Udilova N, and Nohl H. Spin adducts of superoxide, alkoxy, and lipid-derived radicals with EMPO and its derivatives. *Biol Chem* 383: 813–820, 2002.
78. Stolze K, Udilova N, and Nohl H. ESR analysis of spin adducts of alkoxy and lipid-derived radicals with the spin trap Trazon. *Biochem Pharmacol* 63: 1465–1470, 2002.
79. Tsuchiya K, Takasugi M, Minakuchi K, and Fukuzawa K. Sensitive quantification of nitric oxide by EPR spectroscopy. *Free Radic Biol Med* 21: 733–737, 1996.
80. Tsuchiya K, Yoshizumi M, Houchi H, and Mason RP. Nitric oxide-forming reaction between the iron-N-methyl-D-glucamine dithiocarbamate complex and nitrite. *J Biol Chem* 2000: 1551–1556, 2000.
81. Turner MJ and Rosen GM. Spin trapping of superoxide and hydroxyl radicals with substituted pyrroline 1-oxides. *J Med Chem* 29: 2439–2444, 1986.

82. Ueno T, Suzuki Y, Fujii S, Vanin AF, and Yoshimura T. In vivo distribution and behavior of paramagnetic dinitrosyl dithiolato iron complex in the abdomen of mouse. *Free Radic Res* 31: 525–534, 1999.
83. Vanin AF. Iron diethyldithiocarbamate as spin trap for nitric oxide detection. *Methods Enzymol* 301: 269–279, 1999.
84. Vanin AF, Liu X, Samouilov A, Stukan RA, and Zweier JL. Redox properties of iron-dithiocarbamates and their nitrosyl derivatives: implications for their use as traps of nitric oxide in biological system. *Biochim Biophys Acta* 1474: 365–377, 2000.
85. Vanin AF, Huisman A, and Van Faassen EE. Iron dithiocarbamate as spin trap for nitric oxide detection: pitfalls and successes. *Methods Enzymol* 359: 27–42, 2002.
86. Villamena F and Zweier J. Superoxide radical trapping and spin adduct decay of 5-tert-butoxycarbonyl-5-methyl-1-pyrroline N-oxide (BocMPO): kinetics and theoretical analysis. *J Chem Soc Perkin Trans 2*: 1340–1344, 2002.
87. Villamena F, Hadad CM, and Zweier J. Kinetic study and theoretical analysis of hydroxyl radical trapping and spin adduct decay of alkoxycarbonyl- and dialkoxyporphoryl-nitrones in aqueous media. *J Phys Chem A* 107: 4407–4414, 2003.
88. Walton JW and Park JSB. Exploratory study of diazoketones as spin traps for nitroxide. *J Chem Soc Perkin Trans 2*: 2585–2588, 1997.
89. Weil JA, Bolton JR, and Wertz JE. *Electron Paramagnetic Resonance: Elementary Theory and Practical Applications*. New York: John Wiley and Sons, 1994.
90. Williams DLH. A chemist's view of the nitric oxide story. *Org Biomol Chem* 1: 441–449, 2003.
91. Woldman YY, Khramtsov V, Grigor'ev IA, Kiriljuk IA, and Utepbergenov DI. Spin trapping of nitric oxide by nitronyl nitroxides: measurements of the activity of NO synthase from rat cerebellum. *Biochem Biophys Res Commun* 202: 195–203, 1994.
92. Xia Y and Zweier JL. Superoxide and peroxynitrite generation from inducible nitric oxide synthase in macrophages. *Proc Natl Acad Sci U S A* 94: 6954–6958, 1997.
93. Xia Y, Roman LJ, Masters SS, and Zweier JL. Inducible nitric-oxide synthase generates superoxide from the reductase domain. *J Biol Chem* 273: 22635–22639, 1998.
94. Xia Y, Tsai AL, Berka V, and Zweier JL. Superoxide generation from endothelial nitric oxide synthase: a Ca^{2+} /calmodulin dependent and tetrahydrobiopterin regulatory process. *J Biol Chem* 273: 25804–25808, 1998.
95. Xu YK, Chen ZW, Sun J, Liu K, Chen W, Shi W, Wang HM, Zhang XK, and Liu Y. Synthesis and ESR study of a novel phosphorylated lipophilic spin trap. *J Org Chem* 67: 7624–7630, 2002.
96. Yamasaki I, Piette LH, and Grover TA. Kinetic studies on the spin trapping of superoxide and hydroxyl radicals generated in NADPH-cytochrome P-450 reductase-paraquat systems. *J Biol Chem* 265: 652–659, 1990.
97. Zang LY and Misra HP. EPR kinetic studies of superoxide radicals generated during the autoxidation of 1-methyl-4-phenyl-2,3-dihydropyridinium, a bioactivated intermediate of parkinsonian-inducing neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *J Biol Chem* 267: 23601–23608, 1992.
98. Zhang H, Joseph J, Vasquez-Vivar J, Karoui H, Nsanzumuhire C, Martasek P, Tordo P, and Kalyanaraman B. Detection of superoxide anion using an isotopically labeled nitron spin trap: potential biological applications. *FEBS Lett* 473: 58–62, 2000.
99. Zweier JL. Iron-mediated formation of an oxidized adriamycin free radical. *Biochim Biophys Acta* 839: 209–213, 1985.
100. Zweier JL and Kuppusamy P. Electron paramagnetic resonance measurements of free radicals in the intact beating heart: a technique for detection and characterization of free radicals in whole biological tissues. *Proc Natl Acad Sci U S A* 85: 5703–5707, 1988.
101. Zweier JL and Villamena FV. Chemistry of free radicals in biological systems. In: *Oxidative Stress and Cardiac Failure*, edited by Kukin ML and Fuster V. Armonk, NY: Futura Publishing Co., 2003, pp. 67–95.
102. Zweier JL, Kuppusamy P, and Luty GA. Measurement of endothelial cell free radical generation: evidence for a central mechanism of free radical injury in postschismic tissues. *Proc Natl Acad Sci U S A* 85: 4046–4050, 1988.
103. Zweier JL, Kuppusamy P, Williams R, Rayburn BK, Smith D, Weisfeldt ML, and Flaherty JT. Measurement and characterization of postschismic free radical generation in the isolated perfused heart. *J Biol Chem* 264: 18890–18895, 1989.
104. Zweier JL, Samouilov A, and Chzhan M. Measurement of nitric oxide with a solid-state-clear EPR probe. *J Magn Reson B* 109: 259–263, 1995.
105. Zweier JL, Samouilov A, and Kuppusamy P. Non-enzymatic nitric oxide synthesis in biological systems. *Biochim Biophys Acta* 1411: 250–262, 1999.

Address reprint requests to:

Jay L. Zweier, M.D.

110G Davis Heart and Lung Research Institute

473 West 12th Avenue

Columbus, OH 43210-1252

E-mail: Zweier-1@medctr.osu.edu

Received for publication July 3, 2003; accepted February 19, 2004.