

PhD Math Camp

Relations and Functions

Ignacio Urbina

Relations

Understanding Relations

- **Definition:** A relation is a set of ordered pairs, a subset of the Cartesian product of two sets.
- **Components:** Consists of a domain and a range.
- **Visualization:** Often represented using Venn diagrams or graphs.
- **Non-Numeric Example:** Consider the relation R on the set $X = \{\text{students}\}$ and $Y = \{\text{majors}\}$:
 - Let $X = \{\text{Jordan, Bob, Carol}\}$
 - Let $Y = \{\text{History, Engineering, Biology}\}$
 - Define R as follows:

$$R = \{(\text{Jordan, History}), (\text{Bob, Engineering}), (\text{Carol, Biology})\}$$

- Here, R relates each student to their major.

Basic Types of Relations

- **Reflexive:** Every element is related to itself.
- **Symmetric:** If x is related to y , then y is related to x .
- **Transitive:** If x is related to y and y is related to z , then x is related to z .

Special Types of Relations

- **Functions:** Each element of the domain is associated with exactly one element of the range.
- **Equivalence Relations:** Reflexive, symmetric, and transitive.
- **Order Relations:** Reflexive, antisymmetric, and transitive. Important in hierarchical structures.

Applications in Social Sciences

- **Social Networks:** Analysis of friendships, social media connections.
- **Organizational Structures:** Supervision and hierarchy in corporations.
- **Market Relations:** Consumer-product interactions, market dynamics.
- **Sociological Models:** Social mobility and status relations.

Introduction of Functions

A mathematical function is a “mapping” (i.e., specific directions), which gives a correspondence from one measure onto exactly one other for that value.



Introduction of Functions

A function is a mapping from one defined space to another.

$$f : A \rightarrow B$$

- A is the domain of the function, B is the codomain of the function.
- The set of all values actually reached by running each $x \in A$ through f is known as the range (or image).

Function as a Box, Process, or Recipe



Introduction of Functions

Functions can be “many-to-one” or “one-to-one”, but cannot be “one-to-many”.

Examples:

- Function, many-to-one: $y = f(x) = x^2$
- Function, one-to-one: $y = f(x) = 2x + 4$
- Relation that is not a function: $y^2 = 5x$

Relation that is not a function




Figure: The parabola described by $y^2 = 5x$

Dimensions

$$f : \mathbb{R}^1 \rightarrow \mathbb{R}^1$$

Examples:

- $f(x) = x + 1$
- $f(x) = 2x + 3$
- $f(x) = x^2 - 4x + 4$
- $f(x) = \sin(x)$
- $f(x) = e^x$

Dimensions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^1, \quad \text{e.g. } f(x, y) = x^2 + y^2$$





Figure: $f(x, y) = x^2 + y^2$


Dimensions

$$f(x, y) = x^2 + y^2$$



Dimensions

$$f(x, y) = x^2 + y^2$$



Dimensions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^1, \quad \text{e.g. } f(x, y) = x^3 + 2y^2$$





Figure: $f(x, y) = x^3 + 2y^2$

Dimensions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^1, \quad \text{e.g. } f(x, y) = 2x + y$$

For each ordered pair (called tuple) (x, y) , $f(x, y)$ assigns the number $2x + y$



Function Composition (Nested Function)

If we have $f : A \rightarrow B$ and $g : B \rightarrow C$, then $g(f(x)) : A \rightarrow C$.

- Can also be denoted as $g \circ f(x)$.
- E.g., if $f(x) = 2x$ and $g(x) = x^3$, then $g \circ f(x) = (2x)^3 = 8x^3$.

Inverse Function

Function that when composed with the original function returns the identity function.

$$f(x) = y \Leftrightarrow f^{-1}(y) = x$$

i.e. $f^{-1}(f(x)) = x$

- E.g., if $f(x) = 3x + 2$, then $f^{-1}(x) = \frac{x-2}{3}$.
- If $f(x) = x^3 + 2$, then $f^{-1}(x) = \sqrt[3]{x-2}$.

Polynomial Functions

Polynomial functions of x are functions that have components that raise x to some power, e.g.,

$$f(x) = x^3 + x^2 + x + 1$$


Solving Equations

Roots of a polynomial function: where the curve of the function crosses the x-axis, i.e. $f(x) = 0$.

- E.g., $x^3 - 5x^2 + 2x + 8 = 0$

- Factoring polynomials:

$$x^3 - 5x^2 + 2x + 8 = (x + 1)(x - 2)(x - 4) = 0$$



Solving Quadratic Equations by Formula

Roots of a quadratic polynomial: if $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- Exercise: Solve for x : $f(x) = x^2 + 3x - 4 = 0$


Logarithmic and Exponential

$$y = \log_a(x) \Leftrightarrow a^y = x$$

- The log function can be thought of as an inverse for exponential functions.
- a is referred to as the “base” of the logarithm.

Logarithmic and Exponential with e

A comparison of the functions $y = e^x$, $y = x$, and $y = \ln(x)$ on the same graph.



Properties of Exponential Functions

- $a^x \cdot a^y = a^{x+y}$
- $a^{-x} = \frac{1}{a^x}$
- $\frac{a^x}{a^y} = a^{x-y}$
- $(a^x)^y = a^{xy}$
- $a^0 = 1$

Basic Properties of Logarithms

- **Zero/One** $\log_b(1) = 0$
- **Multiplication** $\log(x \cdot y) = \log(x) + \log(y)$
- **Division** $\log\left(\frac{x}{y}\right) = \log(x) - \log(y)$
- **Exponentiation** $\log(x^y) = y \log(x)$
- **Basis** $\log_b(b^x) = x$, and $b^{\log_b(x)} = x$

These properties hold for any base.

- Exercise: Simplify $\log\left(\frac{x^9 y^5}{z^3}\right)$

Monotonic Functions

A monotonic function is a function that:

- Preserves order.
- Is (strictly) increasing over its domain.

Which is important in optimization!

Monotonic Functions (cont'd)

A function $f(x)$ is said to be monotonic if it is either entirely non-increasing or non-decreasing.

- For example, $f(x) = 2x + 3$ is strictly increasing.