# Introduction to Statistical Methods in Political Science

Quiz 4

Ignacio Urbina

Ph.D. Candidate in Political Science

### Quiz 4

## Question 1 - Discrete Random Variable and Probability Calculation

Consider a non-standard six-sided die with faces  $\{1, 2, 2, 3, 3, 4\}$ . The face "2" appears twice, as does "3", while "1" and "4" each appear once. You roll this die twice. Let  $X_1$  be the outcome of the first roll and  $X_2$  be the outcome of the second roll.

Define  $Z = X_1 + X_2$ , as the *random variable* representing the sum of the outcomes of the two rolls.

**Question:** Compute P(Z > 7).

#### Question and Correct Answer

- Question 1. What is the probability that Z is larger than 7?
  - 1/36 ← Correct Answer
  - -15/36
  - 50%
  - 0%
  - 6.25%

### Question 2 - Bayes' Rule

A burglary has been committed in a residential neighborhood. The police have identified a suspect but do not have definitive proof. Initially, based on past crime data, only  $\bf 1$  in  $\bf 500$  (0.2%) of individuals in similar circumstances have actually committed a burglary.

A rare type of shoeprint is found at the crime scene, and the suspect owns the same type of shoes (*Evidence*). From previous forensic studies, we know:

- If a person **did** commit the crime, the probability that their shoeprint would be found is  $P(Evidence \mid Guilty) = 0.75$ .
- If a person **did not** commit the crime, the probability that they own the same rare shoe type is  $P(Evidence \mid Not \; Guilty) = 0.05$ .

**Question:** Using **Bayes' Rule**, compute the updated probability that the suspect actually committed the crime given that their shoeprint type was found at the scene – i.e., compute  $P(Guilty \mid Evidence)$ .

#### Question and Correct Answer

- Question 2. What is the probability that the subject is truly guilty, given that their shoeprint type was found at the scene? (hint: use Bayes' rule).
  - 2.9% ← Correct Answer
  - 0.2%
  - 75%
  - 93.75%
  - 70%
  - 40%
  - 1.5%

#### Question 3 - Binomial Distribution and Complement Rule

A 5-member congressional committee is voting on an amendment to a given bill. Each member independently supports the amendment with probability p=0.60. Let Y be the number of members (out of 5) who vote in favor.

#### Questions:

• Compute the probability that at least two committee members vote in favor of the amendment, i.e.,  $P(Y \ge 2)$ .

#### Question and Correct Answer

- Question 3. What is the probability that at least two committee members will vote against the amendment?
  - 66.30% ← Correct Answer
  - 34.56%
  - 68.26%
  - 80%
  - 16%
  - 40%
  - 33.70%