Split-plot Analysis

Diego Della Lunga, Ashmita Upadhyay, Igor Kuivjogi Fernandes, Timothy Goodhart, Amanda Lay

Introduction

- ► Split-plot design (RCBD)
- ► EDA
- Linear Mixed Model
- ► Mean Comparisons

Factors

Whole-plot: Irrigation

- 1. moist
- 2. saturated
- 3. flooded

Sub-plot: Fertilizer

- 1. 0
- 2. 50
- 3. 100
- 4. 150

Creating the design

```
sp <- FielDHub::split_plot(
  wp = 3,  # whole-plot
  sp = 4,  # sub-plot
  reps = 3,  # blocks
  type = 2,  # for RCBD
  seed = 2023
)</pre>
```

Field layout

Split Plot Design (RCBD) 12X3 2|3 3|4 1|3 2|4 3|1 1|4 2|2 3|3 1|2 2|1 3|2 1|1 1|3 2|2 3|1 2|4 1|2 3|3 ROWS 1|4 2|3 3|4 1|1 2|1 3|2 1|2 2|2 3|4 1|3 2|3 3|1 1|4 2|4 3|2 1|1 2|1 3|3

COLUMNS

Simulating effects of treatments

```
# effect of irrigation
split_plot$y_irr <- NA
split_plot$y_irr[split_plot$irrigation == "flooded"] <- 6</pre>
split_plot$y_irr[split_plot$irrigation == "saturated"] <- 12</pre>
split_plot$y_irr[split_plot$irrigation == "moist"] <- 15</pre>
# effect of fertilizer
split_plot$y_fert <- NA
split_plot$y_fert[split_plot$fertilizer == "0"] <- 5</pre>
split_plot$y_fert[split_plot$fertilizer == "50"] <- 8</pre>
split_plot$y_fert[split_plot$fertilizer == "100"] <- 10</pre>
split_plot$y_fert[split_plot$fertilizer == "150"] <- 12</pre>
# effect of interaction
split_plot$y_inter <- (</pre>
  split_plot$y_irr * split_plot$y_fert
) * 0.01
```

Simulating effects of blocks and error


```
# effect of block
split_plot$y_b <- NA
split_plot$y_b[split_plot$REP == 1] <- -10</pre>
split_plot$y_b[split_plot$REP == 2] <- 5</pre>
split_plot$y_b[split_plot$REP == 3] <- -5</pre>
split_plot$y_b[split_plot$REP == 4] <- -10</pre>
# effect of error
set.seed(2023)
split_plot$error <- rnorm(36, 0, 2.5)</pre>
# creating response
split_plot$y <- (
  split_plot$y_irr + split_plot$y_fert +
  split_plot$y_inter + split_plot$error + split_plot$y_b
```

```
with(split_plot, table(block))
block
 1 2 3
12 12 12
with(split_plot, table(irrigation))
irrigation
    moist saturated flooded
       12
                 12
                           12
with(split_plot, table(fertilizer))
fertilizer
    50 100 150
```


```
with (\verb|split_plot|, addmargins(table(irrigation, fertilizer)))\\
```

```
fertilizer
irrigation 0 50 100 150 Sum
 moist
           3 3
                 3
                    3 12
 saturated
           3 3
                    3 12
           3 3 3
 flooded
                    3 12
 Sum
             9
                 9
                    9 36
```


hist(split_plot\$y, main = 'Histogram')




boxplot(y ~ irrigation, data = split_plot)



boxplot(y ~ fertilizer, data = split_plot)





Model

```
mod.lme <- nlme::lme(
    y ~ irrigation * fertilizer,
    random = ~1 | block/irrigation,
    data = split_plot
)
res <- residuals(mod.lme, type = 'pearson')
anova(mod.lme)</pre>
```


```
    numDF
    denDF
    F-value
    p-value


    (Intercept)
    1
    18
    12.624677
    0.0023

    irrigation
    2
    4
    12.401598
    0.0193


    fertilizer
    3
    18
    22.922619
    <.0001</td>

    irrigation:fertilizer
    6
    18
    1.067825
    0.4172
```






Residuals VS Fitted



```
qqnorm(res, main = 'Normal Q-Q')
qqline(res, lty = 2, col = 'grey')
```



Residuals VS Exp. Units

