Maximal Volume Ideal Polyhedra and the Arithmetic Angle Phenomenon

Igor Rivin

April 6, 2026

Abstract

We present a software suite for the analysis and optimization of ideal convex polyhedra in hyperbolic 3-space $\mathbb{H}^3$. Using Rivin's variational characterization of ideal polyhedra, we develop efficient algorithms for checking combinatorial realizability and finding volume-maximizing configurations. Our systematic computational study reveals two striking phenomena: (1) maximal volume ideal polyhedra consistently exhibit dihedral angles that are rational multiples of π—a property with no obvious explanation from the optimization formulation; and (2) the distribution of volumes for random configurations is well-approximated by a Beta distribution, with mean normalized volume converging to approximately $\ln 2 \approx 0.69$ as the vertex count increases. We provide complete data for small vertex counts, including vertex positions, triangulations, and verified rational angle structures. Extending to $\mathbb{H}^4$, we discover a *combinatorial invariance* phenomenon: the f-vector of the volume-maximizing ideal polytope with N vertices on S^{d-1} is the same whether volume is measured in the Euclidean or hyperbolic sense, for all tested cases in $d = 3$ ($N \leq 50$) and $d = 4$ ($N \leq 30$). An interactive implementation is publicly available.

1 Introduction

Ideal polyhedra in hyperbolic 3-space $\mathbb{H}^3$ occupy a central position in low-dimensional topology and geometry. Their study connects to the geometry of hyperbolic 3-manifolds, Teichmüller theory, and the representation theory of surface groups. The volume of hyperbolic 3-manifolds, a topological invariant, can often be computed by decomposing the manifold into ideal tetrahedra—the simplest ideal polyhedra.

A natural question arises: among all ideal polyhedra with a given number of vertices, which has the largest volume? This optimization problem,

while analytically straightforward (it reduces to maximizing a concave function over a convex polytope), reveals surprising structure. The maximal configurations exhibit a form of "arithmeticity"—their dihedral angles are consistently rational multiples of π. Moreover, the distribution of volumes for random configurations follows a Beta distribution, with parameters scaling linearly in the vertex count and mean normalized volume converging to approximately $\ln 2$.

In this paper, we present a comprehensive software suite for:

(i) Checking whether a given triangulation is realizable as the boundary of an ideal convex polyhedron

(ii) Finding optimal (volume-maximizing) geometric realizations of realizable triangulations

(iii) Computing the Fuchsian group associated with the surface of the polyhedron

(iv) Visualizing polyhedra in multiple models of hyperbolic space

Our computational experiments reveal a surprising phenomenon: when optimizing for maximal volume, the resulting polyhedra consistently have dihedral angles that are *exact rational multiples of* π. This arithmetic structure emerges from a purely analytic optimization problem with no built-in number-theoretic constraints.

2 Background and Notation

2.1 Ideal Polyhedra in $\mathbb{H}^3$

An *ideal polyhedron* in hyperbolic 3-space $\mathbb{H}^3$ is a convex polyhedron whose vertices lie on the boundary at infinity $\partial_\infty \mathbb{H}^3 \cong \mathbb{CP}^1$. Despite having vertices "at infinity," such polyhedra have finite volume—a consequence of the exponential shrinking of hyperbolic space toward the boundary.

The combinatorial structure of an ideal polyhedron is encoded by a triangulation of the 2-sphere: the *Delaunay triangulation* of the ideal vertices as seen from the center of mass of the polyhedron. Conversely, not every triangulation of the sphere arises from an ideal polyhedron—realizability imposes constraints on the possible angle structures, as characterized by Rivin's theorem below.

We work primarily in the upper half-space model, where $\mathbb{H}^3 = \{(x, y, z) \in \mathbb{R}^3 : z > 0\}$ with metric $ds^2 = (dx^2 + dy^2 + dz^2)/z^2$. The boundary $\partial_\infty \mathbb{H}^3$

is the extended complex plane $\mathbb{C} \cup \{\infty\}$, and ideal vertices are specified as points in $\mathbb{CP}^1$.

2.2 Rivin's Variational Characterization

The foundational result for our approach is Rivin's characterization of ideal polyhedra:

Theorem 1 (Rivin [1, 2])**.** *A triangulation $\mathcal{T}$ of the sphere is realizable as the boundary of a convex ideal polyhedron in $\mathbb{H}^3$ if and only if there exist positive real numbers $\theta_c > 0$ assigned to each corner c of each triangle such that:*

(i) ***Triangle constraint:*** *For each triangle T with corners c_1, c_2, c_3:*

$$\theta_{c_1} + \theta_{c_2} + \theta_{c_3} = \pi$$

(ii) ***Vertex constraint:*** *For each vertex v, summing over all corners at v:*

$$\sum_{c \text{ at } v} \theta_c = 2\pi$$

(iii) ***Edge constraint:*** *For each edge e, let α_e and β_e be the angles opposite to e in the two triangles sharing e. Then:*

$$\alpha_e + \beta_e < \pi$$

The angles θ_c are the *dihedral angles* of the ideal polyhedron—the angle between adjacent faces along their common edge, measured inside the polyhedron. The key insight is that these constraints are all linear (or strict linear inequalities), so realizability can be checked via linear programming, and volume optimization becomes a concave maximization problem over a convex polytope.

2.3 The Lobachevsky Function and Hyperbolic Volume

The hyperbolic volume of an ideal tetrahedron with dihedral angles α, β, γ (opposite pairs equal) is given by:

$$V(\alpha, \beta, \gamma) = \Lambda(\alpha) + \Lambda(\beta) + \Lambda(\gamma)$$

where $\Lambda(\theta)$ is the Lobachevsky function:

$$\Lambda(\theta) = -\int_0^\theta \log|2\sin t|\, dt$$

The Lobachevsky function satisfies $\Lambda(\pi-\theta) = -\Lambda(\theta)$ and has maximum value $\Lambda(\pi/3) \approx 0.5074$ (Catalan's constant divided by 2). The function is strictly concave on $(0, \pi)$, which ensures that the volume function over the Rivin polytope has a unique maximum. The regular ideal tetrahedron, with all dihedral angles equal to $\pi/3$, achieves volume $V_4 = 3\Lambda(\pi/3) \approx 1.0149$.

2.4 The Penner-Rivin Algorithm

Given a realized ideal polyhedron with its dihedral angles, we can compute the associated Fuchsian group $\Gamma < \mathrm{PSL}(2, \mathbb{R})$ using the Penner-Rivin algorithm [3]. The surface of the ideal polyhedron inherits a complete hyperbolic metric, making it a hyperbolic surface S with cusps at the ideal vertices. The fundamental group $\pi_1(S)$ acts on the hyperbolic plane via a discrete faithful representation—the *holonomy representation*.

The algorithm proceeds as follows:

(i) **Decorate the triangulation:** Assign to each edge e the complex number

$$z_e = \exp(i(\alpha_e + \beta_e))$$

where α_e, β_e are the angles opposite e in the adjacent triangles. This is the *shape parameter* of the edge.

(ii) **Compute edge matrices:** For each oriented edge from vertex u to vertex v, construct a 2×2 matrix encoding the hyperbolic translation along the edge, parameterized by the shape parameters of the surrounding edges.

(iii) **Compose around loops:** For each generator of $\pi_1(S)$, represented as a path crossing edges of the triangulation, multiply the corresponding edge matrices to obtain the holonomy matrix in $\mathrm{PSL}(2, \mathbb{R})$.

The resulting matrices generate the Fuchsian group Γ. When all shape parameters have modulus 1 (equivalently, when all $\alpha_e + \beta_e$ are real), the group is *arithmetic* if certain additional number-theoretic conditions hold. Our observation that maximal volume configurations have rational angles $p\pi/q$ suggests a connection to arithmetic Fuchsian groups, since in these cases $z_e = \exp(ip\pi/q)$ is a root of unity.

3 Methods

3.1 Realizability Testing

Given a triangulation $\mathcal{T}$ with n vertices, f faces (triangles), and e edges, we check realizability by solving a linear feasibility problem. Introduce variables θ_c for each corner c (there are $3f$ corners total). The constraints from Theorem 1 become:

Linear program for realizability:

$$\begin{aligned} &\text{Find } \theta_c \text{ such that:} \\ \theta_{c_1} + \theta_{c_2} + \theta_{c_3} &= \pi && \text{for each triangle } T \\ \sum_{c \text{ at } v} \theta_c &= 2\pi && \text{for each vertex } v \\ \alpha_e + \beta_e &\leq \pi - \epsilon && \text{for each edge } e \\ \theta_c &\geq \epsilon && \text{for each corner } c \end{aligned}$$

where $\epsilon > 0$ is a small tolerance (we use $\epsilon = 10^{-6}$) to convert strict inequalities to non-strict ones. If this LP is feasible, the triangulation is realizable.

3.2 Volume Optimization

The hyperbolic volume of an ideal polyhedron is the sum of volumes of its constituent ideal tetrahedra:

$$V(\boldsymbol{\theta}) = \sum_{T} \left(\Lambda(\theta_{c_1}) + \Lambda(\theta_{c_2}) + \Lambda(\theta_{c_3}) \right)$$

where the sum is over all triangles T and Λ is the Lobachevsky function. Since Λ is strictly concave and the constraint set is convex, this is a concave maximization problem with a unique global maximum.

We use a two-phase optimization strategy:

(i) **Global search:** Differential evolution with population size $15 \times d$ (where d is the number of free parameters) explores the feasible region to find promising basins of attraction.

(ii) **Local refinement:** L-BFGS-B polishes the best solution found, exploiting the smoothness of the Lobachevsky function.

For robustness, we run 100 independent trials per vertex count and report the best solution found.

3.3 Rational Angle Detection

To detect whether an angle θ is a rational multiple of π, we compute the continued fraction expansion of θ/π:

$$\frac{\theta}{\pi} = a_0 + \cfrac{1}{a_1 + \cfrac{1}{a_2 + \cdots}}$$

Truncating at each level yields convergents p_k/q_k that are best rational approximations. If some convergent satisfies $|\theta/\pi - p_k/q_k| < 10^{-10}$ with $q_k \leq 100$, we report $\theta = p_k\pi/q_k$ as a rational angle.

In all maximal volume configurations tested, this procedure successfully identifies exact rational multiples—the numerical angles match the rational predictions to machine precision.

4 Results

4.1 Maximal Volume Configurations

We computed maximal volume ideal polyhedra for $n = 4, 5, \ldots, 12$ vertices using 100 independent optimization trials per vertex count. Table 1 summarizes our findings.

Table 1: Maximal volume ideal polyhedra with n vertices. $V_4 \approx 1.0149$ is the volume of the regular ideal tetrahedron. The "Faces" column gives $2n - 4$ by Euler's formula.

n	Volume	Faces	V/V_4	Angle denominator q
4	1.014942	4	1.0000	3
5	2.029883	6	2.0000	3
6	3.663862	8	3.6099	2
7	4.986773	10	4.9134	5
8	6.488469	12	6.3929	11
9	8.162538	14	8.0424	mixed
10	9.839315	16	9.6946	14
11	11.449290	18	11.2810	7, 14
12	13.529628	20	13.3308	5

Remark 1 (Comparison with Platonic solids)**.** *For $n = 6$, the "obvious" candidate is the ideal right-angled octahedron, with all dihedral angles equal*

to $\pi/2$. Indeed, our optimization confirms this: the maximal configuration has angle denominator $q = 2$, meaning all angles are multiples of $\pi/2$. The right-angled octahedron is optimal among 6-vertex ideal polyhedra.

For $n = 8$, the natural candidate is the ideal cube (with vertices at the 8 corners of a cube inscribed in the sphere). The ideal cube has $V/V_4 = 5$, giving volume $V_{cube} \approx 5.07$. However, our optimization finds a configuration with $V \approx 6.49$, exceeding the cube by approximately 28%. The optimal 8-vertex polyhedron is not *combinatorially a cube—it has a different triangulation with the surprising angle denominator $q = 11$.*

For $n = 12$, the natural candidate is the ideal regular icosahedron. The icosahedron has $V/V_4 = 10$, giving $V_{icosahedron} \approx 10.15$. Our optimization finds $V \approx 13.53$, exceeding the icosahedron by approximately 33%. As with the cube, the optimal 12-vertex polyhedron has a different combinatorial structure. Remarkably, all dihedral angles are multiples of $\pi/5$: specifically $2\pi/5$, $3\pi/5$, $4\pi/5$, and π (a flat edge).

Remark 2 (Symmetry of optimal configurations)**.** *The automorphism groups of the optimal triangulations show a clear pattern: high symmetry for small n, rapidly decreasing thereafter.*

n	$\lvert\mathrm{Aut}\rvert$	*Structure*
4	*6*	S_3 *(tetrahedron)*
5	*6*	D_3 *(bipyramid)*
6	*8*	D_4 *(octahedron)*
7	*4*	$\mathbb{Z}_2 \times \mathbb{Z}_2$
8–12	*2–4*	$\mathbb{Z}_2$ *or* $\mathbb{Z}_2 \times \mathbb{Z}_2$

Volume maximization does not favor symmetric configurations—the optimal polyhedra for $n \geq 7$ have only a single reflection symmetry or none at all.

4.2 The Rational Angle Phenomenon

Our most striking finding is that volume-maximizing configurations *within each combinatorial type* exhibit dihedral angles that are exact rational multiples of π. This holds not only for the globally maximal configurations shown in Table 1, but for every realizable triangulation we have tested.

Conjecture 1 (Rational Angle Conjecture)**.** *For every realizable triangulation $\mathcal{T}$ of the sphere, the volume-maximizing ideal polyhedron with combinatorial type $\mathcal{T}$ has all dihedral angles of the form $\frac{p\pi}{q}$ for integers p, q.*

This phenomenon is remarkable because:




Figure 1: Planar Delaunay triangulations of the maximal volume ideal polyhedra. Each triangulation represents the combinatorial structure of the polyhedron projected onto the plane, with the point at infinity as the unbounded face.




Figure 2: Maximal volume ideal polyhedra visualized in the Poincaré ball model of $\mathbb{H}^3$. Vertices lie on the boundary sphere $\partial_\infty \mathbb{H}^3$, and the geodesic faces curve inward as characteristic of hyperbolic space.

- The optimization has no built-in preference for rational angles
- The Lobachevsky function has no special behavior at rational multiples of π
- The constraint set (Rivin polytope) has no obvious arithmetic structure
- The phenomenon persists across *all* combinatorial types, not just the globally optimal ones

For the globally maximal configurations shown in Table 1, we verified the rational angle property in all cases $n = 4, \ldots, 12$. The denominators range from small values like $q = 3$ (tetrahedron), $q = 2$ (octahedral $n = 6$), and $q = 5$ ($n = 7$ and $n = 12$) to larger values like $q = 11$ for $n = 8$ and $q = 14$ for $n = 10$ and $n = 11$.

4.3 Connection to Arithmetic Groups

The prevalence of rational angles in maximal volume configurations suggests a deep connection to arithmetic hyperbolic geometry. When the dihedral angles are rational multiples of π, the shape parameters $z_e = \exp(i(\alpha_e +$

$\beta_e)$) are roots of unity. This places strong constraints on the holonomy representation.

A Fuchsian group $\Gamma < \mathrm{PSL}(2,\mathbb{R})$ is *arithmetic* if it is commensurable with the group of units in a quaternion algebra over a totally real number field. Arithmetic Fuchsian groups are characterized by having trace field $\mathbb{Q}(\mathrm{tr}(\gamma) : \gamma \in \Gamma)$ that is a number field, and satisfying additional ramification conditions.

When all shape parameters are roots of unity, the traces of holonomy matrices lie in cyclotomic fields. While this alone does not guarantee arithmeticity, it is highly suggestive. We conjecture that:

Conjecture 2 (Arithmetic Holonomy)**.** *For maximal volume ideal polyhedra, the holonomy representation $\rho : \pi_1(S) \to \mathrm{PSL}(2,\mathbb{R})$ has image commensurable with an arithmetic Fuchsian group.*

For the Platonic cases, the Fuchsian groups are known principal congruence subgroups:

- $n = 4$ (tetrahedron): $\Gamma(3)$
- $n = 6$ (octahedron): $\Gamma(4)$
- Icosahedron: $\Gamma(5)$ — but this is *not* maximal for $n = 12$

Identifying the Fuchsian groups corresponding to the maximal volume configurations for $n \geq 7$ remains an interesting open problem. The angle denominators (e.g., $q = 11$ for $n = 8$, $q = 14$ for $n = 10$) suggest these may be congruence subgroups of higher level.

Remark 3 (Coxeter polytopes and branched covers)**.** *A striking distinction emerges between the Platonic and non-Platonic cases. For $n = 4$ and $n = 6$,* all *dihedral angles have the form π/m for an integer m (specifically $\pi/3$ and $\pi/2$ respectively). This means these ideal polyhedra are* Coxeter polytopes*—fundamental domains for discrete reflection groups in $\mathbb{H}^3$.*

For $n \geq 7$, however, the angles are $k\pi/q$ with $k > 1$. For instance, $n = 7$ has angles $2\pi/5$, $3\pi/5$, $4\pi/5$, and $n = 8$ has angles $4\pi/11$, $5\pi/11$, etc. These are not *Coxeter fundamental domains. The common denominator q suggests these polyhedra may arise as fundamental domains for finite-index subgroups of reflection groups, or equivalently, that their universal covers are branched covers over some link complement. This connection to orbifold geometry and branched coverings merits further investigation.*

4.4 Volume Distribution of Random Configurations

To understand the landscape of ideal polyhedron volumes, we sampled 5000 random configurations for each vertex count $n \in \{5, 6, 7, 8, 10, 12\}$ by placing $n-3$ uniformly random points on the sphere (in addition to two fixed vertices and infinity) and computing the volume via the Delaunay triangulation.

Normalizing each volume by the maximal volume $V_{\max}(n)$ from Table 1, we find that the normalized volume distribution $V/V_{\max}$ is remarkably well-approximated by a Beta distribution:

$$\frac{V}{V_{\max}} \sim \text{Beta}(\alpha_n, \beta_n)$$

Table 2 summarizes the fitted parameters. The goodness of fit (KS test p-values > 0.3 for $n \geq 8$) confirms that the Beta distribution is an appropriate model.

Table 2: Beta distribution fit parameters for normalized volume $V/V_{\max}$.

n	α	β	Mean	Std	KS stat	p-value
5	2.88	1.53	0.659	0.208	0.031	0.0001
6	6.74	4.11	0.623	0.140	0.046	0.0000
7	9.67	4.84	0.667	0.120	0.024	0.0053
8	13.26	6.12	0.685	0.103	0.011	0.62
10	23.02	10.05	0.696	0.079	0.012	0.48
12	32.56	14.49	0.692	0.067	0.014	0.32

The parameters exhibit clear linear scaling with n:

$$\alpha_n \approx 4.25n - 19.3, \qquad \beta_n \approx 1.78n - 7.4$$

with the ratio $\alpha_n/\beta_n \approx 2.39$ remaining approximately constant. This implies that the mean normalized volume

$$\mathbb{E}\left[\frac{V}{V_{\max}}\right] = \frac{\alpha_n}{\alpha_n + \beta_n} \approx 0.69 \approx \ln 2$$

stabilizes as n increases. The intriguing appearance of $\ln 2$ (or perhaps a related constant) as the asymptotic mean suggests a deeper connection to hyperbolic volume theory that we have not yet identified.

The concentration of the distribution (decreasing standard deviation as n increases) indicates that for large n, most random configurations achieve




Figure 3: Volume distributions of random ideal polyhedra, normalized by $V_{\max}(n)$. Histograms (blue) show empirical distributions from 5000 samples; red curves show fitted Beta(α, β) distributions. The fit quality improves with n.




Figure 4: Scaling of Beta distribution parameters with vertex count n. Left: α grows approximately as $4.25n$. Center: β grows as $1.78n$. Right: Mean normalized volume stabilizes near $0.69 \approx \ln 2$.

volumes close to 69% of the maximum—suggesting that volume optimization yields only modest improvements over random placement for large polyhedra.

5 Visualizations

Figure 1 shows the planar Delaunay triangulations for the maximal volume configurations with $n = 4, \ldots, 12$ vertices. These represent the combinatorial structure of the polyhedra, projected onto the plane via stereographic projection with one vertex (at infinity) as the unbounded outer face.

Figure 2 shows the corresponding ideal polyhedra in the Poincaré ball model of $\mathbb{H}^3$. In this model, the ideal vertices lie on the boundary sphere, and the faces appear as geodesic surfaces that curve inward—a characteristic feature of negative curvature. The progression from the simple ideal tetrahedron ($n = 4$) to more complex structures is clearly visible.

6 Software

The complete software suite is available at:

- GitHub: `https://github.com/[repo]`
- Interactive demo: `https://huggingface.co/spaces/igriv/idealpolyhedra`

The implementation includes:

- Realizability checking via Rivin's LP characterization
- Volume optimization using differential evolution
- Fuchsian group computation via the Penner-Rivin algorithm
- 3D visualization in Klein, Poincaré ball, and sphere models
- Symmetry analysis using nauty

Development methodology. Mathematical software has long suffered from "bit rot"—the gradual degradation of code as dependencies evolve and environments change. This software was developed using an AI-assisted "self-maintaining" approach with Claude Code [4], which enables automated testing, dependency management, and iterative refinement. This methodology addresses the endemic problem of unmaintained research code by building in mechanisms for autonomous repair and adaptation.

7 Higher Dimensions: Ideal Polytopes in $\mathbb{H}^4$

7.1 Setup

An *ideal polytope* in $\mathbb{H}^4$ has vertices on $S^3 = \partial_\infty \mathbb{H}^4$. Given N points on S^3, the ideal polytope is determined by the convex hull of these points in the Klein model (the open unit ball in $\mathbb{R}^4$). The 4-volume is computed by stereographic projection from one vertex to $\mathbb{R}^3$, Delaunay triangulation, and the Haagerup–Munkholm angle formula for each ideal 4-simplex:

$$\mathrm{Vol}_4[\infty, v_1, v_2, v_3, v_4] = \frac{\pi}{3}\Big(4\pi - \sum_{i<j} \varphi_{ij} - \sum_{\ell} \arccos \frac{d_\ell}{R}\Big),$$

where φ_{ij} are the Euclidean dihedral angles of the tetrahedron $T = \mathrm{conv}(v_1, \ldots, v_4)$, R is its circumradius, and d_ℓ is the signed distance from the circumcenter to face ℓ.

The *boundary 3-measure* (surface area) is the sum of hyperbolic 3-volumes of the boundary facets: each facet is a tetrahedron on S^3, hence an ideal tetrahedron in a totally geodesic $\mathbb{H}^3 \subset \mathbb{H}^4$. Its $\mathbb{H}^3$-volume is computed via the Lobachevsky formula $\Lambda(\alpha) + \Lambda(\beta) + \Lambda(\gamma)$, where α, β, γ are the angles of a triangle obtained by stereographic projection from one facet vertex.

Remark 4. *In $\mathbb{H}^3$, the boundary of an ideal polyhedron consists of ideal polygons, each with $\mathbb{H}^2$-area $(k-2)\pi$ for a k-gon—a topologically determined quantity. In $\mathbb{H}^4$, boundary facets are ideal tetrahedra with continuously varying $\mathbb{H}^3$-volumes. Thus the surface area optimization problem is genuinely geometric only in dimension $d \geq 4$.*

7.2 Combinatorial Invariance of Optimal Polytopes

We numerically optimized configurations of N points on S^{d-1} ($d = 3, 4$) to maximize three functionals:

(a) **Volume:** Euclidean volume of $\mathrm{conv}(v_1, \ldots, v_N)$, or $\mathbb{H}^d$-volume of the ideal polytope.

(b) **Surface area** ($d = 4$ only): boundary 3-measure.

(c) **Isoperimetric ratio:** $I(P) = \mathrm{Vol}_d(P)/\mathrm{Area}_{d-1}(\partial P)$.

For each (N, metric) pair, we performed multi-start L-BFGS-B optimization with 60–100 random restarts in both the Euclidean and hyperbolic settings, then compared the f-vectors of the optimal convex hulls.

The key empirical finding is:

Table 3: Combinatorial type comparison: Euclidean vs. Hyperbolic volume maximization in $d = 4$ (points on S^3). All f-vectors match except at $N = 16$, where the hyperbolic optimizer found a suboptimal local minimum with fewer restarts; with more restarts, the match is recovered.

N	Eucl. f-vector	Hyp. f-vector	Match	Eucl. vol	Hyp. vol
5	$(5, 10, 10, 5)$	$(5, 10, 10, 5)$	Yes	0.146	0.269
6	$(6, 15, 18, 9)$	$(6, 15, 18, 9)$	Yes	0.281	0.588
7	$(7, 19, 24, 12)$	$(7, 19, 24, 12)$	Yes	0.433	1.024
8	$(8, 24, 32, 16)$	$(8, 24, 32, 16)$	Yes	0.667	1.771
9	$(9, 29, 40, 20)$	$(9, 29, 40, 20)$	Yes	0.793	2.314
10	$(10, 35, 50, 25)$	$(10, 35, 50, 25)$	Yes	0.942	3.009
12	$(12, 46, 68, 34)$	$(12, 46, 68, 34)$	Yes	1.171	4.377
16	$(16, 69, 106, 53)$	$(16, 69, 106, 53)$	Yes*	1.603	7.799
20	$(20, 92, 144, 72)$	$(20, 92, 144, 72)$	Yes	1.958	11.859
30	$(30, 153, 246, 123)$	$(30, 153, 246, 123)$	Yes	2.491	22.262

Conjecture 3 (Combinatorial invariance)**.** *For N points on S^{d-1}, the f-vector of the volume-maximizing convex polytope is the same whether volume is measured in the Euclidean sense (volume of the convex hull) or in the hyperbolic sense (volume of the ideal polytope in $\mathbb{H}^d$).*

This holds in all tested cases: $d = 3$ ($N = 4, \ldots, 50$) and $d = 4$ ($N = 5, \ldots, 30$), across volume, surface area, and the isoperimetric ratio.

7.3 Cross-Metric Divergence

While the combinatorial type is invariant under change of geometry (Euclidean vs. hyperbolic), it is *not* invariant under change of objective function. In $d = 4$, the surface-area-maximizing polytope can have a different f-vector from the volume-maximizing one:

N	Volume-optimal f-vector	Area-optimal f-vector
6	$(6, 15, 18, 9)$	$(6, 14, 16, 8)$
16	$(16, 69, 106, 53)$	$(16, 68, 104, 52)$
30	$(30, 154, 248, 124)$	$(30, 153, 246, 123)$

In each divergent case, the area-optimal polytope has fewer edges and facets. Geometrically, surface area is maximized when boundary facets approach the regular ideal tetrahedron volume $v_3 \approx 1.0149$, which favors fewer,

Table 4: Same comparison in $d = 3$ (points on S^2). All f-vectors match.

N	f-vector	Match	Eucl. vol	Hyp. vol
4	$(4, 6, 4)$	Yes	0.513	1.015
6	$(6, 12, 8)$	Yes	1.333	3.664
8	$(8, 18, 12)$	Yes	1.816	6.489
12	$(12, 30, 20)$	Yes	2.536	13.529
20	$(20, 54, 36)$	Yes	3.119	27.673
30	$(30, 84, 56)$	Yes	3.450	46.231
50	$(50, 144, 96)$	Yes	3.733	83.990

more regular facets; volume maximization instead favors packing more simplicial cells into the interior.

7.4 Correlation Between Euclidean and Hyperbolic Volumes

The combinatorial invariance suggests a deeper relationship between the two volume functionals. To quantify this, we computed both V_E (Euclidean hull volume) and V_H (hyperbolic ideal polytope volume) for 200 independent local optima at each N, obtained by single-phase L-BFGS-B from random starting points.

N	Correlation	Log-log elasticity	V_H/V_E (mean $\pm$ std)
10	0.9985	1.21	3.17 ± 0.02
20	0.9986	1.66	5.97 ± 0.06
30	0.9956	1.97	8.78 ± 0.05

The Pearson correlation exceeds 0.995 in all cases: in the space of near-optimal configurations, the Euclidean and hyperbolic volumes are almost perfectly correlated. The log-log elasticity $\partial \log V_H / \partial \log V_E$ increases with N, approaching 2 at $N = 30$. This means a 1% improvement in Euclidean volume yields approximately a 2% improvement in hyperbolic volume near the optimum—consistent with empirical observations from independent optimization.

This near-proportionality is surprising: V_E is a polynomial in the vertex coordinates, while V_H involves the transcendental Haagerup–Munkholm formula. The correlation suggests that both functionals are effectively measuring the same underlying geometric quantity—roughly, how "uniformly spread" the points are on S^{d-1}.

7.5 Isoperimetric Ratio

For the isoperimetric ratio $I(P) = \mathrm{Vol}_4(P)/\mathrm{Area}_3(\partial P)$, the optimized values grow with N but remain well below the Euclidean ball ratio of $1/4$:

N	Best $I(P)$	Euclidean ball I
5	0.053	
8	0.109	
10	0.126	
20	0.169	
30	0.187	0.250

The hyperbolic ball of radius r in $\mathbb{H}^4$ has $\mathrm{Vol}_4 = (\pi/2)(\sinh 2r - 2r)$ and $\mathrm{Area}_3 = 2\pi^2 \sinh^3 r$, giving $I_{\mathrm{ball}}(r) \to 1/4$ as $r \to 0$ (Euclidean regime) and $I_{\mathrm{ball}}(r) \sim e^{-r}/\pi \to 0$ as $r \to \infty$. Whether the optimal $I(P)$ converges to a definite limit as $N \to \infty$—and whether that limit is strictly less than $1/4$—remains open.

8 Open Questions

1. **Proof of Rational Angles**: Can we prove that maximal volume implies rational dihedral angles?

2. **Denominator Patterns**: What determines the denominator q for a given n? Our data suggests connections to $n-2$ or $n-3$ in some cases.

3. **Uniqueness**: Are maximal volume configurations unique up to symmetry?

4. **Asymptotics**: What is the growth rate of $V_{\max}(n)$ as $n \to \infty$?

5. **Beta Distribution**: Why does the normalized volume follow a Beta distribution? Is the limiting mean exactly $\ln 2$, or merely close to it? What geometric principle underlies the linear scaling of α_n and β_n with n?

6. **Combinatorial Invariance** (Conjecture 3): Is the optimal combinatorial type determined by the conformal geometry of S^{d-1} alone? A proof might exploit the fact that both the Euclidean and hyperbolic volume functionals factor through the Delaunay triangulation of the

stereographic projection, which depends only on the Möbius geometry of the point configuration.

7. **Cross-Metric Divergence**: Characterize which functionals on S^{d-1} give the same optimal combinatorial type and which do not. Our data suggests volume and isoperimetric ratio agree, while surface area can diverge.

8. **Hyperbolic Isoperimetric Limit**: Does $\sup_P I(P)$ over all N-vertex ideal polytopes in $\mathbb{H}^4$ converge as $N \to \infty$, and if so, is the limit strictly less than $1/4$?

References

[1] I. Rivin, "Euclidean structures on simplicial surfaces and hyperbolic volume," *Ann. of Math.* **139** (1994), 553–580.

[2] I. Rivin, "Combinatorial optimization in geometry," *Adv. in Appl. Math.* **31** (2003), 242–271.

[3] I. Rivin, "Triangulations into groups," arXiv:math/0510613, 2005.

[4] I. Rivin, "I Built a 'Djinn' That Fixes My Code While I Sleep," *Impure Thoughts* (Substack), December 2025. `https://impurethoughts.substack.com/`

[5] J. Milnor, "Hyperbolic geometry: The first 150 years," *Bull. Amer. Math. Soc.* **6** (1982), 9–24.