

MURILÉ BIOSCIENCE



PEPTIDE RECONSTITUTION & ADMINISTRATION GUIDE

Glow Smart. Live Well.

murilebioscience.netlify.app

WhatsApp: +63 945 986 2888

IMPORTANT MEDICAL DISCLAIMER

This guide is for general informational and educational purposes only. Murilé BioScience strongly recommends all peptide protocols be supervised by a qualified healthcare professional. This does not constitute medical advice and should not replace consultation with a licensed physician. Never self-diagnose or self-treat. Always consult your doctor before starting any new supplement or wellness protocol, especially if you have pre-existing medical conditions, are pregnant, breastfeeding, or taking medications. Results may vary.

By using this guide you acknowledge that:

- You are 18 years of age or older
- You understand peptides are general wellness supplements, not medicines
- You will consult a qualified healthcare professional before beginning any protocol
- Murilé BioScience is not liable for any adverse outcomes from improper use

STEP 1 — WHAT YOU WILL NEED

Gather all supplies before you begin. Having everything ready ensures a clean, safe and efficient process.

Your peptide vial	Refrigerate until 15 min before use
Bacteriostatic Water (Bac Water)	3ml or 10ml — do NOT substitute
Insulin syringe 31G 0.5ml	Thin, short — minimal discomfort
Alcohol swabs (70% isopropyl)	Wipe vial tops and injection site
Clean flat surface	Wipe with alcohol first
Sharps container	Never put needles in regular trash
Good lighting	Clearly read syringe markings
Gloves (optional)	For extra hygiene

TIP: Bacteriostatic Water contains benzyl alcohol to prevent bacterial growth, keeping your reconstituted peptide stable for 4–6 weeks. Never use regular water — it will degrade your peptide.

STEP 2 — PREPARING YOUR WORKSPACE

1. Wash hands with soap for at least 20 seconds
2. Wipe work surface with alcohol swab — let dry completely
3. Lay all supplies on the clean surface
4. Remove vials from fridge — allow 10–15 minutes to warm
5. Do not touch needle tips or vial stoppers after cleaning

STEP 3 — RECONSTITUTING YOUR PEPTIDE

Dissolving the freeze-dried peptide powder with bacteriostatic water. Always add water slowly along the SIDE of the vial — never directly onto the powder.

◆ Wipe both vial stoppers

Fresh alcohol swab on each. Let dry 10 seconds.

◆ Draw Bac Water into syringe

Insert into Bac Water vial. Pull plunger to draw required volume.

◆ Inject slowly along vial wall

Insert needle at 45°. Let water run down the SIDE of the vial.

◆ Swirl gently

Remove syringe. Rotate vial in circular motions. DO NOT SHAKE.

◆ Check clarity

Solution must be completely clear. If cloudy — discard.

HOW MUCH BAC WATER TO USE

Bac Water Added	Concentration	Typical Use
1ml (100 units)	Peptide mg ÷ 100 per unit	Lower-dose peptides
2ml (200 units)	Peptide mg ÷ 200 per unit	GHK-CU, Snap-8, skin peptides
3ml (300 units)	Peptide mg ÷ 300 per unit	Tirzepatide, NAD+, weight peptides

EXAMPLE: Tirzepatide 15mg + 3ml Bac Water = 0.05mg per unit. A 2.5mg starting dose = 50 units (pull syringe to the 50 mark).

STEP 4 — DRAWING YOUR DOSE

1. Wipe reconstituted vial stopper with fresh alcohol swab
2. Insert a FRESH insulin syringe into the vial
3. Invert the vial — pull plunger to draw your required dose
4. Check for air bubbles — tap syringe and push air out slowly
5. Confirm the dose on the syringe barrel

STEP 5 — SUBCUTANEOUS INJECTION

Most peptides are administered subcutaneously (under the skin). With a 31G insulin needle, most people feel only a mild pinch.

BEST INJECTION SITES

Site	Notes
Abdomen	2–3 inches from navel. Most recommended — less painful.
Outer Thigh	Upper outer thigh. Good abdomen alternative.
Upper Arm	Outer area only. Ask someone to assist.

INJECTION STEPS

1. Clean site with alcohol swab — let dry 10 seconds
2. Gently pinch a fold of skin with non-dominant hand
3. Hold syringe like a pencil at 45–90° angle
4. Insert needle swiftly and smoothly into the pinched skin
5. Release pinch — slowly press plunger to inject full dose
6. Remove needle smoothly — press lightly with clean swab
7. Dispose of used needle in sharps container IMMEDIATELY

CAUTION: Stop and seek medical advice if you experience severe pain, spreading redness, fever, or signs of allergic reaction (rash, difficulty breathing, facial swelling). Never reuse syringes.

STEP 6 — STORAGE & HANDLING

Type	Temperature	Duration	Notes
Powder (lyophilized)	2–8°C fridge	Up to 12 months	Away from light
Reconstituted	2–8°C fridge	4–6 weeks	Never freeze. Keep upright.
Room temperature	Max 4 hours	Discard after	Do not leave out overnight

STEP 7 — COMMON QUESTIONS

Q: Does it hurt?

With a 31G needle, most feel only a mild pinch. Let alcohol dry fully and rotate sites to reduce discomfort.

Q: What time of day?

NAD+ and Tirzepatide: morning. GHK-CU and recovery peptides: afternoon or before bed. Ask our team for specific guidance.

Q: What if I miss a dose?

Skip it and resume your normal schedule next day. Never double dose.

Q: Can I mix peptides?

Some can be stacked but only under healthcare professional guidance. Do not mix without proper advice.

Q: What if the solution is cloudy?

Discard immediately. A properly reconstituted peptide should be completely clear.

NEED HELP? WhatsApp: +63 945 986 2888 Website: murilebioscience.netlify.app TikTok: @muriletirz1986 | GCash: 09459862888 (Jemross Erida)