

ADULT AND CHILD IMMUNISATION SCHEDULE TASMANIA

(UPDATED OCTOBER 2011*)

AGE	VACCINES	NOTES
BIRTH	✓ Hep B (HB Vax II 0.5mL - paediatric)	Must be given within 7 days of birth.
2 MONTHS (MAY BE GIVEN FROM 6 WEEKS)	✓ DTPa/HIB/HepB/IPV (Infanrix Hexa) ✓ Pneumococcal 13vPCV (Prevenar 13) ✓ Rotavirus (Rotarix Oral Vaccine) *	* Rotarix administered orally. Refer to the 9th edition of the <i>Australian Immunisation Handbook</i> for additional information on interchangeability and age limits for rotavirus vaccines.
4 MONTHS	✓ DTPa/HIB/HepB/IPV (Infanrix Hexa) ✓ Pneumococcal 13vPCV (Prevenar13) ✓ Rotavirus (Rotarix Oral Vaccine)	
6 MONTHS	✓ DTPa/HIB/HepB/IPV (Infanrix Hexa) ✓ Pneumococcal 13vPCV (Prevenar13)	
6MTHS TO 64 YEARS AT MEDICAL RISK (INCLUDING PREGNANT WOMEN)	✓ Influenza (Fluvax*, Vaxigrip or Influvac) #	# for children under 10 years receiving influenza vaccine for the 1 st time a 2 nd dose is required 1 month after the 1 st dose * Fluvax is not for use in children under 5 years of age and is not recommended for children 5 -9 unless there is no alternative vaccine available. Refer to the 9th edition of <i>The Australian Immunisation Handbook</i> for further detail.
12 MONTHS	✓ Measles-Mumps-Rubella (Priorix) ✓ Meningococcal (NeisVacC or Meningitec) ✓ Hib (Hiberix) ✓ Pneumococcal Booster 13vPCV (Prevenar13) <i>predisposing medical conditions only</i>	
12 TO 35 MONTHS – SUPPLEMENTARY DOSE	✓ Pneumococcal 13vPCV (Prevenar13) #	# 1 dose for children who have received a full course of Prevenar 7 - 1 October 2011-30 September 2012 only
18 MONTHS	✓ Varicella (Varilrix or Varivax) +	*Only use if non-immune. Medical Practitioners should advise the Australian Childhood Immunisation Register (ACIR) in writing of a child's natural immunity.
3 – 4 YEARS (MEDICALLY AT RISK)	✓ Pneumococcal 13vPCV (Prevenar 13)# *	# Contact DHHS * Pneumococcal 13vPCV and Pneumococcal Booster 23vPPV must be given 8 weeks apart
4 YEARS (* MAY BE GIVEN FROM 3 YEARS 6 MONTHS)	✓ DTPa/IPV (Infanrix IPV or Quadracel)* ✓ Measles-Mumps-Rubella (Priorix)* ✓ Pneumococcal Booster 23vPPV (Pneumovax 23) <i>predisposing medical conditions only</i>	* May be given from 3 years and 6 months
SCHOOL YEAR 6 OR 7	✓ Varicella (Varilrix or Varivax) #	# <i>May be administered at GP surgeries if no prior history of varicella infection and previous dose not received at 18 months or via a school program.</i>
SCHOOL YEAR 6 OR 7	✓ Adult HepB (HB Vax II 1mL - adult) #	# 2 doses, 4 to 6 months apart. # <i>May be administered at GP surgeries if not received via a school program.</i>
SCHOOL YEAR 6 OR 7 GIRLS ONLY	✓ Human Papillomavirus (Gardasil) † #	† 3 doses - 0, 2 and 6 months # <i>May be administered at GP surgeries if not received via a school program.</i>
SCHOOL YEAR 10	✓ dTpa (Adacel or Boostrix) #	# <i>May be administered at GP surgeries if not received via a school program.</i>
50 YEARS AND OLDER	✓ Pneumococcal (Pneumovax 23) # ✓ dTpa (Adacel or Boostrix) *	# For medically at risk – recommended but not funded under the National Immunisation Program. * Recommended but not funded Refer to the 9th edition of the <i>Australian Immunisation Handbook</i> for further details
65 YEARS AND OLDER	✓ Pneumococcal (Pneumovax 23)* ✓ Influenza (Fluvax, Vaxigrip or Influvac)*	* Funded for people 65 year olds and older Refer to the 9th edition of the <i>Australian Immunisation Handbook</i> for further details

* This resource updated October 2011 check with your local Division of General Practice website for the latest version

Refer to the 9th edition of *The Australian Immunisation Handbook* for further details www.immunise.health.gov.au

ABORIGINAL AND TORRES STRAIT ISLANDER ADULT AND CHILD IMMUNISATION SCHEDULE TASMANIA

(UPDATED OCTOBER 2011*)

AGE	VACCINES	NOTES
BIRTH	✓ Hep B (HB Vax II 0.5mL - paediatric)	Must be given within 7 days of birth.
2 MONTHS (MAY BE GIVEN FROM 6 WEEKS)	✓ DTPa/HIB/HepB/IPV (Infanrix Hexa) ✓ Pneumococcal 13vPCV (Prevenar13) ✓ Rotavirus (Rotarix Oral Vaccine)*	* Rotarix administered orally. See the 9th edition of the <i>Australian Immunisation Handbook</i> for additional information on interchangeability and age limits for rotavirus vaccines.
4 MONTHS	✓ DTPa/HIB/HepB/IPV (Infanrix Hexa) ✓ Pneumococcal 13vPCV (Prevenar13) ✓ Rotavirus (Rotarix Oral Vaccine)	
6 MONTHS	✓ DTPa/HIB/HepB/IPV (Infanrix Hexa) ✓ Pneumococcal 13vPCV (Prevenar13)	
6 MONTHS TO 14 YEARS AT MEDICAL RISK	✓ Influenza (Fluvax*, Vaxigrip or Influvac) #	# for children under 10 years receiving influenza vaccine for the 1 st time a 2 nd dose is required 1 month after the 1 st dose * Fluvax is not for use in children under 5 years of age and is not recommended for children 5 -9 unless there is no alternative vaccine available. Refer to the 9th edition of <i>The Australian Immunisation Handbook</i> for further detail.
12 MONTHS	✓ Measles-Mumps-Rubella (Priorix) ✓ Meningococcal (NeisVacC or Meningitec) ✓ HIB (Hiberix) ✓ Pneumococcal Booster 13vPCV (Prevenar13) <i>predisposing medical conditions only</i>	
12 TO 35 MONTHS – SUPPLEMENTARY DOSE	✓ Pneumococcal 13vPCV (Prevenar13) #	# 1 dose for children who have received a full course of Prevenar 7 - 1 October 2011-30 September 2012 only
18 MONTHS	✓ Varicella (Varilrix or Varivax) +	+Only use if non-immune. Medical Practitioners should advise the Australian Childhood Immunisation Register (ACIR) in writing of a child's natural immunity.
3 – 4 YEARS (MEDICALLY AT RISK)	✓ Pneumococcal 13vPCV (Prevenar 13)# *	# Contact DHHS * Pneumococcal 13vPCV and Pneumococcal Booster 23vPPV must be given 8 weeks apart
4 YEARS (*MAY BE GIVEN FROM 3 YEARS 6 MONTHS)	✓ DTPa/IPV (Infanrix IPV or Quadracel)* ✓ Measles-Mumps-Rubella (Priorix)* ✓ Pneumococcal Booster (Pneumovax 23) <i>predisposing medical conditions only</i>	* May be given from 3 years and 6 months
SCHOOL YEAR 6 OR 7	✓ Varicella (Varilrix or Varivax) #	# <i>May be administered at GP surgeries if no prior history of varicella infection and previous dose not received at 18 months or via a school program.</i>
SCHOOL YEAR 6 OR 7	✓ Adult HepB (HB Vax II 1mL – adult) * #	*2 doses, 4 to 6 months apart. # <i>May be administered at GP surgeries if not received via a school program.</i>
SCHOOL YEAR 6 OR 7 GIRLS ONLY	✓ Human Papillomavirus (Gardasil) † #	† 3 doses - 0, 2 and 6 months after first dose. # <i>May be administered at GP surgeries if not received via a school program.</i>
SCHOOL YEAR 10	✓ dTpa (Adacel or Boostrix) #	# <i>May be administered at GP surgeries if not received via a school program.</i>
15 TO 49 YEARS AT MEDICAL RISK	✓ Pneumococcal (Pneumovax 23)*	* Funded from the age of 15 for people considered to have medical risk factors
15 YEARS AND OLDER	✓ Influenza (Fluvax, Vaxigrip or Influvac)	
50 YEARS AND OLDER	✓ Pneumococcal (Pneumovax 23)# ✓ dTpa (Adacel or Boostrix) *	# Funded from the age of 50 years for all Aboriginal and Torres Strait Islander People * Recommended but not funded Refer to the 9th edition of the <i>Australian Immunisation Handbook</i> for further details

* Immunisation Schedule for Aboriginal and Torres Strait Islander people living in other States and Territories may be different from above

This resource updated October 2011 check with your local Division of General Practice website for the latest version

** Refer to the 9th edition of *The Australian Immunisation Handbook* for further details
www.immunise.health.gov.au