The set of all affine combinations of points in some set $C \subseteq \mathbf{R}^n$ is called the *affine hull* of C, and denoted $\mathbf{aff}\, C$:

$$\mathbf{aff}\, C = \{\theta_1 x_1 + \cdots + \theta_k x_k \mid x_1, \ldots, x_k \in C,\ \theta_1 + \cdots + \theta_k = 1\}.$$

The affine hull is the smallest affine set that contains C, in the following sense: if S is any affine set with $C \subseteq S$, then $\mathbf{aff}\, C \subseteq S$.

2.1.3 Affine dimension and relative interior

We define the *affine dimension* of a set C as the dimension of its affine hull. Affine dimension is useful in the context of convex analysis and optimization, but is not always consistent with other definitions of dimension. As an example consider the unit circle in $\mathbf{R}^2$, *i.e.*, $\{x \in \mathbf{R}^2 \mid x_1^2 + x_2^2 = 1\}$. Its affine hull is all of $\mathbf{R}^2$, so its affine dimension is two. By most definitions of dimension, however, the unit circle in $\mathbf{R}^2$ has dimension one.

If the affine dimension of a set $C \subseteq \mathbf{R}^n$ is less than n, then the set lies in the affine set $\mathbf{aff}\, C \neq \mathbf{R}^n$. We define the *relative interior* of the set C, denoted $\mathbf{relint}\, C$, as its interior relative to $\mathbf{aff}\, C$:

$$\mathbf{relint}\, C = \{x \in C \mid B(x, r) \cap \mathbf{aff}\, C \subseteq C \text{ for some } r > 0\},$$

where $B(x, r) = \{y \mid \|y - x\| \leq r\}$, the ball of radius r and center x in the norm $\|\cdot\|$. (Here $\|\cdot\|$ is any norm; all norms define the same relative interior.) We can then define the *relative boundary* of a set C as $\mathbf{cl}\, C \setminus \mathbf{relint}\, C$, where $\mathbf{cl}\, C$ is the closure of C.

Example 2.2 Consider a square in the (x_1, x_2)-plane in $\mathbf{R}^3$, defined as

$$C = \{x \in \mathbf{R}^3 \mid -1 \leq x_1 \leq 1,\ -1 \leq x_2 \leq 1,\ x_3 = 0\}.$$

Its affine hull is the (x_1, x_2)-plane, *i.e.*, $\mathbf{aff}\, C = \{x \in \mathbf{R}^3 \mid x_3 = 0\}$. The interior of C is empty, but the relative interior is

$$\mathbf{relint}\, C = \{x \in \mathbf{R}^3 \mid -1 < x_1 < 1,\ -1 < x_2 < 1,\ x_3 = 0\}.$$

Its boundary (in $\mathbf{R}^3$) is itself; its relative boundary is the wire-frame outline,

$$\{x \in \mathbf{R}^3 \mid \max\{|x_1|, |x_2|\} = 1,\ x_3 = 0\}.$$

2.1.4 Convex sets

A set C is *convex* if the line segment between any two points in C lies in C, *i.e.*, if for any $x_1,\ x_2 \in C$ and any θ with $0 \leq \theta \leq 1$, we have

$$\theta x_1 + (1 - \theta) x_2 \in C.$$




Figure 2.2 Some simple convex and nonconvex sets. *Left.* The hexagon, which includes its boundary (shown darker), is convex. *Middle.* The kidney shaped set is not convex, since the line segment between the two points in the set shown as dots is not contained in the set. *Right.* The square contains some boundary points but not others, and is not convex.




Figure 2.3 The convex hulls of two sets in $\mathbf{R}^2$. *Left.* The convex hull of a set of fifteen points (shown as dots) is the pentagon (shown shaded). *Right.* The convex hull of the kidney shaped set in figure 2.2 is the shaded set.

Roughly speaking, a set is convex if every point in the set can be seen by every other point, along an unobstructed straight path between them, where unobstructed means lying in the set. Every affine set is also convex, since it contains the entire line between any two distinct points in it, and therefore also the line segment between the points. Figure 2.2 illustrates some simple convex and nonconvex sets in $\mathbf{R}^2$.

We call a point of the form $\theta_1 x_1 + \cdots + \theta_k x_k$, where $\theta_1 + \cdots + \theta_k = 1$ and $\theta_i \geq 0$, $i = 1, \ldots, k$, a *convex combination* of the points x_1, ..., x_k. As with affine sets, it can be shown that a set is convex if and only if it contains every convex combination of its points. A convex combination of points can be thought of as a *mixture* or *weighted average* of the points, with θ_i the fraction of x_i in the mixture.

The *convex hull* of a set C, denoted $\mathbf{conv}\, C$, is the set of all convex combinations of points in C:

$$\mathbf{conv}\, C = \{\theta_1 x_1 + \cdots + \theta_k x_k \mid x_i \in C,\ \theta_i \geq 0,\ i = 1, \ldots, k,\ \theta_1 + \cdots + \theta_k = 1\}.$$

As the name suggests, the convex hull $\mathbf{conv}\, C$ is always convex. It is the smallest convex set that contains C: If B is any convex set that contains C, then $\mathbf{conv}\, C \subseteq B$. Figure 2.3 illustrates the definition of convex hull.

The idea of a convex combination can be generalized to include infinite sums, integrals, and, in the most general form, probability distributions. Suppose $\theta_1, \theta_2, \ldots$

satisfy

$$\theta_i \geq 0, \quad i = 1, 2, \ldots, \qquad \sum_{i=1}^{\infty} \theta_i = 1,$$

and $x_1, x_2, \ldots \in C$, where $C \subseteq \mathbf{R}^n$ is convex. Then

$$\sum_{i=1}^{\infty} \theta_i x_i \in C,$$

if the series converges. More generally, suppose $p : \mathbf{R}^n \to \mathbf{R}$ satisfies $p(x) \geq 0$ for all $x \in C$ and $\int_C p(x)\, dx = 1$, where $C \subseteq \mathbf{R}^n$ is convex. Then

$$\int_C p(x) x \, dx \in C,$$

if the integral exists.

In the most general form, suppose $C \subseteq \mathbf{R}^n$ is convex and x is a random vector with $x \in C$ with probability one. Then $\mathbf{E}\, x \in C$. Indeed, this form includes all the others as special cases. For example, suppose the random variable x only takes on the two values x_1 and x_2, with $\mathbf{prob}(x = x_1) = \theta$ and $\mathbf{prob}(x = x_2) = 1 - \theta$, where $0 \leq \theta \leq 1$. Then $\mathbf{E}\, x = \theta x_1 + (1 - \theta) x_2$, and we are back to a simple convex combination of two points.

2.1.5 Cones

A set C is called a *cone*, or *nonnegative homogeneous*, if for every $x \in C$ and $\theta \geq 0$ we have $\theta x \in C$. A set C is a *convex cone* if it is convex and a cone, which means that for any $x_1,\ x_2 \in C$ and $\theta_1,\ \theta_2 \geq 0$, we have

$$\theta_1 x_1 + \theta_2 x_2 \in C.$$

Points of this form can be described geometrically as forming the two-dimensional pie slice with apex 0 and edges passing through x_1 and x_2. (See figure 2.4.)

A point of the form $\theta_1 x_1 + \cdots + \theta_k x_k$ with $\theta_1, \ldots, \theta_k \geq 0$ is called a *conic combination* (or a *nonnegative linear combination*) of $x_1, \ldots, x_k$. If x_i are in a convex cone C, then every conic combination of x_i is in C. Conversely, a set C is a convex cone if and only if it contains all conic combinations of its elements. Like convex (or affine) combinations, the idea of conic combination can be generalized to infinite sums and integrals.

The *conic hull* of a set C is the set of all conic combinations of points in C, *i.e.*,

$$\{\theta_1 x_1 + \cdots + \theta_k x_k \mid x_i \in C,\ \theta_i \geq 0,\ i = 1, \ldots, k\},$$

which is also the smallest convex cone that contains C (see figure 2.5).




Figure 2.4 The pie slice shows all points of the form $\theta_1 x_1 + \theta_2 x_2$, where $\theta_1,\ \theta_2 \geq 0$. The apex of the slice (which corresponds to $\theta_1 = \theta_2 = 0$) is at 0; its edges (which correspond to $\theta_1 = 0$ or $\theta_2 = 0$) pass through the points x_1 and x_2.




Figure 2.5 The conic hulls (shown shaded) of the two sets of figure 2.3.