19 Following the method of Problem 18, reduce A to echelon form and look at zero rows. The $\boldsymbol{b}$ column tells which combinations you have taken of the rows:

(a) $\begin{bmatrix} 1 & 2 & b_1 \\ 3 & 4 & b_2 \\ 4 & 6 & b_3 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 2 & b_1 \\ 2 & 3 & b_2 \\ 2 & 4 & b_3 \\ 2 & 5 & b_4 \end{bmatrix}$

From the $\boldsymbol{b}$ column after elimination, read off $m-r$ basis vectors in the left nullspace. Those y's are combinations of rows that give zero rows.

20 (a) Check that the solutions to $A\boldsymbol{x} = \boldsymbol{0}$ are perpendicular to the rows:

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 4 & 1 \end{bmatrix} \begin{bmatrix} 4 & 2 & 0 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} = ER.$$

(b) How many independent solutions to $A^{\mathrm{T}}\boldsymbol{y} = \boldsymbol{0}$? Why is $\boldsymbol{y}^{\mathrm{T}}$ the last row of E^{-1}?

21 Suppose A is the sum of two matrices of rank one: $A = \boldsymbol{u}\boldsymbol{v}^{\mathrm{T}} + \boldsymbol{w}\boldsymbol{z}^{\mathrm{T}}$.

(a) Which vectors span the column space of A?

(b) Which vectors span the row space of A?

(c) The rank is less than 2 if _____ or if _____.

(d) Compute A and its rank if $\boldsymbol{u} = \boldsymbol{z} = (1, 0, 0)$ and $\boldsymbol{v} = \boldsymbol{w} = (0, 0, 1)$.

22 Construct $A = \boldsymbol{u}\boldsymbol{v}^{\mathrm{T}} + \boldsymbol{w}\boldsymbol{z}^{\mathrm{T}}$ whose column space has basis $(1, 2, 4), (2, 2, 1)$ and whose row space has basis $(1, 0), (1, 1)$. Write A as (3 by 2) times (2 by 2).

23 Without multiplying matrices, find bases for the row and column spaces of A:

$$A = \begin{bmatrix} 1 & 2 \\ 4 & 5 \\ 2 & 7 \end{bmatrix} \begin{bmatrix} 3 & 0 & 3 \\ 1 & 1 & 2 \end{bmatrix}.$$

How do you know from these shapes that A cannot be invertible?

24 (Important) $A^{\mathrm{T}}\boldsymbol{y} = \boldsymbol{d}$ is solvable when $\boldsymbol{d}$ is in which of the four subspaces? The solution $\boldsymbol{y}$ is unique when the _____ contains only the zero vector.

25 True or false (with a reason or a counterexample):

(a) A and A^{T} have the same number of pivots.

(b) A and A^{T} have the same left nullspace.

(c) If the row space equals the column space then $A^{\mathrm{T}} = A$.

(d) If $A^{\mathrm{T}} = -A$ then the row space of A equals the column space.

26 (***Rank of AB***) If $AB = C$, the rows of C are combinations of the rows of _____. So the rank of C is not greater than the rank of _____. Since $B^{\mathrm{T}}A^{\mathrm{T}} = C^{\mathrm{T}}$, the rank of C is also not greater than the rank of _____.

27 If a, b, c are given with $a \neq 0$, how would you choose d so that $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ has rank 1? Find a basis for the row space and nullspace. Show they are perpendicular!

28 Find the ranks of the 8 by 8 checkerboard matrix B and the chess matrix C:

$$B = \begin{bmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} r & n & b & q & k & b & n & r \\ p & p & p & p & p & p & p & p \\ & & & \text{four zero rows} & & & & \\ p & p & p & p & p & p & p & p \\ r & n & b & q & k & b & n & r \end{bmatrix}$$

The numbers r, n, b, q, k, p are all different. Find bases for the row space and left nullspace of B and C. Challenge problem: Find a basis for the nullspace of C.

29 Can tic-tac-toe be completed (5 ones and 4 zeros in A) so that rank $(A) = 2$ but neither side passed up a winning move?

Challenge Problems

30 If $A = \boldsymbol{u}\boldsymbol{v}^{\mathrm{T}}$ is a 2 by 2 matrix of rank 1, redraw Figure 3.5 to show clearly the Four Fundamental Subspaces. If B produces those same four subspaces, what is the exact relation of B to A?

31 $\mathbf{M}$ is the space of 3 by 3 matrices. Multiply every matrix X in $\mathbf{M}$ by

$$A = \begin{bmatrix} 1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}. \quad \text{Notice: } A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

(a) Which matrices X lead to $AX =$ zero matrix?

(b) Which matrices have the form AX for some matrix X?

(a) finds the "nullspace" of that operation AX and (b) finds the "column space". What are the dimensions of those two subspaces of $\mathbf{M}$? Why do the dimensions add to $(n - r) + r = 9$?

32 Suppose the m by n matrices A and B have *the same four subspaces.* If they are both in row reduced echelon form, prove that F must equal G:

$$A = \begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix} \qquad B = \begin{bmatrix} I & G \\ 0 & 0 \end{bmatrix}.$$