■ REVIEW OF THE KEY IDEAS ■

1. Computer graphics needs shift operations $T(v) = v + v_0$ as well as linear operations $T(v) = Av$.
2. A shift in $\mathbf{R}^n$ can be executed by a matrix of order $n + 1$, using homogeneous coordinates.
3. The extra component 1 in $[x\ y\ z\ 1]$ is preserved when all matrices have the numbers $0, 0, 0, 1$ as last column.

Problem Set 8.7

1 A typical point in $\mathbf{R}^3$ is $xi + yj + zk$. The coordinate vectors i, j, and k are $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$. The coordinates of the point are (x, y, z).

This point in computer graphics is $xi + yj + zk +$ **origin**. Its homogeneous coordinates are (, , ,). Other coordinates for the same point are (, , ,).

2 A linear transformation T is determined when we know $T(i), T(j), T(k)$. For an affine transformation we also need $T($_____$)$. The input point $(x, y, z, 1)$ is transformed to $xT(i) + yT(j) + zT(k) +$ _____.

3 Multiply the 4 by 4 matrix T for translation along $(1, 4, 3)$ and the matrix T_1 for translation along $(0, 2, 5)$. The product TT_1 is translation along _____.

4 Write down the 4 by 4 matrix S that scales by a constant c. Multiply ST and also TS, where T is translation by $(1, 4, 3)$. To blow up the picture around the center point $(1, 4, 3)$, would you use vST or vTS?

5 What scaling matrix S (in homogeneous coordinates, so 3 by 3) would produce a 1 by 1 square page from a standard 8.5 by 11 page?

6 What 4 by 4 matrix would move a corner of a cube to the origin and then multiply all lengths by 2? The corner of the cube is originally at $(1, 1, 2)$.

7 When the three matrices in equation 1 multiply the unit vector a, show that they give $(\cos\theta)a$ and $(1 - \cos\theta)a$ and $\mathbf{0}$. Addition gives $aQ = a$ and the rotation axis is not moved.

8 If b is perpendicular to a, multiply by the three matrices in 1 to get $(\cos\theta)b$ and $\mathbf{0}$ and a vector perpendicular to b. So Qb makes an angle θ with b. ***This is rotation.***

9 What is the 3 by 3 projection matrix $I - nn^{\mathrm{T}}$ onto the plane $\frac{2}{3}x + \frac{2}{3}y + \frac{1}{3}z = 0$? In homogeneous coordinates add $0, 0, 0, 1$ as an extra row and column in P.

10 With the same 4 by 4 matrix P, multiply $T_- P T_+$ to find the projection matrix onto the plane $\frac{2}{3}x + \frac{2}{3}y + \frac{1}{3}z = 1$. The translation T_- moves a point on that plane (choose one) to $(0, 0, 0, 1)$. The inverse matrix T_+ moves it back.

11 Project $(3, 3, 3)$ onto those planes. Use P in Problem 9 and $T_- P T_+$ in Problem 10.

12 If you project a square onto a plane, what shape do you get?

13 If you project a cube onto a plane, what is the outline of the projection? Make the projection plane perpendicular to a diagonal of the cube.

14 The 3 by 3 mirror matrix that reflects through the plane $\boldsymbol{n}^{\mathrm{T}}\boldsymbol{v} = 0$ is $M = I - 2\boldsymbol{n}\boldsymbol{n}^{\mathrm{T}}$. Find the reflection of the point $(3, 3, 3)$ in the plane $\frac{2}{3}x + \frac{2}{3}y + \frac{1}{3}z = 0$.

15 Find the reflection of $(3, 3, 3)$ in the plane $\frac{2}{3}x + \frac{2}{3}y + \frac{1}{3}z = 1$. Take three steps $T_- M T_+$ using 4 by 4 matrices: translate by T_- so the plane goes through the origin, reflect the translated point $(3, 3, 3, 1)T_-$ in that plane, then translate back by T_+.

16 The vector between the origin $(0, 0, 0, 1)$ and the point $(x, y, z, 1)$ is the difference $v =$ _____. In homogeneous coordinates, vectors end in _____. So we add a _____ to a point, not a point to a point.

17 If you multiply only the *last* coordinate of each point to get (x, y, z, c), you rescale the whole space by the number _____. This is because the point (x, y, z, c) is the same as $(\ ,\ ,\ , 1)$.