We will count the individual multiplications, to see how much is saved. Before the **FFT** was invented, the count was the usual $n^2 = (1024)^2$. This is about a million multiplications. I am not saying that they take a long time. The cost becomes large when we have many, many transforms to do—which is typical. Then the saving by the FFT is also large:

The final count for size $n = 2^\ell$ is reduced from n^2 to $\frac{1}{2}n\ell$.

The number 1024 is 2^{10}, so $\ell = 10$. The original count of $(1024)^2$ is reduced to (5)(1024). The saving is a factor of 200. A million is reduced to five thousand. That is why the FFT has revolutionized signal processing.

Here is the reasoning behind $\frac{1}{2}n\ell$. There are ℓ levels, going from $n = 2^\ell$ down to $n = 1$. Each level has $n/2$ multiplications from the diagonal D's, to reassemble the half-size outputs from the lower level. This yields the final count $\frac{1}{2}n\ell$, which is $\frac{1}{2}n \log_2 n$.

One last note about this remarkable algorithm. There is an amazing rule for the order that the c's enter the FFT, after all the even-odd permutations. Write the numbers 0 to $n-1$ in binary (base 2). *Reverse the order of their digits.* The complete picture shows the bit-reversed order at the start, the $\ell = \log_2 n$ steps of the recursion, and the final output $y_0, \ldots, y_{n-1}$ which is F_n times $\boldsymbol{c}$.

The book ends with that very fundamental idea, a matrix multiplying a vector.

Thank you for studying linear algebra. I hope you enjoyed it, and I very much hope you will use it. It was a pleasure to write about this tremendously useful subject.

Problem Set 10.3

1 Multiply the three matrices in equation (3) and compare with F. In which six entries do you need to know that $i^2 = -1$?

2 Invert the three factors in equation (3) to find a fast factorization of F^{-1}.

3 F is symmetric. So transpose equation (3) to find a new Fast Fourier Transform!

4 All entries in the factorization of F_6 involve powers of $w_6 =$ sixth root of 1:

$$F_6 = \begin{bmatrix} I & D \\ I & -D \end{bmatrix} \begin{bmatrix} F_3 & \\ & F_3 \end{bmatrix} \begin{bmatrix} P \end{bmatrix}.$$

Write down these matrices with $1, w_6, w_6^2$ in D and $w_3 = w_6^2$ in F_3. Multiply!

5 If $\boldsymbol{v} = (1,0,0,0)$ and $\boldsymbol{w} = (1,1,1,1)$, show that $F\boldsymbol{v} = \boldsymbol{w}$ and $F\boldsymbol{w} = 4\boldsymbol{v}$. Therefore $F^{-1}\boldsymbol{w} = \boldsymbol{v}$ and $F^{-1}\boldsymbol{v} =$ _____.

6 What is F^2 and what is F^4 for the 4 by 4 Fourier matrix?

7 Put the vector $\boldsymbol{c} = (1,0,1,0)$ through the three steps of the FFT to find $\boldsymbol{y} = F\boldsymbol{c}$. Do the same for $\boldsymbol{c} = (0,1,0,1)$.

8 Compute $y = F_8 c$ by the three FFT steps for $c = (1,0,1,0,1,0,1,0)$. Repeat the computation for $c = (0,1,0,1,0,1,0,1)$.

9 If $w = e^{2\pi i/64}$ then w^2 and $\sqrt{w}$ are among the _____ and _____ roots of 1.

10 (a) Draw all the sixth roots of 1 on the unit circle. Prove they add to zero.

(b) What are the three cube roots of 1? Do they also add to zero?

11 The columns of the Fourier matrix F are the *eigenvectors* of the cyclic permutation P. Multiply PF to find the eigenvalues λ_1 to λ_4:

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & i^2 & i^3 \\ 1 & i^2 & i^4 & i^6 \\ 1 & i^3 & i^6 & i^9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & i^2 & i^3 \\ 1 & i^2 & i^4 & i^6 \\ 1 & i^3 & i^6 & i^9 \end{bmatrix} \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \lambda_3 & \\ & & & \lambda_4 \end{bmatrix}.$$

This is $PF = F\Lambda$ or $P = F\Lambda F^{-1}$. The eigenvector matrix (usually S) is F.

12 The equation $\det(P - \lambda I) = 0$ is $\lambda^4 = 1$. This shows again that the eigenvalue matrix Λ is _____. Which permutation P has eigenvalues = cube roots of 1?

13 (a) Two eigenvectors of C are $(1,1,1,1)$ and $(1,i,i^2,i^3)$. Find the eigenvalues.

$$\begin{bmatrix} c_0 & c_1 & c_2 & c_3 \\ c_3 & c_0 & c_1 & c_2 \\ c_2 & c_3 & c_0 & c_1 \\ c_1 & c_2 & c_3 & c_0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = e_1 \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \quad \text{and} \quad C \begin{bmatrix} 1 \\ i \\ i^2 \\ i^3 \end{bmatrix} = e_2 \begin{bmatrix} 1 \\ i \\ i^2 \\ i^3 \end{bmatrix}.$$

(b) $P = F\Lambda F^{-1}$ immediately gives $P^2 = F\Lambda^2 F^{-1}$ and $P^3 = F\Lambda^3 F^{-1}$. Then $C = c_0 I + c_1 P + c_2 P^2 + c_3 P^3 = F(c_0 I + c_1 \Lambda + c_2 \Lambda^2 + c_3 \Lambda^3) F^{-1} = FEF^{-1}$. That matrix E in parentheses is diagonal. It contains the _____ of C.

14 Find the eigenvalues of the "periodic" $-1, 2, -1$ matrix from $E = 2I - \Lambda - \Lambda^3$, with the eigenvalues of P in Λ. The -1's in the corners make this matrix periodic:

$$C = \begin{bmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{bmatrix} \quad \text{has } c_0 = 2, c_1 = -1, c_2 = 0, c_3 = -1.$$

15 ***Fast convolution.*** To multiply C times a vector x, we can multiply $F(E(F^{-1}x))$ instead. The direct way uses n^2 separate multiplications. Knowing E and F, the second way uses only $n \log_2 n + n$ multiplications. How many of those come from E, how many from F, and how many from F^{-1}?

16 Why is row i of $\overline{F}$ the same as row $N - i$ of F (numbered 0 to $N - 1$)?