PART
II

Structure of Groups

Section 8 Groups of Permutations
Section 9 Finitely Generated Abelian Groups
Section 10 Cosets and the Theorem of Lagrange
Section 11 Plane Isometries

SECTION 8 GROUPS OF PERMUTATIONS

Let $\phi : G \to G'$ be a function mapping the group G to G'. Recall that the homomorphism property of an isomorphism states that for all $a, b \in G$, $\phi(ab) = \phi(a)\phi(b)$. Whenever a function has this property whether or not the function is one-to-one or onto, we say that ϕ is a **group homomorphism**. Of course any group isomorphism is a group homomorphism, but the reverse is not necessarily true.

8.1 Definition Let G and G' be groups with $\phi : G \to G'$. The map ϕ is a **homomorphism** if the homomorphism property

$$\phi(ab) = \phi(a)\phi(b)$$

holds for all $a, b \in G$. ■

8.2 Example Let $\phi : \mathbb{R} \to U$ (the circle group) be defined by the formula

$$\phi(x) = \cos(2\pi x) + i \sin(2\pi x) = e^{2\pi i x}.$$

Then

$$\phi(a + b) = \cos(2\pi(a + b)) + i \sin(2\pi(a + b)) = e^{2\pi i(a+b)}.$$

Using either the usual properties of the exponential function or the formulas from trigonometry involving the sum of two angles, we see that

$$\phi(a + b) = (\cos(2\pi a) + i \sin(2\pi a))(\cos(2\pi b) + i \sin(2\pi b)) = e^{2\pi a i} e^{2\pi b i},$$

so

$$\phi(a + b) = \phi(a)\phi(b),$$

which says that ϕ is a group homomorphism. Although ϕ maps onto U, it is not one-to-one, so ϕ is not an isomorphism.

The identity $0 \in \mathbb{R}$ maps to 1, the identity in U. Furthermore, for any $x \in \mathbb{R}$,

$$\phi(-x) = e^{-2\pi i x} = \frac{1}{e^{2\pi i x}} = (\phi(x))^{-1}.$$

▲

8.3 Example Recall that $U_n = \{z \in \mathbb{C} \mid z^n = 1\}$. Let $\phi : U_{28} \to U_4$ be given by $\phi(z) = z^7$. To check that ϕ is well defined, we see that if $z \in U_{28}$, then $z^{28} = 1$. Therefore, $(z^7)^4 = 1$, which implies that $z^7 \in U_4$. We check that ϕ is a homomorphism.

$$\phi(z_1 z_2) = (z_1 z_2)^7 = z_1^7 z_2^7 = \phi(z_1)\phi(z_2).$$

As in the previous example, ϕ maps the identity in U_{28}, in this case 1, to the identity 1 in U_4. Furthermore,

$$\phi(z^{-1}) = z^{-7} = (z^7)^{-1} = (\phi(z))^{-1}. \quad \blacktriangle$$

8.4 Definition Let $\phi : X \to Y$ and suppose that $A \subseteq X$ and $B \subseteq Y$. The set $\phi[A] = \{\phi(a) \mid a \in A\}$ is called the **image of A in Y under the mapping ϕ.** The set $\phi^{-1}[B] = \{a \in A \mid \phi(a) \in B\}$ is called the **inverse image of B under the mapping ϕ.** ■

The four properties of a homomorphism given in the theorem that follows are obvious in the case of an isomorphism since we think of an isomorphism as simply relabeling the elements of a group. However, it is not obvious that these properties hold for all homomorphisms whether or not they are one-to-one or onto maps. Consequently, we give careful proofs of all four properties.

8.5 Theorem Let ϕ be a homomorphism of a group G into a group G'.

1. If e is the identity element in G, then $\phi(e)$ is the identity element e' in G'.
2. If $a \in G$, then $\phi(a^{-1}) = \phi(a)^{-1}$.
3. If H is a subgroup of G, then $\phi[H]$ is a subgroup of G'.
4. If K' is a subgroup of G', then $\phi^{-1}[K']$ is a subgroup of G.

Loosely speaking, ϕ preserves the identity element, inverses, and subgroups.

Proof Let ϕ be a homomorphism of G into G'. Then

$$\phi(e) = \phi(ee) = \phi(e)\phi(e).$$

Multiplying on the left by $\phi(e)^{-1}$, we see that $e' = \phi(e)$. Thus $\phi(e)$ must be the identity element e' in G'. The equation

$$e' = \phi(e) = \phi(aa^{-1}) = \phi(a)\phi(a^{-1})$$

shows that $\phi(a^{-1}) = \phi(a)^{-1}$ for all $a \in G$.

Turning to Statement (3), let H be a subgroup of G, and let $\phi(a)$ and $\phi(b)$ be any two elements in $\phi[H]$. Then $\phi(a)\phi(b) = \phi(ab)$, so we see that $\phi(a)\phi(b) \in \phi[H]$; thus, $\phi[H]$ is closed under the operation of G'. The fact that $e' = \phi(e)$ and $\phi(a^{-1}) = \phi(a)^{-1}$ completes the proof that $\phi[H]$ is a subgroup of G'.

Going the other way for Statement (4), let K' be a subgroup of G'. Suppose a and b are in $\phi^{-1}[K']$. Then $\phi(a)\phi(b) \in K'$ since K' is a subgroup. The equation $\phi(ab) = \phi(a)\phi(b)$ shows that $ab \in \phi^{-1}[K']$. Thus $\phi^{-1}[K']$ is closed under the binary operation in G. Also, K' must contain the identity element $e' = \phi(e)$, so $e \in \phi^{-1}[K']$. If $a \in \phi^{-1}[K']$, then $\phi(a) \in K'$, so $\phi(a)^{-1} \in K'$. But $\phi(a)^{-1} = \phi(a^{-1})$, so we must have $a^{-1} \in \phi^{-1}[K']$. Hence $\phi^{-1}[K']$ is a subgroup of G. ◆

Let $\phi : G \to G'$ be a homomorphism and let e' be the identity element of G'. Now $\{e'\}$ is a subgroup of G', so $\phi^{-1}[\{e'\}]$ is a subgroup H of G by Statement (4) in Theorem 8.5. This subgroup is critical to the study of homomorphisms.

8.6 Definition Let $\phi : G \to G'$ be a homomorphism of groups. The subgroup $\phi^{-1}[\{e'\}] = \{x \in G \mid \phi(x) = e'\}$ is the **kernel of ϕ**, denoted by $\text{Ker}(\phi)$. ■

We will use the kernel of a homomorphism when we define the alternating group later in this section.

Another extreme is to let $H = G$ in Statement (3) of Theorem 8.5. In this case, the theorem says that $\phi[G]$ is a subgroup of G'. We use this in the proof of Cayley's Theorem.

8.7 Example In Example 8.2, the homomorphism $\phi : \mathbb{R} \to U$ is defined by $\phi(x) = \cos(2\pi x) + i\sin(2\pi x) = e^{2\pi ix}$. The kernel of ϕ is the set of integers since $\cos(2\pi x) + i\sin(2\pi x) = 1$ if and only if x is an integer.

Let n be a positive integer. Then $\left\langle \frac{1}{n} \right\rangle$ is a subgroup of $\mathbb{R}$ and

$$\begin{aligned}\phi\left[\left\langle \frac{1}{n} \right\rangle\right] &= \phi\left[\left\{ \frac{m}{n} \mid m \in \mathbb{Z} \right\}\right] \\ &= \{\cos(2\pi m/n + i\sin(2\pi m/n)) \mid m \in \mathbb{Z}\} \\ &= U_n.\end{aligned}$$

▲

8.8 Example Let $\phi : \mathbb{Z}_n \to D_n$ be given by $\phi(k) = \rho^k$. We check that ϕ is a homomorphism. Let $a, b \in \mathbb{Z}_n$. If $a + b < n$, then $a +_n b = a + b$, so $\phi(a +_n b) = \phi(a + b) = \rho^{a+b} = \rho^a\rho^b = \phi(a)\phi(b)$. If $a + b \geq n$, then $\phi(a +_n b) = \phi(a + b - n) = \rho^{a+b-n} = \rho^a\rho^b\rho^{-n} = \rho^a\rho^b = \phi(a)\phi(b)$. The image $\phi[\mathbb{Z}_n]$ is $\langle\rho\rangle$. ▲

Cayley's Theorem

Each of the groups we have seen so far is isomorphic to a subgroup of permutations on some set. For example, $\mathbb{Z}_n$ is isomorphic with the cyclic group $\langle(1, 2, 3, \ldots, n)\rangle \leq S_n$. The dihedral group D_n is defined to be the permutations in $S_{\mathbb{Z}_n}$ with the property that the line segment between vertices i and j is an edge in P_n, a regular n-gon, if and only if the line segment between the images of i and j is also an edge. The infinite group $\mathrm{GL}(n, \mathbb{R})$ can be thought of as invertible linear transformations of $\mathbb{R}^n$. Each element of $\mathrm{GL}(n, \mathbb{R})$ permutes the vectors in $\mathbb{R}^n$, which makes $\mathrm{GL}(n, \mathbb{R})$ isomorphic with a permutation group on vectors in $\mathbb{R}^n$. We refer to a subgroup of a permutation group as a **group of permutations**. Cayley's Theorem states that any group is isomorphic with a group of permutations.

At first Cayley's Theorem seems like a remarkable result that could be used to understand all groups. In fact, this is a nice and intriguing classic result. Unfortunately, approaching group theory by trying to determine all possible permutation groups is not feasible. On the other hand, Cayley's theorem does show the strength and generality of permutation groups and it deserves a special place in group theory for that reason. For example, if we wish to find a counterexample to a conjecture about groups, provided that there is one, it will occur in a permutation group.

It may seem a mystery how we could start with an arbitrary group and come up with a permutation group that is isomorphic with the given group. The key is to think about the group table. Each row contains each element of the group exactly once. So each row defines a permutation of the elements of the group by placing the table head as the top row in the two-row representation of a permutation and placing the row corresponding to an element a in the group as the bottom row. Table 8.9 is the group table for D_3. Note that the permutation obtained using the row $\mu\rho$ is

$$\begin{pmatrix} \iota & \rho & \rho^2 & \mu & \mu\rho & \mu\rho^2 \\ \mu\rho & \mu\rho^2 & \mu & \rho^2 & \iota & \rho \end{pmatrix}.$$

8.9 Table

D_3	ι	ρ	ρ^2	μ	$\mu\rho$	$\mu\rho^2$
ι	ι	ρ	ρ^2	μ	$\mu\rho$	$\mu\rho^2$
ρ	ρ	ρ^2	ι	$\mu\rho^2$	μ	$\mu\rho$
ρ^2	ρ^2	ι	ρ	$\mu\rho$	$\mu\rho^2$	μ
μ	μ	$\mu\rho$	$\mu\rho^2$	ι	ρ	ρ^2
$\mu\rho$	$\mu\rho$	$\mu\rho^2$	μ	ρ^2	ι	ρ
$\mu\rho^2$	$\mu\rho^2$	μ	$\mu\rho$	ρ	ρ^2	ι

All that remains to prove Cayley's Theorem, at least when the group is finite, is to check that the permutations obtained from the group table form a group isomorphism with the original group. Let λ_x be the permutation of the elements of G given by the x row of the table for G. Then for any $g \in G$, $\lambda_x(g)$ is the entry in the x row and g column of the group table. In other words, $\lambda_x(g) = xg$, which is perfectly valid in the case of an infinite as well as a finite group. We formalize this connection between G and permutations on G in Definition 8.10.

8.10 Definition Let G be a group. The function $\phi : G \to S_G$ given by $\phi(x) = \lambda_x$ where $\lambda_x(g) = xg$ for all $g \in G$ is called the **left regular representation** of G. ■

In order to be sure that λ_x is a permutation, it should be verified that λ_x is both one-to-one and onto. We see that λ_x is one-to-one since if $\lambda_x(a) = \lambda_x(b)$, $xa = xb$ and cancellation gives $a = b$. Also, λ_x maps onto G because for any $b \in G$, $\lambda_x(x^{-1}b)) = b$. We are now ready to prove Cayley's Theorem.

8.11 Theorem **(Cayley's Theorem)** Every group is isomorphic to a group of permutations.

Proof Let G be a group. The left regular representation provides a map $\phi : G \to S_G$ defined by $\phi(x) = \lambda_x$. We must verify that ϕ is a group homomorphism and that ϕ is one-to-one. Then $\phi[G]$ is a subgroup of S_G by Theorem 8.5 and $\phi : G \to \phi[G]$ is an isomorphism.

We first show that ϕ is one-to-one. Suppose that $a, b \in G$ and $\phi(a) = \phi(b)$. Then the permutations λ_a and λ_b are the same, so $\lambda_a(e) = \lambda_b(e)$. Thus $ae = be$ and $a = b$. So ϕ is one-to-one.

We now need to show that ϕ is a group homomorphism. Let $a, b \in G$. Then $\phi(ab) = \lambda_{ab}$ and $\phi_a\phi_b = \lambda_a\lambda_b$. We must show that the two permutations λ_{ab} and $\lambda_a\lambda_b$ are the same. Let $g \in G$.

$$\lambda_{ab}(g) = (ab)g = a(bg) = \lambda_a(bg) = \lambda_a(\lambda_b(g)) = (\lambda_a\lambda_b)(g).$$

Thus $\lambda_{ab} = \lambda_a\lambda_b$, which implies that $\phi(ab) = \phi(a)\phi(b)$. So ϕ is a one-to-one homomorphism, which completes the proof. ◆

8.12 Example The proof of Cayley's Theorem shows that any group G is isomorphic with a subgroup of S_G, but this is typically not the smallest symmetric group that has a subgroup isomorphic with G. For example, D_n is isomorphic with a subgroup of $S_{\mathbb{Z}_n}$ while the proof of Cayley's Theorem gives a subgroup of S_{D_n} and D_n has $2n$ elements while $\mathbb{Z}_n$ has only n elements. On the surface, it may seem that $\mathbb{Z}_6$ cannot be isomorphic with a subgroup of S_n for $n < 6$, but $(1, 2, 3)(4, 5) \in S_5$ generates a subgroup isomorphic with $\mathbb{Z}_6$. ▲

We defined the left regular representation in Definition 8.10. We now define the right regular representation. Instead of λ_x representing the row for x in the group table, we use σ_x to represent the column with head x. Instead of using ϕ for the function that sends x to λ_x, we use τ, which sends x to $\sigma_{x^{-1}}$.

HISTORICAL NOTE

Arthur Cayley (1821–1895) gave an abstract-sounding definition of a group in a paper of 1854: "A set of symbols, $1, \alpha, \beta, \cdots$, all of them different and such that the product of any two of them (no matter in what order) or the product of any one of them into itself, belongs to the set, is said to be a group." He then proceeded to define a group table and note that every line and column of the table "will contain all the symbols $1, \alpha, \beta, \cdots$." Cayley's symbols, however, always represented operations on sets; it does not seem that he was aware of any other kind of group. He noted, for instance, that the four matrix operations $1, \alpha =$ inversion, $\beta =$ transposition, and $\gamma = \alpha\beta$, form, abstractly, the non-cyclic group of four elements. In any case, his definition went unnoticed for a quarter of a century.

This paper of 1854 was one of about 300 written during the 14 years Cayley was practicing law, being unable to find a suitable teaching post. In 1863, he finally became a professor at Cambridge. In 1878, he returned to the theory of groups by publishing four papers, in one of which he stated Theorem 8.11 of this text; his "proof" was simply to notice from the group table that multiplication by any group element permuted the group elements. However, he wrote, "this does not in any wise show that the best or the easiest mode of treating the general problem [of finding all groups of a given order] is thus to regard it as a problem of [permutations]. It seems clear that the better course is to consider the general problem in itself."

The papers of 1878, unlike the earlier one, found a receptive audience; in fact, they were an important influence on Walther von Dyck's 1882 axiomatic definition of an abstract group, the definition that led to the development of abstract group theory.

8.13 Definition Let G be a group. The map $\tau : G \to S_G$ given by $\tau(x) = \sigma_{x^{-1}}$ where $\sigma_x(g) = gx$ is called the **right regular representation** of G. ■

We could have used the right regular representation to prove Cayley's Theorem instead of the left regular representation. Exercise 54 asks for the details of the proof.

Even and Odd Permutations

It seems reasonable that every reordering of the sequence $1, 2, \ldots, n$ can be achieved by repeated interchange of positions of pairs of numbers. We discuss this a bit more formally.

8.14 Definition A cycle of length 2 is a **transposition**. ■

Thus a transposition leaves all elements but two fixed, and maps each of these onto the other. A computation shows that

$$(a_1, a_2, \cdots, a_n) = (a_1, a_n)(a_1, a_{n-1}) \cdots (a_1, a_3)(a_1, a_2).$$

Therefore any cycle of length n can be written as a product of $n - 1$ transpositions. Since any permutation of a finite set can be written as a product of cycles, we have the following.

8.15 Theorem Any permutation of a finite set containing at least two elements is a product of transpositions. ♦

Naively, this theorem just states that any rearrangement of n objects can be achieved by successively interchanging pairs of them.

8.16 Example Following the remarks prior to the theorem, we see that (1, 6) (2, 5, 3) is the product (1, 6) (2, 3) (2, 5) of transpositions. ▲

8.17 Example In S_n for $n \geq 2$, the identity permutation is the product (1, 2) (1, 2) of transpositions. ▲

We have seen that every permutation of a finite set with at least two elements is a product of transpositions. The transpositions may not be disjoint, and a representation of the permutation in this way is not unique. For example, we can always insert at the beginning the transposition (1, 2) twice, because (1, 2) (1, 2) is the identity permutation. What is true is that the number of transpositions used to represent a given permutation must either always be even or always be odd. This is an important fact. The proof involves counting orbits and was suggested by David M. Bloom.

Let $\sigma \in S_A$ and $a \in A$. We let the **orbit** of a be the set $\{\sigma^k(a) \mid k \in \mathbb{Z}\}$. In the case of $\sigma \in S_n$, a simple way to think of the orbit of a is to think of the elements in the cycle containing a in the disjoint cycle representation of σ.

8.18 Example Let $\sigma = (1, 2, 6)(3, 5) \in S_6$. Then the orbit of 1 is the set $\{1, 2, 6\}$, which is also the orbit of 2 and the orbit of 6. The set $\{3, 5\}$ is the orbit of 3 and the orbit of 5. What about the orbit of 4? Recall that if we include 1-cycles, $\sigma = (1, 2, 6)(3, 5)(4)$, which says the orbit of 4 is $\{4\}$. ▲


8.19 Theorem No permutation in S_n can be expressed both as a product of an even number of transpositions and as a product of an odd number of transpositions.

Proof Let $\sigma \in S_n$ and let $\tau = (i, j)$ be a transposition in S_n. We claim that the number of orbits of σ and of $\tau\sigma$ differ by 1.

Case I Suppose i and j are in different orbits of σ. Write σ as a product of disjoint cycles, the first of which contains j and the second of which contains i, symbolized by the two circles in Fig. 8.20. We may write the product of these two cycles symbolically as

$$(b, j, \times, \times, \times)(a, i, \times, \times)$$

where the symbols $\times$ denote possible other elements in these orbits.



8.20 Figure


Computing the product of the first three cycles in $\tau\sigma = (i, j)\sigma$, we obtain

$$(i, j)(b, j, \times, \times, \times)(a, i, \times, \times) = (a, j, \times, \times, \times, b, i, \times, \times).$$

The original 2 orbits have been joined to form just one in $\tau\sigma$ as symbolized in Fig. 8.20. Exercise 42 asks us to repeat the computation to show that the same thing happens if either one or both of i and j should be the only element of their orbit in σ.

Case II Suppose i and j are in the same orbit of σ. We can then write σ as a product of disjoint cycles with the first cycle of the form

$$(a, i, \times, \times, \times, b, j, \times, \times)$$



8.21 Figure

shown symbolically by the circle in Fig. 8.20. Computing the product of the first two cycles in $\tau\sigma = (i,j)\sigma$, we obtain

$$(i,j)(a,i,\times,\times,\times,b,j,\times,\times) = (a,j,\times,\times)(b,i,\times,\times,\times).$$

The original single orbit has been split into two as symbolized in Fig. 8.21.

We have shown that the number of orbits of $\tau\sigma$ differs from the number of orbits of σ by 1. The identity permutation ι has n orbits, because each element is the only member of its orbit. Now the number of orbits of a given permutation $\sigma \in S_n$ differs from n by either an even or an odd number, but not both. Thus it is impossible to write

$$\sigma = \tau_1\tau_2\tau_3\cdots\tau_m\iota$$

where the τ_k are transpositions in two ways, once with m even and once with m odd. ◆

8.22 Definition A permutation of a finite set is **even** or **odd** according to whether it can be expressed as a product of an even number of transpositions or the product of an odd number of transpositions, respectively. ■

8.23 Example The identity permutation ι in S_n is an even permutation since we have $\iota = (1,2)(1,2)$. If $n = 1$ so that we cannot form this product, we define ι to be even. On the other hand, the permutation (1, 4, 5, 6) (2, 1, 5) in S_6 can be written as

$$(1,4,5,6)(2,1,5) = (1,6)(1,5)(1,4)(2,5)(2,1)$$

which has five transpositions, so this is an odd permutation. ▲

The Alternating Groups

We claim that for $n \geq 2$, the number of even permutations in S_n is the same as the number of odd permutations; that is, S_n is split equally and both numbers are $(n!)/2$. To show this, let A_n be the set of even permutations in S_n and let B_n be the set of odd permutations for $n \geq 2$. We proceed to define a one-to-one function from A_n onto B_n. This is exactly what is needed to show that A_n and B_n have the same number of elements.

Let τ be any fixed transposition in S_n; it exists since $n \geq 2$. We may as well suppose that $\tau = (1,2)$. We define a function

$$\lambda_\tau : A_n \to B_n$$

by

$$\lambda_\tau(\sigma) = \tau\sigma,$$

that is, $\sigma \in A_n$ is mapped into $(1,2)\sigma$ by λ_τ. Observe that since σ is even, the permutation $(1,2)\sigma$ can be expressed as a product of a (1 + even number), or odd number, of transpositions, so $(1,2)\sigma$ is indeed in B_n. If for σ and μ in A_n it is true that $\lambda_\tau(\sigma) = \lambda_\tau(\mu)$, then

$$(1,2)\sigma = (1,2)\mu,$$

and since S_n is a group, we have $\sigma = \mu$. Thus λ_τ is a one-to-one function. Finally,

$$\tau = (1,2) = \tau^{-1},$$

so if $\rho \in B_n$, then

$$\tau^{-1}\rho \in A_n,$$

and

$$\lambda_\tau(\tau^{-1}\rho) = \tau(\tau^{-1}\rho) = \rho.$$

Thus λ_τ maps onto B_n. Hence the number of elements in A_n is the same as the number in B_n since there is a one-to-one correspondence between the elements of the sets.

Note that the product of two even permutations is again even. Also since $n \geq 2$, S_n has the transposition $(1, 2)$ and $\iota = (1, 2)(1, 2)$ is an even permutation. Finally, note that if σ is expressed as a product of transpositions, the product of the same transpositions taken in just the opposite order is σ^{-1}. Thus if σ is an even permutation, σ^{-1} must also be even. Referring to Theorem 5.12, we see that we have proved the following statement.

8.24 Theorem If $n \geq 2$, then the collection of all even permutations of $\{1, 2, 3, \cdots, n\}$ forms a subgroup of order $n!/2$ of the symmetric group S_n.

We can define a function called the **sign of a permutation**, $\text{sgn} : S_n \to \{1, -1\}$ by the formula

$$\text{sgn}(\sigma) = \begin{cases} 1 & \text{if } \sigma \text{ is even} \\ -1 & \text{if } \sigma \text{ is odd.} \end{cases}$$

Thinking of $\{1, -1\}$ as a group under multiplication, it is easy to see that sgn is a homomorphism. Since 1 is the identity in the group $\{1, -1\}$, $\text{Ker}(\text{sgn}) = \text{sgn}^{-1}[\{1\}]$ is a subgroup of S_n consisting of all the even permutations. The homomorphism sgn is used in the standard way of defining the determinant of a square matrix. Exercise 52 asks you to prove some of the standard facts about determinants using this definition.

8.25 Definition The subgroup of S_n consisting of the even permutations of n letters is the **alternating group A_n on n letters**. ■

Both S_n and A_n are very important groups. Cayley's theorem shows that every finite group G is structurally identical to some subgroup of S_n for $n = |G|$. It can be shown that there are no formulas involving just radicals for solution of polynomial equations of degree n for $n \geq 5$. This fact is actually due to the structure of A_n, surprising as that may seem!

EXERCISES 8

Computations

In Exercises 1 through 10 determine whether the given map is a group homomorphism. [Hint: To verify that a map is a homomorphism, you must check the homomorphism property. To check that a map is not a homomorphism you could either find a and b such that $\phi(ab) \neq \phi(a)\phi(b)$, or else you could determine that any of the properties in Theorem 8.5 fail.]

1. Let $\phi : \mathbb{Z}_{10} \to \mathbb{Z}_2$ be given by $\phi(x) =$ the remainder when x is divided by 2.
2. Let $\phi : \mathbb{Z}_9 \to \mathbb{Z}_2$ be given by $\phi(x) =$ the remainder when x is divided by 2.
3. Let $\phi : \mathbb{Q}^* \to \mathbb{Q}^*$ be given by $\phi(x) = |x|$.
4. Let $\phi : \mathbb{R} \to \mathbb{R}^+$ be given by $\phi(x) = 2^x$.
5. Let $\phi : D_4 \to \mathbb{Z}_4$ be given by $\phi(\rho^i) = \phi(\mu\rho^i) = i$ for $0 \leq i \leq 3$.

6. Let F be the additive group of all functions mapping $\mathbb{R}$ to $\mathbb{R}$. Let $\phi : F \to F$ be given by $\phi(f) = g$ where $g(x) = f(x) + x$.

7. Let F be as in Exercise 6 and $\phi : F \to F$ be defined by $\phi(f) = 5f$.

8. Let F be the additive group of all continuous functions mapping $\mathbb{R}$ to $\mathbb{R}$. Let $\phi : F \to \mathbb{R}$ be defined by $\phi(g) = \int_0^1 g(x)\,dx$.

9. Let M_n be the additive group of $n \times n$ matrices with real entries. Let $\phi : M_n \to \mathbb{R}$ be given by $\phi(A) = \det(A)$, the determinant of A.

10. Let M_n be as in Exercise 9 and $\phi : M_n \to \mathbb{R}$ be defined by $\phi(A) = \text{tr}(A)$ where $\text{tr}(A)$ is the trace of A, which is the sum of the entries on the diagonal.

In Exercises 11 through 16, compute the kernel for the given homomorphism ϕ.

11. $\phi : \mathbb{Z} \to \mathbb{Z}_8$ such that $\phi(1) = 6$.

12. $\phi : \mathbb{Z} \to \mathbb{Z}$ such that $\phi(1) = 12$.

13. $\phi : \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z}$ where $\phi(1, 0) = 3$ and $\phi(0, 1) = -5$.

14. $\phi : \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z}$ where $\phi(1, 0) = 6$ and $\phi(0, 1) = 9$.

15. $\phi : \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z} \times \mathbb{Z}$ where $\phi(1, 0) = (2, 5)$ and $\phi(0, 1) = (-3, 2)$.

16. Let D be the additive group of all differentiable functions mapping $\mathbb{R}$ to $\mathbb{R}$ and F the additive group of all functions from $\mathbb{R}$ to $\mathbb{R}$. $\phi : D \to F$ is given by $\phi(f) = f'$, the derivative of f.

In Exercises 17 through 22, find all orbits of the given permutation.

17. $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 1 & 3 & 6 & 2 & 4 \end{pmatrix}$

18. $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 5 & 6 & 2 & 4 & 8 & 3 & 1 & 7 \end{pmatrix}$

19. $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 3 & 5 & 1 & 4 & 6 & 8 & 7 \end{pmatrix}$

20. $\sigma : \mathbb{Z} \to \mathbb{Z}$ where $\sigma(n) = n + 1$

21. $\sigma : \mathbb{Z} \to \mathbb{Z}$ where $\sigma(n) = n + 2$

22. $\sigma : \mathbb{Z} \to \mathbb{Z}$ where $\sigma(n) = n - 3$

In Exercises 23 through 25, express the permutation of {1, 2, 3, 4, 5, 6, 7, 8} as a product of disjoint cycles, and then as a product of transpositions.

23. $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 8 & 2 & 6 & 3 & 7 & 4 & 5 & 1 \end{pmatrix}$

24. $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 3 & 6 & 4 & 1 & 8 & 2 & 5 & 7 \end{pmatrix}$

25. $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 5 & 3 & 2 & 8 & 4 & 7 & 6 & 1 \end{pmatrix}$

26. Figure 8.26 shows a Cayley digraph for the alternating group A_4 using the generating set $S = \{(1, 2, 3), (1, 2)(3, 4)\}$. Continue labeling the other nine vertices with the elements of A_4, expressed as a product of disjoint cycles.


Concepts

In Exercises 27 through 29, correct the definition of the italicized term without reference to the text, if correction is needed, so that it is in a form acceptable for publication.

27. For a permutation σ of a set A, an *orbit* of σ is a nonempty minimal subset of A that is mapped onto itself by σ.

28. The left regular representation of a group G is the map of G into S_G whose value at $g \in G$ is the permutation of G that carries each $x \in G$ into gx.

29. The *alternating group* is the group of all even permutations.



8.26 Figure

30. Before the proof of Cayley's Theorem, it is shown that λ_x is one-to-one. In the proof, one-to-one is shown again. Is it necessary to show one-to-one twice? Explain.

31. Determine whether each of the following is true or false.

a. Every permutation is a cycle.
b. Every cycle is a permutation.
c. The definition of even and odd permutations could have been given equally well before Theorem 8.19.
d. Every nontrivial subgroup H of S_9 containing some odd permutation contains a transposition.
e. A_5 has 120 elements.
f. S_n is not cyclic for any $n \geq 1$.
g. A_3 is a commutative group.
h. S_7 is isomorphic to the subgroup of all those elements of S_8 that leave the number 8 fixed.
i. S_7 is isomorphic to the subgroup of all those elements of S_8 that leave the number 5 fixed.
j. The odd permutations in S_8 form a subgroup of S_8.
k. Every group G is isomorphic with a subgroup of S_G.

32. The dihedral group is defined to be permutations with certain properties. Use the usual notation involving μ and ρ for elements in D_n.

a. Identify which elements in D_3 are even. Do the even elements form a cyclic group?
b. Identify which of elements of D_4 are even. Do the even elements form a cyclic group?
c. For which values of n do the even permutations of D_n form a cyclic group?

Proof Synopsis

33. Give a two-sentence synopsis of the proof of Cayley's Theorem.

34. Give a two-sentence synopsis of the proof of Theorem 8.19.

Theory

35. Suppose that $\phi : G \to G'$ is a group homomorphism and $a \in \text{Ker}\phi$. Show that for any $g \in G$, $gag^{-1} \in \text{Ker}\phi$.

36. Prove that a homomorphism $\phi : G \to G'$ is one-to-one if and only if $\text{Ker}(\phi)$ is the trivial subgroup of G.

37. Let $\phi : G \to G'$ be a group homomorphism. Show that $\phi(a) = \phi(b)$ if and only if $a^{-1}b \in \text{Ker}\phi$.

38. Use Exercise 37 to prove that if $\phi : G \to G'$ is a group homomorphism mapping onto G' and G is a finite group, then for any $b, c \in G'$, $|\phi^{-1}[\{b\}]| = |\phi^{-1}[\{c\}]|$. Conclude that if $|G|$ is a prime number, then either ϕ is an isomorphism or else G' is the trivial group.

39. Show that if $\phi : G \to G'$ and $\gamma : G' \to G''$ are group homomorphisms, then $\gamma \circ \phi : G \to G''$ is also a group homomorphism.

40. Let $\phi : G \to G'$ be a group homomorphism. Show that $\phi[G]$ is abelian if and only if $xyx^{-1}y^{-1} \in \text{Ker}(\phi)$ for all $x, y \in G$.

41. Prove the following about S_n if $n \geq 3$.

a. Every permutation in S_n can be written as a product of at most $n - 1$ transpositions.

b. Every permutation in S_n that is not a cycle can be written as a product of at most $n - 2$ transpositions.

c. Every odd permutation in S_n can be written as a product of $2n + 3$ transpositions, and every even permutation as a product of $2n + 8$ transpositions.

42. a. Draw a figure like Fig. 8.20 to illustrate that if i and j are in different orbits of σ and $\sigma(i) = i$, then the number of orbits of $(i, j)\sigma$ is one less than the number of orbits of σ.

b. Repeat part (a) if $\sigma(j) = j$ also.

43. Show that for every subgroup H of S_n for $n \geq 2$, either all the permutations in H are even or exactly half of them are even.

44. Let σ be a permutation of a set A. We shall say "σ **moves** $a \in A$" if $\sigma(a) \neq a$. If A is a finite set, how many elements are moved by a cycle $\sigma \in S_A$ of length n?

45. Let A be an infinite set. Let H be the set of all $\sigma \in S_A$ such that the number of elements moved by σ (see Exercise 44) is finite. Show that H is a subgroup of S_A.

46. Let A be an infinite set. Let K be the set of all $\sigma \in S_A$ that move (see Exercise 44) at most 50 elements of A. Is K a subgroup of S_A? Why?

47. Consider S_n for a fixed $n \geq 2$ and let σ be a fixed odd permutation. Show that every odd permutation in S_n is a product of σ and some permutation in A_n.

48. Show that if σ is a cycle of odd length, then σ^2 is a cycle.

49. Following the line of thought opened by Exercise 48, complete the following with a condition involving n and r so that the resulting statement is a theorem:

If σ is a cycle of length n, then σ^r is also a cycle if and only if . . .

50. Show that S_n is generated by $\{(1, 2), (1, 2, 3, \cdots, n)\}$. [*Hint:* Show that as r varies, $(1, 2, 3, \cdots, n)^r(1, 2)(1, 2, 3, \cdots, n)^{n-r}$ gives all the transpositions $(1, 2), (2, 3), (3, 4), \cdots, (n - 1, n), (n, 1)$. Then show that any transposition is a product of some of these transpositions and use Theorem 8.15.]

51. Let $\sigma \in S_n$ and define a relation on $\{1, 2, 3, \ldots, n\}$ by $i \sim j$ if and only if $j = \sigma^k(i)$ for some $k \in \mathbb{Z}$.

1. Prove that $\sim$ is an equivalence relation.

2. Prove that for any $1 \leq i \leq n$, the equivalence class of i is the orbit of i.

52. The usual definition for the determinant of an $n \times n$ matrix $A = (a_{i,j})$ is

$$\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1,\sigma(1)} a_{2,\sigma(2)} a_{3,\sigma(3)} \cdots a_{n,\sigma(n)}$$

where $\text{sgn}(\sigma)$ is the sign of σ. Using this definition, prove the following properties of determinants.

a. If a row of matrix A has all zero entries, then $\det(A) = 0$.

b. If two different rows of A are switched to obtain B, then $\det(B) = -\det(A)$.

c. If r times one row of A is added to another row of A to obtain a matrix B, then $\det(A) = \det(B)$

d. If a row of A is multiplied by r to obtain the matrix B, then $\det(B) = r \det(A)$.

53. Prove that any finite group G is isomorphic with a subgroup of $\text{GL}(n, \mathbb{R})$ for some n. [Hint: For each $\sigma \in S_n$, find a matrix in $\text{GL}(n, \mathbb{R})$ that sends each basis vector e_i to $e_{\sigma(i)}$. Use this to show that S_n is isomorphic with a subgroup of $\text{GL}(n, \mathbb{R})$.]

54. Prove Cayley's Theorem using the right regular representation rather than the left regular representation.

55. Let $\sigma \in S_n$. An inversion is a pair (i, j) such that $i < j$ and $\sigma(i) > \sigma(j)$. Prove Theorem 8.19 by showing that multiplying a permutation by a transposition changes the number of inversions by an odd number.