Table 26: Percentage of children 0-59 months of age reported ill during the last two weeks who received increased fluids and continued feeding, Country, Year

		Number of children under 5	Children with illness who drank:				Children with illness who ate:				Received	
	Reported illness in last two weeks		More	Same/Less	Missing/ DK	Total	Somewhat less/same/ more	Much less/none	Missing/ DK	Total	increased fluids and continued eating	Number of sick children
Male	45.2	970	19.9	78.3	1.8	100.0	75.8	22.2	2.0	100.0	14.9	438
Female	45.7	860	21.9	75.7	2.5	100.0	76.4	20.9	2.7	100.0	16.3	393
Central	40.7	663	25.6	73.3	1.2	100.0	81.4	17.4	1.2	100.0	19.4	270
South Central	47.2	370	17.2	81.1	1.7	100.0	74.3	24.0	1.7	100.0	12.5	174
West	37.4	270	16.5	78.8	4.7	100.0	78.9	16.4	4.7	100.0	11.7	101
East	47.4	219	28.7	70.4	.9	100.0	71.3	26.9	1.9	100.0	20.4	. 104
South	59.0	309	15.3	81.6	3.2	100.0	71.1	25.3	3.7	100.0	12.1	182
Urban	40.8	896	24.0	74.9	1.1	100.0	79.7	19.1	1.1	100.0	18.0	366
Rural	49.8	934	18.4	78.8	2.9	100.0	73.2	23.5	3.3	100.0	13.6	465
< 6 months	38.6	215	10.0	84.1	6.0	100.0	78.8	12.9	8.3	100.0	8.7	83
6-11 months	50.4	158	15.3	82.3	2.4	100.0	74.1	21.1	4.8	100.0	14.0	79
12-23 months	54.3	366	16.6	81.5	2.0	100.0	74.1	24.4	1.5	100.0	14.0	199
24-35 months	45.8	422	29.5	69.5	1.0	100.0	79.0	20.5	.5	100.0	22.2	
36-47 months	39.9	345	23.0	76.2	.8	100.0	72.6	26.6	.8	100.0	13.8	138
48-59 months	43.1	321	22.4	74.8	2.8	100.0	77.7	19.5	2.8	100.0	15.2	139
None	55.1	238	20.7	77.7	1.5	100.0	68.8	29.0	2.3	100.0	14.8	131
Mother's education												
Primary	45.2	496	18.2	80.1	1.7	100.0	75.6	21.8	2.6	100.0	15.1	224
Secondary +	43.4	1096	22.1	75.4	2.5	100.0	78.3	19.4	2.3	100.0	16.0	
Total	45.4	1830	20.8	3 77.0	2.1	100.0	76.1	21.6	2.3	100.0	15.5	831

Version date: 00.09.04 28