

# OM Review - XIX

Prof. C. V. Jawahar

IIIT Hyderabad

April 3, 2020

<https://forms.gle/o2f5oioWCPRirUgW8>

## Context for next question

We saw in one of the problems in the last lecture that K-Means is prone to local minima. Consider strategies S1, S2, S3 aimed to solve this issue. Given dataset  $S$  we need to find initial set of cluster centroids  $T$ .

S1: Pick  $k$  points from  $S$  that are farthest away from each other.

S2: Pick  $x \in S$  uniformly at random and set  $T \leftarrow \{x\}$

While  $|T| < k$ :

(i) Pick  $x \in S$  at random, with probabilities directly proportional to

$\min_{z \in T} \|x - z\|$

(ii)  $T \leftarrow T \cup \{x\}$

S3: Pick  $x \in S$  uniformly at random and set  $T \leftarrow \{x\}$

While  $|T| < k$ :

(i) Pick  $x \in S$  at random, with probabilities inversely proportional to

$\min_{z \in T} \|x - z\|$

(ii)  $T \leftarrow T \cup \{x\}$

## Question 1

Which of these are true with regards to above?

- (A) S2 is the best among S1,S2,S3
- (B) S3 is the best among S1,S2,S3
- (C) S1 is prone to outliers
- (D) S1 is not prone to outliers

More than one may be correct.

## Question 2

Which of these is true about following statements on K Means?

- (i) Bad initialization can lead to poor convergence speed
- (ii) Bad initialization can lead to bad overall clustering
- (iii) If we choose  $K = |S|$ , ie size of dataset; then  $J = 0$ , hence it is a good choice for K.

- (A) (i),(ii),(iii)
- (B) None of the others
- (C) (i)
- (D) (i),(ii)

## Question 3

Which of these is true?

- (i) MDS with Euclidean distance gives a linear combination of data as the answer
  - (ii) ISOMAP tries to preserve Euclidean distances between all pairs of points
- (A) Both (i) and (ii)
  - (B) Neither (i) nor (ii)
  - (C) (i) only
  - (D) (ii) only

## Question 4

Which of these is true about ISOMAP?

- (A) It can be obtained by solving for the eigenvectors of  $K = -\frac{1}{2}HDH$  where  $D$  is the Euclidean distance matrix and  $H$  is the centering matrix
- (B) It requires us to solve APSP problem once.
- (C) It requires us to solve APSP problem more than once.

APSP problem: All pairs shortest paths problem

More than one may be correct.

## Question 5

Which of these is true?

- (A) It is possible that LLE can lead to the same solution as PCA
- (B) None of the others
- (C) In Laplacian Eigenmaps, weights are optimized in one of the steps
- (D) In Laplacian Eigenmaps, weights are constants

More than one may be correct