

- Fourier Transforms.
- Discrete Fourier Transform.

- The Fourier transform is one of the most useful and widely used tool in physics – both traditional and computational.

- The Fourier transform is one of the most useful and widely used tool in physics – both traditional and computational.
- Allows one to break down functions/signals into their component parts and analyze, smooth or filter them.

- The Fourier transform is one of the most useful and widely used tool in physics – both traditional and computational.
- Allows one to break down functions/signals into their component parts and analyze, smooth or filter them.
- Also allows one to perform certain kinds of calculations and solve certain differential equations.

Any periodic function  $f(x)$  defined on a finite interval  $0 \leq x \leq L$  can be written as a Fourier series (assuming that the function is bounded and has at most a finite number of discontinuities).

Any periodic function  $f(x)$  defined on a finite interval  $0 \leq x \leq L$  can be written as a Fourier series (assuming that the function is bounded and has at most a finite number of discontinuities).

- If the function is even (i.e. symmetric) about the midpoint ( $x = \frac{L}{2}$ ) then one can write the cosine series:

$$f(x) = \sum_{k=0}^{\infty} \alpha_k \cos \left( \frac{2\pi kx}{L} \right)$$

Any periodic function  $f(x)$  defined on a finite interval  $0 \leq x \leq L$  can be written as a Fourier series (assuming that the function is bounded and has at most a finite number of discontinuities).

- If the function is even (i.e. symmetric) about the midpoint ( $x = \frac{L}{2}$ ) then one can write the cosine series:

$$f(x) = \sum_{k=0}^{\infty} \alpha_k \cos \left( \frac{2\pi kx}{L} \right)$$

- If the function is odd (i.e. antisymmetric) about the midpoint ( $x = \frac{L}{2}$ ) then one can write the sine series:

$$f(x) = \sum_{k=1}^{\infty} \beta_k \sin \left( \frac{2\pi kx}{L} \right)$$

# Fourier Series – periodic vs non periodic

- For a general function (with no special symmetry), one can write the series:

$$f(x) = \sum_{k=-\infty}^{\infty} \gamma_k \exp\left(i \frac{2\pi kx}{L}\right)$$



# Fourier Series – periodic vs non periodic

- For a general function (with no special symmetry), one can write the series:

$$f(x) = \sum_{k=-\infty}^{\infty} \gamma_k \exp\left(i \frac{2\pi kx}{L}\right)$$

- The Fourier series can only be used for periodic functions (ie the function in 0 to L is repeated in both directions).

# Fourier Series – periodic vs non periodic

- For a general function (with no special symmetry), one can write the series:

$$f(x) = \sum_{k=-\infty}^{\infty} \gamma_k \exp \left( i \frac{2\pi kx}{L} \right)$$

- The Fourier series can only be used for periodic functions (ie the function in 0 to L is repeated in both directions).
- If the function is not periodic, and we are only interested in a portion of this non periodic function over a finite interval, 0 to L, we can just take that portion and repeat it to create a periodic function!

- For a general function (with no special symmetry), one can write the series:

$$f(x) = \sum_{k=-\infty}^{\infty} \gamma_k \exp\left(i \frac{2\pi kx}{L}\right)$$

- The Fourier series can only be used for periodic functions (ie the function in 0 to L is repeated in both directions).
- If the function is not periodic, and we are only interested in a portion of this non periodic function over a finite interval, 0 to L, we can just take that portion and repeat it to create a periodic function!
- Then the Fourier coefficients will only give the correct information about the function in the interval 0 to L. Outside this interval, the function will be just repeated (and may not have anything to do with the original function).

The coefficients  $\gamma_k$  are in general complex:

The coefficients  $\gamma_k$  are in general complex:

$$\int_0^L f(x) \exp\left(-i\frac{2\pi kx}{L}\right) dx = \sum_{k'=-\infty}^{\infty} \gamma_{k'} \int_0^L \exp\left(i\frac{2\pi(k' - k)x}{L}\right) dx$$

The coefficients  $\gamma_k$  are in general complex:

$$\int_0^L f(x) \exp\left(-i\frac{2\pi kx}{L}\right) dx = \sum_{k'=-\infty}^{\infty} \gamma_{k'} \int_0^L \exp\left(i\frac{2\pi(k' - k)x}{L}\right) dx$$

If  $k' \neq k$ :

$$\begin{aligned} \int_0^L \exp\left(i\frac{2\pi(k' - k)x}{L}\right) dx &= \frac{L}{i2\pi(k' - k)} \left[ \exp\left(i\frac{2\pi(k' - k)x}{L}\right) \right]_0^L \\ &= \frac{L}{i2\pi(k' - k)} [e^{i2\pi(k' - k)} - 1] \\ &= 0 \end{aligned}$$

For  $k' = k$ :

$$\int_0^L f(x) \exp\left(-i\frac{2\pi kx}{L}\right) dx = L\gamma_k$$

For  $k' = k$ :

$$\int_0^L f(x) \exp \left( -i \frac{2\pi kx}{L} \right) dx = L\gamma_k$$

$$\gamma_k = \frac{1}{L} \int_0^L f(x) \exp \left( -i \frac{2\pi kx}{L} \right) dx$$



For  $k' = k$ :

$$\int_0^L f(x) \exp \left( -i \frac{2\pi kx}{L} \right) dx = L\gamma_k$$

$$\gamma_k = \frac{1}{L} \int_0^L f(x) \exp \left( -i \frac{2\pi kx}{L} \right) dx$$

Thus, given a function  $f(x)$ , we can find the Fourier coefficients  $\gamma_k$ , or given the coefficients, we can find the function  $f(x)$  – we can go back and forth freely between the function and the Fourier coefficients.

- For some functions  $f(x)$ , the Fourier coefficients can be determined analytically.

- For some functions  $f(x)$ , the Fourier coefficients can be determined analytically.
- There are, however, many cases where this is not doable – the integral is not doable because the function is too complicated or the function  $f(x)$  may not even be known in analytic form (for eg. if it is a signal measured in the laboratory experiment).

- For some functions  $f(x)$ , the Fourier coefficients can be determined analytically.
- There are, however, many cases where this is not doable – the integral is not doable because the function is too complicated or the function  $f(x)$  may not even be known in analytic form (for eg. if it is a signal measured in the laboratory experiment).
- In such cases, the integral can be evaluated numerically.

# Discrete Fourier Transform (DFT)

Applying the trapezoidal rule for integration (N slices of width  $h = L/N$ ) to calculate  $\gamma_k$ :

$$\gamma_k = \frac{1}{L} \frac{L}{N} \left[ \frac{f(0)}{2} + \frac{f(L)}{2} + \sum_{n=1}^{N-1} f(x_n) \exp \left( -i \frac{2\pi k x_n}{L} \right) \right]$$

where  $x_n = \frac{n}{N}L$

# Discrete Fourier Transform (DFT)

Applying the trapezoidal rule for integration (N slices of width  $h = L/N$ ) to calculate  $\gamma_k$ :

$$\gamma_k = \frac{1}{L} \frac{L}{N} \left[ \frac{f(0)}{2} + \frac{f(L)}{2} + \sum_{n=1}^{N-1} f(x_n) \exp \left( -i \frac{2\pi k x_n}{L} \right) \right]$$

where  $x_n = \frac{n}{N}L$

However, as  $f(x)$  is periodic,  $f(L) = f(0)$ :

$$\gamma_k = \frac{1}{N} \left[ \sum_{n=0}^{N-1} f(x_n) \exp \left( -i \frac{2\pi k x_n}{L} \right) \right]$$

# Discrete Fourier Transform (DFT)

Applying the trapezoidal rule for integration ( $N$  slices of width  $h = L/N$ ) to calculate  $\gamma_k$ :

$$\gamma_k = \frac{1}{L} \frac{L}{N} \left[ \frac{f(0)}{2} + \frac{f(L)}{2} + \sum_{n=1}^{N-1} f(x_n) \exp \left( -i \frac{2\pi k x_n}{L} \right) \right]$$

where  $x_n = \frac{n}{N} L$

However, as  $f(x)$  is periodic,  $f(L) = f(0)$ :

$$\gamma_k = \frac{1}{N} \left[ \sum_{n=0}^{N-1} f(x_n) \exp \left( -i \frac{2\pi k x_n}{L} \right) \right]$$

This formula can be used to evaluate the coefficients on a computer. A simpler way to write this is as:

$$\gamma_k = \frac{1}{N} \left[ \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi k n}{N} \right) \right]$$

where  $y_n = f(x_n)$

# Discrete Fourier Transform (DFT)

$$\gamma_k = \frac{1}{N} \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi kn}{N} \right)$$

This sum is known as the discrete Fourier transform (DFT).



# Discrete Fourier Transform (DFT)

$$\gamma_k = \frac{1}{N} \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi kn}{N} \right)$$

This sum is known as the discrete Fourier transform (DFT).  
Similarly one can define:

$$c_k = \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi kn}{N} \right)$$

# Discrete Fourier Transform (DFT)

$$\gamma_k = \frac{1}{N} \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi kn}{N} \right)$$

This sum is known as the discrete Fourier transform (DFT). Similarly one can define:

$$c_k = \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi kn}{N} \right)$$

The quantities  $\gamma_k$  and  $c_k$  only differ by the constant  $1/N$  factor. For our purpose they are both equal, and we define the latter as the definition of DFT.

# Discrete Fourier Transform (DFT)

So far, we have described the results of using Trapezoidal rule for the integral.. However, in principle, this is exact!:

# Discrete Fourier Transform (DFT)

So far, we have described the results of using Trapezoidal rule for the integral.. However, in principle, this is exact!:

$$\begin{aligned}\sum_{k=0}^{N-1} c_k \exp\left(i\frac{2\pi kn}{N}\right) &= \sum_{k=0}^{N-1} \sum_{n'=0}^{N-1} y_{n'} \exp\left(-i\frac{2\pi kn'}{N}\right) \exp\left(i\frac{2\pi kn}{N}\right) \\&= \sum_{n'=0}^{N-1} y_{n'} \sum_{k=0}^{N-1} \exp\left(i\frac{2\pi k(n' - n)}{N}\right) \\&= \sum_{n'=0}^{N-1} y_{n'} \frac{\exp(i2\pi(n' - n)) - 1}{\exp(i2\pi(n' - n)/N) - 1} = \sum_{n'=0}^{N-1} y_{n'} N\delta_{n,n'} \\&= Ny_n \quad \text{assuming } 0 \leq n \leq N\end{aligned}$$

# Discrete Fourier Transform (DFT)

So far, we have described the results of using Trapezoidal rule for the integral.. However, in principle, this is exact!:

$$\begin{aligned}\sum_{k=0}^{N-1} c_k \exp\left(i\frac{2\pi kn}{N}\right) &= \sum_{k=0}^{N-1} \sum_{n'=0}^{N-1} y_{n'} \exp\left(-i\frac{2\pi kn'}{N}\right) \exp\left(i\frac{2\pi kn}{N}\right) \\ &= \sum_{n'=0}^{N-1} y_{n'} \sum_{k=0}^{N-1} \exp\left(i\frac{2\pi k(n' - n)}{N}\right) \\ &= \sum_{n'=0}^{N-1} y_{n'} \frac{\exp(i2\pi(n' - n)) - 1}{\exp(i2\pi(n' - n)/N) - 1} = \sum_{n'=0}^{N-1} y_{n'} N\delta_{n,n'} \\ &= Ny_n \quad \text{assuming } 0 \leq n \leq N\end{aligned}$$

Equivalently,

$$y_n = \frac{1}{N} \sum_{k=0}^{N-1} c_k \exp\left(i\frac{2\pi kn}{N}\right)$$

This is called the "Inverse Discrete Fourier Transform."

# Discrete Fourier Transform (DFT)

- This inverse transform is the counterpart to the "forward" transform.

# Discrete Fourier Transform (DFT)

- This inverse transform is the counterpart to the "forward" transform.
- It tells us given the coefficients  $c_k$ , we can recover the values of the sample  $y_n$  they came from "exactly"!

# Discrete Fourier Transform (DFT)

- This inverse transform is the counterpart to the "forward" transform.
- It tells us given the coefficients  $c_k$ , we can recover the values of the sample  $y_n$  they came from "exactly"!
- Thus both  $c_k$  and  $y_n$  give a complete representation of the data and we can move from one to the other without losing any detail.



# Discrete Fourier Transform (DFT)

- This inverse transform is the counterpart to the "forward" transform.
- It tells us given the coefficients  $c_k$ , we can recover the values of the sample  $y_n$  they came from "exactly"!
- Thus both  $c_k$  and  $y_n$  give a complete representation of the data and we can move from one to the other without losing any detail.
- It is important to appreciate that unlike the original Fourier series, the discrete version only gives us the sample values at  $y_n = f(x_n)$ . It tells us nothing about the value of the function  $f(x)$  in between the points.

# Discrete Fourier Transform (DFT)

- This inverse transform is the counterpart to the "forward" transform.
- It tells us given the coefficients  $c_k$ , we can recover the values of the sample  $y_n$  they came from "exactly"!
- Thus both  $c_k$  and  $y_n$  give a complete representation of the data and we can move from one to the other without losing any detail.
- It is important to appreciate that unlike the original Fourier series, the discrete version only gives us the sample values at  $y_n = f(x_n)$ . It tells us nothing about the value of the function  $f(x)$  in between the points.
- So, two different functions with same values at the sample points will have the same DFT – no matter what they do in between the points!

Suppose all the  $y_n$  are real and consider the value of  $c_k$  for some  $k$  that is less than  $N$  but greater than  $\frac{N}{2}$ .

Suppose all the  $y_n$  are real and consider the value of  $c_k$  for some  $k$  that is less than  $N$  but greater than  $\frac{N}{2}$ .

We can write  $k = N - r$  with  $1 \leq r \leq \frac{N}{2}$

Suppose all the  $y_n$  are real and consider the value of  $c_k$  for some  $k$  that is less than  $N$  but greater than  $\frac{N}{2}$ .

We can write  $k = N - r$  with  $1 \leq r \leq \frac{N}{2}$

$$\begin{aligned} c_{N-r} &= \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi(N-r)n}{N} \right) \\ &= \sum_{n=0}^{N-1} y_n \exp(-i2\pi n) \exp \left( i \frac{2\pi r n}{N} \right) \\ &= \sum_{n=0}^{N-1} y_n \exp \left( i \frac{2\pi r n}{N} \right) = c_r^* \end{aligned}$$

# DFT for real functions

Suppose all the  $y_n$  are real and consider the value of  $c_k$  for some  $k$  that is less than  $N$  but greater than  $\frac{N}{2}$ .

We can write  $k = N - r$  with  $1 \leq r \leq \frac{N}{2}$

$$\begin{aligned} c_{N-r} &= \sum_{n=0}^{N-1} y_n \exp \left( -i \frac{2\pi(N-r)n}{N} \right) \\ &= \sum_{n=0}^{N-1} y_n \exp(-i2\pi n) \exp \left( i \frac{2\pi r n}{N} \right) \\ &= \sum_{n=0}^{N-1} y_n \exp \left( i \frac{2\pi r n}{N} \right) = c_r^* \end{aligned}$$

Thus:  $c_{N-1} = c_1^*$   $c_{N-2} = c_2^*$  and so forth. That means that Fourier coefficients  $c_k$  of a real function only has to be calculated for  $0 \leq k \leq \frac{N}{2}$ .

```
from numpy import zeros
from cmath import exp, pi

def dft(y):
    N = len(y)
    c = zeros(N//2 + 1, complex)
    for k in range(N//2+1):
        for n in range(N):
            c[k] += y[n]*exp(-2j*pi*k*n/N)
    return c
```

One thing to notice about the DFT is that we can shift the sample points along the  $x$ -axis, and not much changes.

Suppose:

$$x'_n = x_n + \Delta = \frac{n}{N}L + \Delta$$



One thing to notice about the DFT is that we can shift the sample points along the  $x$ -axis, and not much changes.

Suppose:

$$x'_n = x_n + \Delta = \frac{n}{N}L + \Delta$$

Then the DFT is:

$$\begin{aligned} c_k &= \sum_{n=0}^{N-1} f(x_n + \Delta) \exp \left( -i \frac{2\pi k(x_n + \Delta)}{L} \right) \\ &= \exp \left( -i \frac{2\pi k\Delta}{L} \right) \sum_{n=0}^{N-1} f(x'_n) \exp \left( -i \frac{2\pi kx_n}{L} \right) \\ &= \exp \left( -i \frac{2\pi k\Delta}{L} \right) \sum_{n=0}^{N-1} y'_n \exp \left( -i \frac{2\pi kn}{N} \right) \end{aligned}$$

$$c_k = \exp\left(-i\frac{2\pi k\Delta}{L}\right) \sum_{n=0}^{N-1} y'_n \exp\left(-i\frac{2\pi kn}{N}\right)$$

$$c_k = \exp\left(-i\frac{2\pi k\Delta}{L}\right) \sum_{n=0}^{N-1} y'_n \exp\left(-i\frac{2\pi kn}{N}\right)$$

But this is the same as the original DFT except for a (k-dependent) phase factor. Thus the DFT is really independent of where we choose to place the samples – only the coefficients change by a phase factor.

## Two-dimensional Fourier Transforms

Functions of two variables  $f(x, y)$  can also be Fourier transformed, using a two dimensional Fourier transform. This simply means that one first transform with respect to one variable and then the other!

# Two-dimensional Fourier Transforms

Functions of two variables  $f(x, y)$  can also be Fourier transformed, using a two dimensional Fourier transform. This simply means that one first transform with respect to one variable and then the other!

Suppose that we have an  $M \times N$  grid points of samples  $y_{mn}$ . We first transform on each of the  $M$  rows:

$$c'_{ml} = \sum_{n=0}^{N-1} y_{mn} \exp \left( -i \frac{2\pi l n}{N} \right)$$

# Two-dimensional Fourier Transforms

Functions of two variables  $f(x, y)$  can also be Fourier transformed, using a two dimensional Fourier transform. This simply means that one first transform with respect to one variable and then the other!

Suppose that we have an  $M \times N$  grid points of samples  $y_{mn}$ . We first transform on each of the  $M$  rows:

$$c'_{ml} = \sum_{n=0}^{N-1} y_{mn} \exp \left( -i \frac{2\pi l n}{N} \right)$$

Then we take the  $l^{th}$  coefficient of each row and Fourier transform them :

$$c_{kl} = \sum_{m=0}^{M-1} c'_{ml} \exp \left( -i \frac{2\pi k m}{M} \right)$$

Alternatively, we can write a single expression for the complete Fourier transform in two dimensions:

$$c_{kl} = \sum_{m=0}^{M-1} \sum_{n=0}^{N-1} y_{mn} \exp \left[ -i2\pi \left( \frac{ln}{N} + \frac{km}{M} \right) \right]$$

Alternatively, we can write a single expression for the complete Fourier transform in two dimensions:

$$c_{kl} = \sum_{m=0}^{M-1} \sum_{n=0}^{N-1} y_{mn} \exp \left[ -i2\pi \left( \frac{ln}{N} + \frac{km}{M} \right) \right]$$

The corresponding inverse transform is:

$$y_{mn} = \frac{1}{MN} \sum_{k=0}^{M-1} \sum_{l=0}^{N-1} c_{kl} \exp \left[ i2\pi \left( \frac{ln}{N} + \frac{km}{M} \right) \right]$$