

- Derivatives.
- Errors.
- Higher order derivatives.

In general, derivatives command much less attention than integrals. This is because

- The basic techniques for numerical derivatives are quite simple.

In general, derivatives command much less attention than integrals. This is because

- The basic techniques for numerical derivatives are quite simple.
- Derivatives of known functions can always be calculated analytically. So there is less of a need to calculate it numerically.

In general, derivatives command much less attention than integrals. This is because

- The basic techniques for numerical derivatives are quite simple.
- Derivatives of known functions can always be calculated analytically. So there is less of a need to calculate it numerically.
- There are significant practical problems with numerical derivatives. As a result they are used less often than numerical integration.

Standard definition of derivatives that you see in calculus is:

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

# Forward and Backward differences

Standard definition of derivatives that you see in calculus is:

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

The basic method of calculating derivatives is just implemetation of the formula (ofcourse the limit cannot be taken in practice):

$$\frac{df}{dx} \simeq \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

This is called the forward difference.

# Forward and Backward differences

Standard definition of derivatives that you see in calculus is:

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

The basic method of calculating derivatives is just implemetation of the formula (ofcourse the limit cannot be taken in practice):

$$\frac{df}{dx} \simeq \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

This is called the forward difference.

There is also backward difference:

$$\frac{df}{dx} \simeq \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h}$$

There are two sources of errors in calculating derivatives:

- Round off errors.
- Approximation error (as we cannot take  $h \rightarrow 0$ ).



There are two sources of errors in calculating derivatives:

- Round off errors.
- Approximation error (as we cannot take  $h \rightarrow 0$ ).

Consider the forward difference. We can write the Taylor expansion of  $f(x + h)$  as

$$f(x + h) = f(x) + hf'(x) + \frac{1}{2}h^2f''(x) + \dots$$
$$f'(x) = \frac{f(x + h) - f(x)}{h} - \frac{1}{2}h^2f''(x) \dots$$

The approximation error is  $\frac{1}{2}h^2f''(x)$ .

There are two sources of errors in calculating derivatives:

- Round off errors.
- Approximation error (as we cannot take  $h \rightarrow 0$ ).

Consider the forward difference. We can write the Taylor expansion of  $f(x + h)$  as

$$f(x + h) = f(x) + hf'(x) + \frac{1}{2}h^2f''(x) + \dots$$
$$f'(x) = \frac{f(x + h) - f(x)}{h} - \frac{1}{2}h^2f''(x) \dots$$

The approximation error is  $\frac{1}{2}h^2f''(x)$ .

But because we are subtracting numbers very close to each other (if  $h$  is small), then the round off error also comes into play!

- On a computer, we can typically calculate  $f(x)$  to an accuracy  $Cf(x)$  where  $C \sim 10^{-16}$ .

- On a computer, we can typically calculate  $f(x)$  to an accuracy  $Cf(x)$  where  $C \sim 10^{-16}$ .
- Since  $f(x+h)$  is close to  $f(x)$ , the error in  $f(x+h) - f(x)$  is  $\sim 2C|f(x)|$ .

- On a computer, we can typically calculate  $f(x)$  to an accuracy  $Cf(x)$  where  $C \sim 10^{-16}$ .
- Since  $f(x+h)$  is close to  $f(x)$ , the error in  $f(x+h) - f(x)$  is  $\sim 2C|f(x)|$ .
- The total error  $\epsilon$  is:

$$\epsilon = \frac{2C|f(x)|}{h} + \frac{1}{2}hf''(x)$$

- On a computer, we can typically calculate  $f(x)$  to an accuracy  $Cf(x)$  where  $C \sim 10^{-16}$ .
- Since  $f(x+h)$  is close to  $f(x)$ , the error in  $f(x+h) - f(x)$  is  $\sim 2C|f(x)|$ .
- The total error  $\epsilon$  is:

$$\epsilon = \frac{2C|f(x)|}{h} + \frac{1}{2}hf''(x)$$

- Minimizing this error with respect to  $h$ :

$$0 = -\frac{2C|f(x)|}{h^2} + \frac{1}{2}f''(x)$$

$$h = \sqrt{\frac{4C|f(x)|}{|f''(x)|}}$$

- On a computer, we can typically calculate  $f(x)$  to an accuracy  $Cf(x)$  where  $C \sim 10^{-16}$ .
- Since  $f(x+h)$  is close to  $f(x)$ , the error in  $f(x+h) - f(x)$  is  $\sim 2C|f(x)|$ .
- The total error  $\epsilon$  is:

$$\epsilon = \frac{2C|f(x)|}{h} + \frac{1}{2}hf''(x)$$

- Minimizing this error with respect to  $h$ :

$$0 = -\frac{2C|f(x)|}{h^2} + \frac{1}{2}f''(x)$$
$$h = \sqrt{\frac{4C|f(x)|}{|f''(x)|}}$$

- If  $|f(x)|$  and  $|f''(x)|$  are order 1,  $h \sim \sqrt{C} \sim 10^{-8}$ .

$$\frac{df}{dx} \simeq \frac{f(x + h/2) - f(x - h/2)}{h}$$

The error analysis gives:

$$h = \left( 24C \left| \frac{f(x)}{f'''(x)} \right| \right)^{1/3}$$

If  $f(x)$  and  $f'''(x)$  are of order 1, then  $h \sim C^{1/3} \sim 10^{-5}$ .

Similar ideas can be extended to higher order derivatives.