

- Monte Carlo integration.
- Non-uniform distributions.
- Random walk.

- There are better ways of evaluating this integral:

$$I = \int_a^b f(x) dx$$

- There are better ways of evaluating this integral:

$$I = \int_a^b f(x)dx$$

- Average value  $\langle f \rangle$  in range  $a$  to  $b$  is:

$$\langle f \rangle = \frac{1}{b-a} \int_a^b f(x)dx = \frac{I}{b-a}$$

- There are better ways of evaluating this integral:

$$I = \int_a^b f(x) dx$$

- Average value  $\langle f \rangle$  in range  $a$  to  $b$  is:

$$\langle f \rangle = \frac{1}{b-a} \int_a^b f(x) dx = \frac{I}{b-a}$$

- A simple way to estimate  $\langle f \rangle$  is to just measure  $f(x)$  at  $N$  points,  $x_1, x_2, \dots, x_N$  chosen uniformly between  $a$  and  $b$ :

$$\langle f \rangle = \frac{1}{N} \sum_{i=1}^N f(x_i)$$

$$I \simeq \frac{b-a}{N} \sum_{i=1}^N f(x_i)$$

- Variance of the sum on  $N$  independent random numbers is equal to  $N$  times the variance of a single one.

- Variance of the sum on  $N$  independent random numbers is equal to  $N$  times the variance of a single one.
- Random numbers in this case are the values  $f(x_i)$  and we can estimate the variance of a single one of them  $\text{var} f = \langle f^2 \rangle - \langle f \rangle^2$  with:

$$\langle f \rangle = \frac{1}{N} \sum_{i=1}^N f(x_i) \quad \langle f^2 \rangle = \frac{1}{N} \sum_{i=1}^N [f(x_i)]^2$$

- Variance of the sum on  $N$  independent random numbers is equal to  $N$  times the variance of a single one.
- Random numbers in this case are the values  $f(x_i)$  and we can estimate the variance of a single one of them  $\text{var} f = \langle f^2 \rangle - \langle f \rangle^2$  with:

$$\langle f \rangle = \frac{1}{N} \sum_{i=1}^N f(x_i) \quad \langle f^2 \rangle = \frac{1}{N} \sum_{i=1}^N [f(x_i)]^2$$

- Variance on the sum is  $N$  times the variance on a single term or  $N \text{var} f$ .

- Variance of the sum on  $N$  independent random numbers is equal to  $N$  times the variance of a single one.
- Random numbers in this case are the values  $f(x_i)$  and we can estimate the variance of a single one of them  $\text{var} f = \langle f^2 \rangle - \langle f \rangle^2$  with:

$$\langle f \rangle = \frac{1}{N} \sum_{i=1}^N f(x_i) \quad \langle f^2 \rangle = \frac{1}{N} \sum_{i=1}^N [f(x_i)]^2$$

- Variance on the sum is  $N$  times the variance on a single term or  $N \text{var} f$ .
- Error/standard deviation on the integral:

$$\sigma = \frac{b-a}{N} \sqrt{N \text{var} f} = (b-a) \frac{\sqrt{\text{var} f}}{\sqrt{N}}$$

which goes as  $1/\sqrt{N}$  but the variance is smaller!



In a large number of situations, random numbers with nonuniform distribution are needed (rather than uniform distribution)

In a large number of situations, random numbers with nonuniform distribution are needed (rather than uniform distribution)

- Radioactive decay

In a large number of situations, random numbers with nonuniform distribution are needed (rather than uniform distribution)

- Radioactive decay
- Experiments with different types of distributions

How does one generate non-uniform random number distributions with a uniform random number generators?

Generating a non-uniform distribution with a probability distribution  $w(x)$ :

Generating a non-uniform distribution with a probability distribution  $w(x)$ :

- Generate two random numbers  $x_i$  on  $[x_{min}, x_{max}]$  and  $y_i$  on  $[y_{min}, y_{max}]$

Generating a non-uniform distribution with a probability distribution  $w(x)$ :

- Generate two random numbers  $x_i$  on  $[x_{min}, x_{max}]$  and  $y_i$  on  $[y_{min}, y_{max}]$
- If  $y_i < w(x_i)$  accept  $x_i$ .

Generating a non-uniform distribution with a probability distribution  $w(x)$ :


- Generate two random numbers  $x_i$  on  $[x_{min}, x_{max}]$  and  $y_i$  on  $[y_{min}, y_{max}]$
- If  $y_i < w(x_i)$  accept  $x_i$ .
- If  $y_i > w(x_i)$  reject  $x_i$ .

Generating a non-uniform distribution with a probability distribution  $w(x)$ :

- Generate two random numbers  $x_i$  on  $[x_{min}, x_{max}]$  and  $y_i$  on  $[y_{min}, y_{max}]$
- If  $y_i < w(x_i)$  accept  $x_i$ .
- If  $y_i > w(x_i)$  reject  $x_i$ .
- The  $x_i$  so accepted will have the weighting  $w(x_i)$ .



$$w(x) = e^{-x^2} \quad x \in [0, 2]$$



- Suppose we have a source of random numbers  $z$  drawn from a distribution with probability density  $q(z)$  – the probability of generating a number between  $z$  and  $z + dz$  is  $q(z)dz$ .

- Suppose we have a source of random numbers  $z$  drawn from a distribution with probability density  $q(z)$  – the probability of generating a number between  $z$  and  $z + dz$  is  $q(z)dz$ .
- Suppose we have a function  $x = x(z)$ .

- Suppose we have a source of random numbers  $z$  drawn from a distribution with probability density  $q(z)$  – the probability of generating a number between  $z$  and  $z + dz$  is  $q(z)dz$ .
- Suppose we have a function  $x = x(z)$ .
- If  $z$  is random,  $x(z)$  will also be random, with a different distribution  $p(x)$ .

- Suppose we have a source of random numbers  $z$  drawn from a distribution with probability density  $q(z)$  – the probability of generating a number between  $z$  and  $z + dz$  is  $q(z)dz$ .
- Suppose we have a function  $x = x(z)$ .
- If  $z$  is random,  $x(z)$  will also be random, with a different distribution  $p(x)$ .
- Our goal is to choose the function  $x(z)$  such that  $x$  has the distribution we want.

- The probability of generating a value of  $x$  between  $x$  and  $x + dx$  is by definition equal to the probability of generating a value of  $z$  in the corresponding  $z$  interval:

$$p(x)dx = q(z)dz$$

- The probability of generating a value of  $x$  between  $x$  and  $x + dx$  is by definition equal to the probability of generating a value of  $z$  in the corresponding  $z$  interval:

$$p(x)dx = q(z)dz$$

- In most common cases,  $q(z) = 1$  in the interval  $[0, 1]$ .  
Then integrating both sides:

$$\int_{-\infty}^{x(z)} p(x')dx' = \int_0^z dz' = z$$

- The probability of generating a value of  $x$  between  $x$  and  $x + dx$  is by definition equal to the probability of generating a value of  $z$  in the corresponding  $z$  interval:

$$p(x)dx = q(z)dz$$

- In most common cases,  $q(z) = 1$  in the interval  $[0, 1]$ . Then integrating both sides:

$$\int_{-\infty}^{x(z)} p(x')dx' = \int_0^z dz' = z$$

- If we can do the integral on the left and solve the equation, we will have the required  $x(z)$ !



Example: Suppose we want to generate real random numbers  $x$  in the interval  $[0, \infty]$  with the exponential probability distribution:

$$p(x) = \frac{1}{\mu} e^{-\mu x}$$

Example: Suppose we want to generate real random numbers  $x$  in the interval  $[0, \infty]$  with the exponential probability distribution:

$$p(x) = \frac{1}{\mu} e^{-\mu x}$$

$$\mu \int_0^{x(z)} e^{-\mu x'} dx' = 1 - e^{-\mu x} = z$$

$$x = -\frac{1}{\mu} \ln(1 - z)$$

Example: Suppose we want to generate real random numbers  $x$  in the interval  $[0, \infty]$  with the exponential probability distribution:

$$p(x) = \frac{1}{\mu} e^{-\mu x}$$

$$\mu \int_0^{x(z)} e^{-\mu x'} dx' = 1 - e^{-\mu x} = z$$

$$x = -\frac{1}{\mu} \ln(1 - z)$$

Thus if we feed the above equation uniformly distributed  $z$  in interval  $[0, 1]$ , it will generate the exponential distribution  $x$  for us.

- A common problem in physics calculations is the generation of random numbers drawn from a Gaussian (or normal) distribution:

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

- A common problem in physics calculations is the generation of random numbers drawn from a Gaussian (or normal) distribution:

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

- This does not allow inversion! As the integral is not analytic...

- A common problem in physics calculations is the generation of random numbers drawn from a Gaussian (or normal) distribution:

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

- This does not allow inversion! As the integral is not analytic...
- However, consider two independent random numbers  $x$  and  $y$  drawn from a Gaussian distribution with the same  $\sigma$ . The probability that a point  $(x, y)$  falls in an element  $dxdy$  of the  $xy$  plane:

$$p(x)dx \times p(y)dy = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{x^2 + y^2}{2\sigma^2}\right) dxdy$$

- In polar coordinates:

$$p(r, \theta) dr d\theta = \frac{r}{\sigma^2} \exp\left(-\frac{r^2}{2\sigma^2}\right) dr \times \frac{d\theta}{2\pi} \equiv p(r) dr \times p(\theta) d\theta$$

- In polar coordinates:

$$p(r, \theta) dr d\theta = \frac{r}{\sigma^2} \exp\left(-\frac{r^2}{2\sigma^2}\right) dr \times \frac{d\theta}{2\pi} \equiv p(r) dr \times p(\theta) d\theta$$

- In this expression,  $\theta$  is just a uniform distribution

$$p(\theta) = \frac{1}{2\pi}$$



- In polar coordinates:

$$p(r, \theta) dr d\theta = \frac{r}{\sigma^2} \exp\left(-\frac{r^2}{2\sigma^2}\right) dr \times \frac{d\theta}{2\pi} \equiv p(r) dr \times p(\theta) d\theta$$

- In this expression,  $\theta$  is just a uniform distribution

$$p(\theta) = \frac{1}{2\pi}$$

- The distribution of  $r$  can be inverted with  $z$  uniformly distributed:

$$r = \sqrt{-2\sigma^2 \ln(1 - z)}$$

- In polar coordinates:

$$p(r, \theta) dr d\theta = \frac{r}{\sigma^2} \exp\left(-\frac{r^2}{2\sigma^2}\right) dr \times \frac{d\theta}{2\pi} \equiv p(r) dr \times p(\theta) d\theta$$

- In this expression,  $\theta$  is just a uniform distribution

$$p(\theta) = \frac{1}{2\pi}$$

- The distribution of  $r$  can be inverted with  $z$  uniformly distributed:

$$r = \sqrt{-2\sigma^2 \ln(1 - z)}$$

- Then one can construct  $x$  and  $y$  as:

$$x = r \cos(\theta) \quad y = r \sin(\theta)$$

- A simple random walk is a sequence of unit steps where each step is taken in the direction of one of the coordinate axis, and each possible direction has equal probability of being chosen.

- A simple random walk is a sequence of unit steps where each step is taken in the direction of one of the coordinate axis, and each possible direction has equal probability of being chosen.
- Random walk on a lattice:

- A simple random walk is a sequence of unit steps where each step is taken in the direction of one of the coordinate axis, and each possible direction has equal probability of being chosen.
- Random walk on a lattice:
  - In two dimensions, a single step starting at the point with integer coordinates  $(x,y)$  would be equally likely to move to any of one of the four neighbors  $(x+1,y)$ ,  $(x-1,y)$ ,  $(x,y+1)$  and  $(x,y-1)$ .

- A simple random walk is a sequence of unit steps where each step is taken in the direction of one of the coordinate axis, and each possible direction has equal probability of being chosen.
- Random walk on a lattice:
  - In two dimensions, a single step starting at the point with integer coordinates  $(x,y)$  would be equally likely to move to any of one of the four neighbors  $(x+1,y)$ ,  $(x-1,y)$ ,  $(x,y+1)$  and  $(x,y-1)$ .
  - In one dimension walk there are two possible neighbors

- A simple random walk is a sequence of unit steps where each step is taken in the direction of one of the coordinate axis, and each possible direction has equal probability of being chosen.
- Random walk on a lattice:
  - In two dimensions, a single step starting at the point with integer coordinates  $(x,y)$  would be equally likely to move to any of one of the four neighbors  $(x+1,y)$ ,  $(x-1,y)$ ,  $(x,y+1)$  and  $(x,y-1)$ .
  - In one dimension walk there are two possible neighbors
  - In three dimensions there are six possible neighbors.

- Brownian motion (answer the question - how many collisions, on average, a particle must take to travel a distance  $R$ ).




- Brownian motion (answer the question - how many collisions, on average, a particle must take to travel a distance  $R$ ).
- Electron transport in metals

- Brownian motion (answer the question - how many collisions, on average, a particle must take to travel a distance  $R$ ).
- Electron transport in metals
- Polymer simulations.

- Brownian motion (answer the question - how many collisions, on average, a particle must take to travel a distance  $R$ ).
- Electron transport in metals
- Polymer simulations.
- ...

# Simple Random walk

In 100 steps,  $\langle r \rangle \sim 8.9$



- Persistent random walk

- Persistent random walk
- Restricted random walk

- Persistent random walk
- Restricted random walk
- Self-avoiding random walk

- Persistent random walk
- Restricted random walk
- Self-avoiding random walk
- ...



- Persistent random walk
- Restricted random walk
- Self-avoiding random walk
- ...

Examples of applications:

Spread of infectious diseases and effects of immunization

Spreading of fire

- A persistent random walk in 2 dimensions in a city with  $n \times n$  blocks

# A persistent random walk


- A persistent random walk in 2 dimensions in a city with  $n \times n$  blocks
- Condition: the walker can not step back

# A persistent random walk

- A persistent random walk in 2 dimensions in a city with  $n \times n$  blocks
- Condition: the walker can not step back
- Goal: find average number of steps to get out the city

# Persistent Random walk

To escape  $24 \times 24$ ,  $\langle n \rangle \sim 92$



- Importance Sampling and Statistical mechanics.
- Markov Chain Monte Carlo.
- Simulated Annealing

- One of the problems in statistical mechanics is to calculate the average (or expectation) value of a quantity of interest in a physical system in thermal equilibrium at temperature  $T$ .

# Importance Sampling and Statistical Mechanics

- One of the problems in statistical mechanics is to calculate the average (or expectation) value of a quantity of interest in a physical system in thermal equilibrium at temperature  $T$ .
- One in general doesn't know the exact state of the physical system.



- One of the problems in statistical mechanics is to calculate the average (or expectation) value of a quantity of interest in a physical system in thermal equilibrium at temperature  $T$ .
- One in general doesn't know the exact state of the physical system.
- Instead we know that at temperature  $T$ , a system will pass through a succession of states such that the probability of occupying a state  $i$  with energy  $E_i$  is:

$$P(E_i) = \frac{e^{-\beta E_i}}{Z}, \quad Z = \sum_i e^{-\beta E_i}$$

- One of the problems in statistical mechanics is to calculate the average (or expectation) value of a quantity of interest in a physical system in thermal equilibrium at temperature  $T$ .
- One in general doesn't know the exact state of the physical system.
- Instead we know that at temperature  $T$ , a system will pass through a succession of states such that the probability of occupying a state  $i$  with energy  $E_i$  is:

$$P(E_i) = \frac{e^{-\beta E_i}}{Z}, \quad Z = \sum_i e^{-\beta E_i}$$

- The average value of  $X$ :

$$\langle X \rangle = \sum_i X_i P(E_i)$$

- In rare few cases, this can be done analytically..

- In rare few cases, this can be done analytically..
- In most cases, this has to be done numerically – however a brute force summing will be out of question (in most cases) – example if one has  $10^{23}$  gas molecules, and even if each had 2 quantum states then the total number of states is  $2^{10^{23}}$ !!!!

- In rare few cases, this can be done analytically..
- In most cases, this has to be done numerically – however a brute force summing will be out of question (in most cases) – example if one has  $10^{23}$  gas molecules, and even if each had 2 quantum states then the total number of states is  $2^{10^{23}}$ !!!!
- We can take the approach that we will calculate the sum via Monte Carlo (random sampling). In that case, we will chose  $N$  random states,  $k = 1 \dots N$  and calculate:

$$\langle X \rangle \simeq \frac{\sum_{k=1}^N X_k P(E_k)}{\sum_{k=1}^N P(E_k)}$$

The denominator is needed to normalize the weighted average correctly..

- Unfortunately, even this will not work correctly!

- Unfortunately, even this will not work correctly!
- In most cases, with  $E_i \gg k_B T$ , the Boltzmann probability will be exponentially small...

- Unfortunately, even this will not work correctly!
- In most cases, with  $E_i \gg k_B T$ , the Boltzmann probability will be exponentially small...
- That is most of the states we choose will have have very little contribution to the original average.



- Unfortunately, even this will not work correctly!
- In most cases, with  $E_i \gg k_B T$ , the Boltzmann probability will be exponentially small...
- That is most of the states we choose will have have very little contribution to the original average.
- Typically, only a small fraction of states will actually contribute significantly to the value of the sum – and choosing states at random, we are unlikely to pick these important states and hence get a poor estimate of the true value.

- Unfortunately, even this will not work correctly!
- In most cases, with  $E_i \gg k_B T$ , the Boltzmann probability will be exponentially small...
- That is most of the states we choose will have have very little contribution to the original average.
- Typically, only a small fraction of states will actually contribute significantly to the value of the sum – and choosing states at random, we are unlikely to pick these important states and hence get a poor estimate of the true value.
- But this is ideally suited for importance sampling!

- Importance sampling calculates the correct value of a sum using a set of samples drawn non-uniformly.

- Importance sampling calculates the correct value of a sum using a set of samples drawn non-uniformly.
- For any quantity  $g_i$  that depends on states  $i$  we can define a weighted average over states:

$$\langle g \rangle_w = \frac{\sum_i w_i g_i}{\sum_i w_i}$$

where  $w_i$  is any sets of weights we choose.

- Importance sampling calculates the correct value of a sum using a set of samples drawn non-uniformly.
- For any quantity  $g_i$  that depends on states  $i$  we can define a weighted average over states:

$$\langle g \rangle_w = \frac{\sum_i w_i g_i}{\sum_i w_i}$$

where  $w_i$  is any sets of weights we choose.

- Making the particular choice  $g_i = X_i P(E_i) / w_i$ :

$$\left\langle \frac{X_i P(E_i)}{w_i} \right\rangle_w = \frac{\sum_i X_i P(E_i)}{\sum_i w_i} = \frac{\langle X \rangle}{\sum_i w_i}$$

- Importance sampling calculates the correct value of a sum using a set of samples drawn non-uniformly.
- For any quantity  $g_i$  that depends on states  $i$  we can define a weighted average over states:

$$\langle g \rangle_w = \frac{\sum_i w_i g_i}{\sum_i w_i}$$

where  $w_i$  is any sets of weights we choose.

- Making the particular choice  $g_i = X_i P(E_i) / w_i$ :

$$\left\langle \frac{X_i P(E_i)}{w_i} \right\rangle_w = \frac{\sum_i X_i P(E_i)}{\sum_i w_i} = \frac{\langle X \rangle}{\sum_i w_i}$$

- Then:

$$\langle X \rangle = \left\langle \frac{X_i P(E_i)}{w_i} \right\rangle_w \sum_i w_i$$

- We can evaluate this expression approximately by selecting a set of  $N$  sample states randomly but non-uniformly such that the probability of choosing a state  $i$  is:

$$p_i = \frac{w_i}{\sum_j w_j}$$

- We can evaluate this expression approximately by selecting a set of  $N$  sample states randomly but non-uniformly such that the probability of choosing a state  $i$  is:

$$p_i = \frac{w_i}{\sum_j w_j}$$

- In this case, we get:

$$\langle X \rangle \simeq \frac{1}{N} \sum_{k=1}^N \frac{X_k P(E_k)}{w_k} \sum_i w_i$$

Note that the first sum is only over the states that we sample, but the second sum is over all states  $i$  and has to be calculated analytically!



- Our goal is to choose  $w_i$  so that most of the samples are in the region where  $P(E_i)$  is big and very few in the region where it is small.

- Our goal is to choose  $w_i$  so that most of the samples are in the region where  $P(E_i)$  is big and very few in the region where it is small.
- We can choose  $w_i = P(E_i)$ , in which case the above condition is satisfied as well as  $\sum_i w_i = 1$ .

- Our goal is to choose  $w_i$  so that most of the samples are in the region where  $P(E_i)$  is big and very few in the region where it is small.
- We can choose  $w_i = P(E_i)$ , in which case the above condition is satisfied as well as  $\sum_i w_i = 1$ .
- In other words, we just choose  $N$  states in proportion to their Boltzmann probabilities and take the average of  $X$  over them..

- Our goal is to choose  $w_i$  so that most of the samples are in the region where  $P(E_i)$  is big and very few in the region where it is small.
- We can choose  $w_i = P(E_i)$ , in which case the above condition is satisfied as well as  $\sum_i w_i = 1$ .
- In other words, we just choose  $N$  states in proportion to their Boltzmann probabilities and take the average of  $X$  over them..
- Unfortunately, we are not done yet – The catch is that it is not easy to pick states with probability  $P(E_i)$ . This is because to calculate  $P(E_i)$ , we need to know the partition function,  $Z$ .

- Remarkably, it turns out that we can choose states with probability  $P(E_i)$ , without knowing the partition function, using a device called Markov Chain.

- Remarkably, it turns out that we can choose states with probability  $P(E_i)$ , without knowing the partition function, using a device called Markov Chain.
- In this, we want states for the sum  $\langle X \rangle = \frac{1}{N} \sum_{k=1}^N X_k$ , which we will get by generating a Markov chain.

- Remarkably, it turns out that we can choose states with probability  $P(E_i)$ , without knowing the partition function, using a device called Markov Chain.
- In this, we want states for the sum  $\langle X \rangle = \frac{1}{N} \sum_{k=1}^N X_k$ , which we will get by generating a Markov chain.
- Consider a single step in the process – Suppose that the previous state for the step before this one, was state  $i$ .

- Remarkably, it turns out that we can choose states with probability  $P(E_i)$ , without knowing the partition function, using a device called Markov Chain.
- In this, we want states for the sum  $\langle X \rangle = \frac{1}{N} \sum_{k=1}^N X_k$ , which we will get by generating a Markov chain.
- Consider a single step in the process – Suppose that the previous state for the step before this one, was state  $i$ .
- For the new state, instead of choosing randomly, we will make some change (usually small) to the state  $i$  so as to create a new state.



- Remarkably, it turns out that we can choose states with probability  $P(E_i)$ , without knowing the partition function, using a device called Markov Chain.
- In this, we want states for the sum  $\langle X \rangle = \frac{1}{N} \sum_{k=1}^N X_k$ , which we will get by generating a Markov chain.
- Consider a single step in the process – Suppose that the previous state for the step before this one, was state  $i$ .
- For the new state, instead of choosing randomly, we will make some change (usually small) to the state  $i$  so as to create a new state.
- The choice of the new state is determined probabilistically by a set of transition probabilities  $T_{ij}$  that give the probability of changing from state  $i$  to  $j$ .

- If we choose  $T_{ij}$  correctly, we can arrange that the probability of visiting any particular state on any step of the Markov chain to be precisely the Boltzmann probability,  $P(E_i)$ .

- If we choose  $T_{ij}$  correctly, we can arrange that the probability of visiting any particular state on any step of the Markov chain to be precisely the Boltzmann probability,  $P(E_i)$ .
- When we take many steps and generate the entire chain, the complete set of states that we move through is a correct sample of the Boltzmann distribution and we can average any quantity we like over these states.

- The trick lies in choosing  $T_{ij}$ :

$$\sum_j T_{ij} = 1$$

and also

$$\frac{T_{ij}}{T_{ji}} = \frac{P(E_j)}{P(E_i)} = \frac{e^{-\beta E_j} / Z}{e^{-\beta E_i} / Z} = e^{-\beta(E_j - E_i)}$$

In other words, we are choosing a particular value for the ratio of the probability to go from  $i$  to  $j$  and the probability to go back from  $j$  to  $i$ .

- The trick lies in choosing  $T_{ij}$ :

$$\sum_j T_{ij} = 1$$

and also

$$\frac{T_{ij}}{T_{ji}} = \frac{P(E_j)}{P(E_i)} = \frac{e^{-\beta E_j} / Z}{e^{-\beta E_i} / Z} = e^{-\beta(E_j - E_i)}$$

In other words, we are choosing a particular value for the ratio of the probability to go from  $i$  to  $j$  and the probability to go back from  $j$  to  $i$ .

- The Boltzmann distribution is a fixed point of the Markov chain.

- While this is all good – we still need to figure out what  $T_{ij}$  should be.

# Metropolis algorithm

- While this is all good – we still need to figure out what  $T_{ij}$  should be.
- The most common choice is the choice that leads to the Metropolis algorithm.

# Metropolis algorithm

- While this is all good – we still need to figure out what  $T_{ij}$  should be.
- The most common choice is the choice that leads to the Metropolis algorithm.
- Note that we are allowed to visit the same state more than once in the Markov chain..



# Metropolis algorithm

- While this is all good – we still need to figure out what  $T_{ij}$  should be.
- The most common choice is the choice that leads to the Metropolis algorithm.
- Note that we are allowed to visit the same state more than once in the Markov chain..
- Suppose we generate a new state  $j$  after making some change in state  $i$ . We chose the particular change we make uniformly at random from a specified set of possible changes – called the move set.

# Metropolis algorithm

- While this is all good – we still need to figure out what  $T_{ij}$  should be.
- The most common choice is the choice that leads to the Metropolis algorithm.
- Note that we are allowed to visit the same state more than once in the Markov chain..
- Suppose we generate a new state  $j$  after making some change in state  $i$ . We chose the particular change we make uniformly at random from a specified set of possible changes – called the move set.
- We will accept or reject the new state with probability  $P_a$ :

$$P_a = \begin{cases} 1 & \text{if } E_j \leq E_i, \\ e^{-\beta(E_j - E_i)} & \text{if } E_j > E_i \end{cases}$$

- While this is all good – we still need to figure out what  $T_{ij}$  should be.
- The most common choice is the choice that leads to the Metropolis algorithm.
- Note that we are allowed to visit the same state more than once in the Markov chain..
- Suppose we generate a new state  $j$  after making some change in state  $i$ . We chose the particular change we make uniformly at random from a specified set of possible changes – called the move set.
- We will accept or reject the new state with probability  $P_a$ :

$$P_a = \begin{cases} 1 & \text{if } E_j \leq E_i, \\ e^{-\beta(E_j - E_i)} & \text{if } E_j > E_i \end{cases}$$

- This scheme will satisfy all the criterion of the transition probability:  $\frac{T_{ij}}{T_{ji}} = e^{-\beta(E_i - E_j)}$

- 1 Choose a random starting state.

# Metropolis Algorithm

- 1 Choose a random starting state.
- 2 Choose a move uniformly at random from an allowed set of moves.

# Metropolis Algorithm

- 1 Choose a random starting state.
- 2 Choose a move uniformly at random from an allowed set of moves.
- 3 Calculate the probability,  $P_a$  to accept or reject the move.

- 1 Choose a random starting state.
- 2 Choose a move uniformly at random from an allowed set of moves.
- 3 Calculate the probability,  $P_a$  to accept or reject the move.
- 4 With probability  $P_a$ , accept the move – ie system changes to the new state; OR reject the move – ie system stays in the current state.

- 1 Choose a random starting state.
- 2 Choose a move uniformly at random from an allowed set of moves.
- 3 Calculate the probability,  $P_a$  to accept or reject the move.
- 4 With probability  $P_a$ , accept the move – ie system changes to the new state; OR reject the move – ie system stays in the current state.
- 5 Measure the value of the quantity of interest and add it to the running sum.



- 1 Choose a random starting state.
- 2 Choose a move uniformly at random from an allowed set of moves.
- 3 Calculate the probability,  $P_a$  to accept or reject the move.
- 4 With probability  $P_a$ , accept the move – ie system changes to the new state; OR reject the move – ie system stays in the current state.
- 5 Measure the value of the quantity of interest and add it to the running sum.
- 6 Go to step 2.





- The steps where you reject the move that don't change the state of the system do count as steps.

- The steps where you reject the move that don't change the state of the system do count as steps.
- The number of moves that you chose to take you from  $i$  to  $j$  should be the same as from  $j$  to  $i$ .

- The steps where you reject the move that don't change the state of the system do count as steps.
- The number of moves that you chose to take you from  $i$  to  $j$  should be the same as from  $j$  to  $i$ .
- One has to choose a move set such that every possible state is reachable.

- The steps where you reject the move that don't change the state of the system do count as steps.
- The number of moves that you chose to take you from  $i$  to  $j$  should be the same as from  $j$  to  $i$ .
- One has to choose a move set such that every possible state is reachable.
- That the Markov chain will go to Boltzmann distribution is proved but, how long will it take to reach equilibrium is not known.

$$E = - \sum_{\langle ij \rangle} s_i s_j$$



- If we are interested in the ground state energy of a system (classical or quantum), we can use simulated annealing.

- If we are interested in the ground state energy of a system (classical or quantum), we can use simulated annealing.
- The probability that the system is in a state  $i$  is:

$$P(E_i) = \frac{e^{-\beta E_i}}{Z} \quad Z = \sum_i e^{-\beta E_i}$$



- If we are interested in the ground state energy of a system (classical or quantum), we can use simulated annealing.
- The probability that the system is in a state  $i$  is:

$$P(E_i) = \frac{e^{-\beta E_i}}{Z} \quad Z = \sum_i e^{-\beta E_i}$$

- Let us assume that the system has a single unique ground state and let us choose our energy scale such that  $E_i = 0$  in the ground state and  $E_i > 0$  for all others.

- If we are interested in the ground state energy of a system (classical or quantum), we can use simulated annealing.
- The probability that the system is in a state  $i$  is:

$$P(E_i) = \frac{e^{-\beta E_i}}{Z} \quad Z = \sum_i e^{-\beta E_i}$$

- Let us assume that the system has a single unique ground state and let us choose our energy scale such that  $E_i = 0$  in the ground state and  $E_i > 0$  for all others.
- Then in the limit  $T \rightarrow 0$ ,  $\beta \rightarrow \infty$  and  $e^{-\beta E_i} \rightarrow 0$  except in the ground state where  $e^0 = 1$ .

- If we are interested in the ground state energy of a system (classical or quantum), we can use simulated annealing.
- The probability that the system is in a state  $i$  is:

$$P(E_i) = \frac{e^{-\beta E_i}}{Z} \quad Z = \sum_i e^{-\beta E_i}$$

- Let us assume that the system has a single unique ground state and let us choose our energy scale such that  $E_i = 0$  in the ground state and  $E_i > 0$  for all others.
- Then in the limit  $T \rightarrow 0$ ,  $\beta \rightarrow \infty$  and  $e^{-\beta E_i} \rightarrow 0$  except in the ground state where  $e^0 = 1$ .
- This in this limit:

$$P(E_i) = \begin{cases} 1 & \text{for } E_i = 0 \\ 0 & \text{for } E_i > 0 \end{cases}$$

- One of the ways to find the ground state is then to slowly lowering the temperature, and see what state the system lands in.

- One of the ways to find the ground state is then to slowly lowering the temperature, and see what state the system lands in.
- As one lowers the temperature, the system should land in the ground-state!