



**University College Dublin**

**An Coláiste Ollscoile, Baile Átha Cliath**

# **PHYC20040 Exploring the Solar System**

Experiment No.3 The Classification of Stellar Spectra

**26 February 2025**

by Joana C.C. Adao (Student No. 23311051)



# Contents

<b>Abstract</b>	<b>2</b>
<b>1 Theory</b>	<b>2</b>
1.1 Spectra . . . . .	2
1.1.1 Absorption Line Spectra . . . . .	3
1.1.2 Emission Line Spectra . . . . .	3
1.2 Blackbody Radiation . . . . .	3
1.2.1 Wien's Displacement Law . . . . .	3
1.3 Apparent and Absolute Magnitude . . . . .	4
1.4 Stellar Classification . . . . .	4
1.4.1 Harvard Classification of Spectral Types . . . . .	4
1.4.2 MK System of Luminosity . . . . .	5
1.4.3 Main Sequence Stars . . . . .	6
<b>2 Methodology</b>	<b>6</b>
<b>3 Results and Calculations</b>	<b>8</b>
3.1 Spectral Classification (CLEA List) . . . . .	8
3.1.1 SIMBAD Analysis . . . . .	9
3.1.2 Jacoby-Hunter-Christian Atlas Analysis . . . . .	9
3.1.3 A Comment on SAO 81292 . . . . .	9
3.2 Spectral Classification (Telescope) . . . . .	10
3.2.1 SIMBAD Analysis . . . . .	11
3.2.2 (ADDITIONAL) Spectral Classification (SIMBAD Coordinates) . . . . .	11
<b>4 Conclusion</b>	<b>12</b>
<b>References</b>	<b>13</b>
<b>List of Figures</b>	<b>14</b>
<b>List of Tables</b>	<b>14</b>
<b>Appendix</b>	<b>15</b>
Code . . . . .	15
CLEA Atlas of Standard Spectra . . . . .	16


## Abstract

This experiment made use of the CLEA (Contemporary Laboratory Experience in Astronomy) 'Classifying Stellar Spectra' software to practice classifying a star's spectral type. The program had a list of 25 pre-determined stars that were analysed and classified on their spectral type based off of their absorption spectra when compared to the known spectra of main sequence stars from O-type to M-type. When these estimated spectral types were cross-checked to the spectral types as defined by the SIMBAD database and JHC Atlas, **30.4%** and **45.8%** of the results matched, respectively. The same process was done for randomly selected stars in the simulated nightsky offered by CLEA, and these chosen stars were compared to ones available on the SIMBAD database via input of their coordinates.

# 1 Theory

## 1.1 Spectra

The electromagnetic spectrum describes all forms of electromagnetic radiation, including visible light, as it varies with wavelength and frequency [1, 2]. Spectroscopes are equipments used to visually observe the spectra and spectrographs photograph and map the studied spectra [1]. There are three main ways that the spectra can be classified, as illustrated in figure 1 [3].



**Figure 1:** Types of spectra: continuous, emission, absorption [3]

**Spectrometry** puts quantitative values to the theory explored by spectroscopy (*theoretical* interaction between radiation and matter) by measuring the interactions between light (electromagnetic radiation) and matter [4].

### 1.1.1 Absorption Line Spectra


The **absorption spectrum** is measured when light from a continuous source, like a star, passes through a cloud of cooler gas. The wavelengths that will be absorbed depend on the composition and elements of the gas, as well as its temperature and density. The wavelengths that didn't pass through are what appear as dark lines on the continuous spectrum known as the absorption line spectrum [5].

### 1.1.2 Emission Line Spectra

The **emission spectrum** is measured when atoms become excited after light, like from a star, passes through a cloud of gas. The light can heat up the cloud of gas, thus exciting the atoms and causing them to release light. The light which the gas releases depends on the composition and elements, temperature, and density of the gas cloud. The light emitted will appear as coloured lines.

## 1.2 Blackbody Radiation

Blackbodies are idealised surfaces that can absorb any wavelength of incident radiation without any reflection, and that can emit electromagnetic (EM) radiation at maximum possible monochromatic intensities for a range of wavelengths (a continuous spectrum) based on temperature [6, 7, 8].



**Figure 2:** Diagram of blackbody radiation curves for stars of 10 000K, 6 000K, 3 500K, with the visible light spectrum shown on the x-axis [8].

The characteristic curve of the continuous spectrum of blackbody radiation can be used to determine the temperatures of stars and other cosmic objects by finding their "colour" [8]. Hotter stars emit shorter wavelengths and therefore appear bluer, while colder stars emit longer wavelengths and therefore appear redder (see figure 4).

### 1.2.1 Wien's Displacement Law

Wien's Law relates the peak wavelength of a blackbody's peak emission spectrum ( $\lambda_{\text{peak}}$ ) to temperature ( $T$ ) [9]. This is given by the following, with  $\lambda_{\text{peak}}$  in metres (m) and temperature  $T$  in Kelvin (K) [9]:

$$\lambda_{\text{peak}}(\text{m}) = \frac{2.89777 \times 10^{-3}}{T} \quad (1)$$

The wavelength  $\lambda_{\text{peak}}$  can also be found in Ångstroms (Å) by changing the numerator constant to be in Ångstrom-Kelvin instead of metre-Kelvin:

$$\lambda_{\text{peak}}(\text{Å}) = \frac{2.9 \times 10^7}{T} \quad (2)$$

### 1.3 Apparent and Absolute Magnitude

Magnitude is a logarithmic measure of the brightness of a star or other celestial body. The brighter the object, the lower the magnitude (number), and they can be negative for particularly bright stars [10]. The magnitude of the celestial objects are divided into two types of observation:

- **Apparent magnitude,  $m$** , is used to describe how bright a celestial object appears from the view on Earth [11].
- **Absolute magnitude,  $M$** , also in reference to *luminosity*, is defined as the magnitude of the star if the distance between it and Earth were 10 parsecs (pc) [12, 13]. When at a set distance, astronomers are then able to compare intrinsic brightness of stars [13].

Absolute ( $M$ ) and apparent ( $m$ ) magnitudes can be used in equation 3 to calculate  $D$ , the distance in parsecs (pc). The distance modulus is then  $m - M$ . The magnitudes do not have units [13].

$$M = m + 5 - 5(\log_{10} D) \quad , \quad m - M = 5 \log_{10} \left( \frac{D}{10} \right) \quad (3)$$


The above equation (3) can then be manipulated to find the distance,  $D$ :

$$\log_{10} D = \frac{m - M + 5}{5} \quad \implies \quad D = 10^{\frac{m - M + 5}{5}} \quad (4)$$

### 1.4 Stellar Classification

#### 1.4.1 Harvard Classification of Spectral Types

The currently-used system of classification of stars was created by a team at Harvard in 1924 [14]. The different classes are **OBAFGKM**, left to right: hottest to coolest. The different classes and temperature thresholds are illustrated in figure 3.



**Figure 3:** Chart for classifying of the main star types as per the Harvard classification [15].

Subclasses of numbers ranging from 0-9 allow for finer distinctions in temperature within the OBAFGKM classes. For example, the Sun is spectral type **G2** with a temperature of about 5 700 K (Kelvin) [16].

### 1.4.2 MK System of Luminosity

The Morgan-Keega (MK) Luminosity Class is an addition to the Harvard stellar classification scheme (see §1.4.1), given in the form of (primarily) Roman numerals. This addition to the system was devised in order to be able to distinguish between stars of similar temperatures but differing *luminosities*.

The MK System was originally from I to V, but has since been expanded to differentiate more between I-type stars and further beyond V. The table containing the luminosity classifications is shown below [17]:


**Table 1:** Table of the MK Luminosity Classes [17].

Class	Star Type
Ia-O	extremely luminous supergiants
Ia	luminous supergiants
Ib	less luminous supergiants
II	bright giants
III	normal giants
IV	subgiants
V	main sequence dwarf stars
VI, or sd	subdwarfs
D	white dwarfs

### 1.4.3 Main Sequence Stars

Main sequence stars have nuclear reactions at their core through the fusion of hydrogen atoms into helium that allow them to sustain themselves [18, 19, 20]. The radiation pressure from the nuclear reactions and the gravitational pressure work against each other in such a way that the star remains stable [19] in a process known as **hydrostatic equilibrium** [18].

Figure 4 shows the graph of the absorption line spectrum for specific main sequence stars ranging from A to M, excluding classes B and O.



**Figure 4:** Stellar spectra of the main sequence (V) stars (A5 to M0) [21].

## 2 Methodology

This experiment made use of the *Contemporary Laboratory Experience in Astronomy*, CLEA, software and the *Classification of Stellar Spectra* program included in the package. CLEA is an online software developed to simulate and illustrate modern astronomical techniques in a 'real' night sky. This experiment is divided into two separate yet related exercises. [22].

For the first part of the experiment, graphs of different *absorption spectra* for 25 unknown stars from a list were compared and contrasted against the absorption spectra of main sequence stars of 13 different spectral types sourced from an Atlas of Standard Spectra. By comparing the known to the unknown, the spectral type for the unknown stars could be approximated and noted. By identifying the approximate spectral type, astronomers could then estimate the temperature of the star.

The program offered the simulated use of a spectrometer attached to a 0.4m (16") research telescope for the second part of the experiment in which ideally 3 stars are found and studied. The aim is to find at least both a bright and dim star and utilise the spectrometer attached to the telescope to produce an absorption spectrum for that star. From there, the same as was done for the first part of the experiment can be done and the spectral type of the star can be estimated by comparing and contrasting to the Atlas of Standard Spectra provided within the software [23].

Windows for both programs of the experiment are illustrated in figure 5.




Figure 5: Screenshot of the Stellar Spectra CLEA program.

The spectral type identification program allowed also for the use of different visualisation and data analysis methods (not pictured):

- **Grayscale Photo:** this setting showed a photographed representation of the spectrum in black and white.
- **Graphical Trace:** default setting, graphical representation of the spectra as pictured in figure 5.
- **Comb. (Photo & Trace):** combination of both the above settings; centre panel shows the photographic representation of the unknown star whilst the bottom panel shows the graphical representation.

These settings are useful to help differentiate between the strongest dips/lines of that spectra when it is not very clear on another representation.



### 3 Results and Calculations

#### 3.1 Spectral Classification (CLEA List)

The estimated spectral types for the list of the 25 stars provided by CLEA has been formatted into a compact table below, table 2, with reasons for why that spectral type was chosen for that star:

**Table 2:** Table of the estimated spectral class for the CLEA star list, with reasons listed.

STAR	SP TYPE	REASONS
HD 124320	A3	HI lines v. strong; Ca II line betw. A1 and A5
HD 37767	B3	Strong Ca II lines, He II and HI strong; betw. B0 and B6
HD 35619	O7	Strong cont. spectrum, weak dips; betw. O5 and B0
HD 23733	F3	Ca II lines v. strong; betw. F0 and F5
O1015	B8	HI lines similar to B6, slightly stronger dips; betw. B6 and A1
HD 24189	G3	Strong Ca II lines, strong He II line; betw. G0 and G6
HD 107399	F7	He I line similar to G0; betw. F5 and G0
HD 240344	B4	Strong HI line; betw. B0 and B6
HD 17647	G5	Weak HI line, strong Ca II (K) line; betw. G0 and G6
BD +63 137	M1	Strong dip at 4227 Å; betw. M0 and M5
HD 66171	G5	Strong Ca II lines, weaker HI (H-gamma); betw. G0 and G6
HZ 948	F7	Strong HI line, minimal difference slope; betw. F5 and G0
HD 35215	B3	Strong HI (H-epsilon), minor difference slope; betw. B0 and B6
Feige 40	B4	Strong HI (H-delta); betw. B0 and B6
Feige 41	A0	Strong HI (H-epsilon, H-gamma, H-delta) lines; betw. B6 and A1
HD 6111	F8	Strong Ca II line, strong HI (H-delta) line; betw. F5 and G0
HD 23863	A5	Smoother cont. spectrum, otherwise identical to A5
HD 221741	A3	Strong Ca II lines, smoother cont. spectrum; betw. A1 and A5
HD 242936	O5	Nearly identical to O5
HD 5351	K4	Strong Fe I lines and surrounding dips; betw. K0 and K5
SAO 81292	M3	(EMISSION) v. strong Ca II lines, strong HI (H-gamma); betw. M5 and M0
HD 27685	G6	Nearly identical to G6
HD 21619	A6	Smoother spectrum; nearly A5
HD 23511	F6	Difference slope slightly below axis; nearly F5
HD 158659	B0	V. slightly smoother cont. spectrum; nearly identical to B0

The 'HD' prefixed stars are sourced from the *Henry-Draper* catalogue, 'BD' prefixed stars are sourced from the *Bonner Durchmusterung* catalogue, 'HZ' prefixed stars are sourced from the *Humason-Zwicky* catalogue, 'Feige' prefixed stars are sourced from the *Feige* catalogue, and

'SAO' prefixed stars are sourced from the *Smithsonian Astrophysical Observatory* catalogue. It is unclear where the 'O' prefixed stars are sourced from.

### 3.1.1 SIMBAD Analysis

The SIMBAD Astronomical Database is made of use to cross-check the stars provided and selected [24]. As of the 2nd of March 2025, SIMBAD currently has 20 202 386 objects registered on the database.

**Table 3:** Table of the actual spectral types of the CLEA star list as per SIMBAD.

STAR	SP TYPE (SIMBAD)	STAR	SP TYPE (SIMBAD)	STAR	SP TYPE (SIMBAD)	STAR	SP TYPE (SIMBAD)	STAR	SP TYPE (SIMBAD)
HD 124320	A5V	HD 24189	F6V	HD 66171	G2V	HD 6111	G5 E	SAO 81292	dM3 C
HD 37767	B3V	HD 107399	G0 E	HZ 948	N/A	HD 23863	A7Vn C	HD 27685	G2 E
HD 35619	O7.5V	HD 240344	B7 E	HD 35215	B1V	HD 221741	A2 E	HD 21619	A6V C
HD 23733	A9V	HD 17647	G5 D	Feige 40	B4V	HD 242936	A2V	HD 23511	F5V C
O1015	N/A	BD +63 137	K7V	Feige 41	A1V	HD 5351	K4V C	HD 158659	B5(Ib/II) D

Out of the 23 stars available on the SIMBAD database from the 25 of the CLEA list, **30.4%** matched what was estimated.

### 3.1.2 Jacoby-Hunter-Christian Atlas Analysis

The Jacoby-Hunter-Christian (JHC) Atlas is made of use to cross-check the stars provided [25]. The JHC Atlas was originally published in 1984 and contains 161 spectra of stars of spectral classes O to M for luminosity classes V, III, and I.


**Table 4:** Table of the actual spectral types of the CLEA star list as per the JHC Atlas.

STAR	SP TYPE (JHC)	STAR	SP TYPE (JHC)	STAR	SP TYPE (JHC)	STAR	SP TYPE (JHC)	STAR	SP TYPE (JHC)
HD 124320	A2V	HD 24189	F6V	HD 66171	G2V	HD 6111	F8V	SAO 81292	M4.5 Ve
HD 37767	B3V	HD 107399	F9V	HZ 948	F3V	HD 23863	A7V	HD 27685	G7V
HD 35619	O7V	HD 240344	B4V	HD 35215	B1.5V	HD 221741	A3V	HD 21619	A6V
HD 23733	A9V	HD 17647	G1V	Feige 40	B4V	HD 242936	N/A	HD 23511	F4V
O1015	B8V	BD +63 137	M1V	Feige 41	A1V	HD 5351	K4V	HD 158659	B0V

Out of the 24 stars available on the JHC Atlas from the 25 of the CLEA list, **45.8%** matched what was estimated.


### 3.1.3 A Comment on SAO 81292

During the experiment star SAO 81292 was identified to have emission lines (see figure 6).



**Figure 6:** Image of the grayscale spectrum for star SAO 81292.

This could indicate a particularly active late-stage star or a flare star. Through analysis of the lightcurve graph from data gathered by TESS, we can indeed infer that this star is a flare star (see figure 7).



**Figure 7:** Lightcurve graph plot of the SAO 81292 star, TESS-SPOC data (2023) [26].

The flux of light (y-axis) has been gathered over a period of 20 seconds (x-axis), and in that period 4 spikes in the flux can be seen, indicating flares. Known flare stars have spectral types of late-K through late-M, which is consistent with the information found on both the SIMBAD database and the JHC Atlas. Additionally, flare stars often have detectable Ca II and H lines, which is also consistent with what was observed from the experiment [27]. As such, SAO 81292 is most likely a flare star and hence the emission lines visible on the spectra in figure 6.

### 3.2 Spectral Classification (Telescope)

Three stars, one dim and two bright, were then located on the vast simulated expanse of the sky and the spectrographs for each star was found. Stars were chosen to be relatively far away from one another and the findings were formatted into a compact table, table 5, below with the same structure for spectral type and reasoning as before:

**Table 5:** Table of the estimated spectral types for self-selected 1 dim and 2 bright stars.

Star #	Star Name	RA (h m s)	DE (d m s)	App. Mag. (m)	S/N Ratio	SP Type	Reasons
1	436	6 14 20	33 21 24	4.03	1001	G7	Strong Ca II lines, strong G-band; betw. G6 and K0
2	57	6 22 28	30 52 21	14.65	45	A6	Sharper dips, noisy (S/N ratio); betw. A5 and F0
3	49	6 07 19	30 12 32	5.98	879	B5	Strong H I (H-epsilon, H-delta, H-gamma) lines, stronger Fe I line; betw. B0 and B6

It was important to get the S/N (signal to noise) ratio of each spectrum to be as high as possible (higher number) so the spectra would not be as noisy. If the spectrum is noisy, it is much harder to identify the most prominent absorption lines. This can be helped by choosing to use a larger telescope (CLEA offers both 1.0m and 4.0m telescopes to be used) as this way the telescope can collect more light quicker, reducing the noise present in the spectrograph.

### 3.2.1 SIMBAD Analysis

Much like before, the SIMBAD database can be made of use to cross-check the stars found on the digitally simulated nightsky provided by CLEA by inputting the RA and DE coordinates found.

**Table 6:** Stars found through telescope cross-matched on SIMBAD to a radius of 5 arc min.

Star #	Star Name	RA (h m s)	DE (d m s)	Mag. (unspecified)	SP Type (SIMBAD)
1	HD 253515	6 14 <b>26</b>	+33 21 <b>01</b>	11.3	kA1hA2mA3
2	HD 44300	6 22 28	+30 <b>56 43</b>	8.82	F0
3	EM* MWC 790	6 07 <b>24</b>	+30 <b>11 44</b>	12.77	Be

The change in coordinates from those gathered has been highlighted in bold, and only the results where the spectral type was provided have been taken (as it is the aim of the experiment).

According to SIMBAD, star 1 (HD 253515) is of spectral type kA1hA2mA3. This is an Am (metallic-lined) A star with Ca II K-line bands corresponding to that of an A1 spectral type, Hydrogen (H) lines corresponding to that of an A2 spectral type, and metallic lines (such as Iron (Fe)) corresponding to that of an A3 spectral type. Am stars are chemically peculiar, hence the deviation from standard spectral type classification.

Also according to SIMBAD [28], the lowercase 'e' that follows star 3's (EM\* MWC 790) spectral type B indicates emission lines in the spectrum.

### 3.2.2 (ADDITIONAL) Spectral Classification (SIMBAD Coordinates)

With reference to these new SIMBAD coordinates instead, the telescope was accessed again (not in the laboratory) and the experiment was done again to cross-check. As this new rendition of the experiment was not done in the laboratory, the newest version of the CLEA VIREO (VIRtual Educational Observatory) software was able to be accessed. This version allows for a much wider expanse of the simulated nightsky to be observed, the following results were obtained:

**Table 7:** Estimated spectral types of stars as per the SIMBAD approximate coordinates (non-laboratory rendition).

Star #	Object Name	RA (h m s)	DE (d m s)	App. Mag. (m)	S/N Ratio	SP Type	Reasons
1	N3000-01024	6 14 26.1	+33 21 01	11.63	521	B8	Strong HI (H-gamma) line, strong He II line; betw. B6 and A1
2	N3000-00640	6 22 28.4	+30 56 43	8.35	1000	F0	Identical to F0
3	N3000-01059	6 07 23.7	+30 11 42	11.67	620	A3	Slightly weaker Ca II (K) line; betw. A1 and A5

This addition to the experiment was not necessary, but merely a means to compare the data gathered from the SIMBAD database to what would have been obtained from the laboratory experiment. Identical approaches were taken, different softwares and settings.

## 4 Conclusion

Through the use of the CLEA software, this experiment was successful in practicing classifying spectral types of stars based on this spectra. When compared to two different databases, it was found that **30.4%** of the estimated spectral types matched with the SIMBAD database, while **45.8%** of the estimated spectral types matched with the JHC Atlas. The fact that these results do not match could be due to discrepancies in data-gathering methods and complexities of the stars, or classification threshold differences as SIMBAD has a lot more variable descriptors for spectral types.

For the telescope-based section of the experiment, the same approach as before was taken. The factor of the signal-to-noise ratio meant that there was likely some error in estimating the spectral type for stars as some spectral types are more reliant on the 'smoothness' of the spectrum. Higher ratios also meant more noise, making it difficult to tell what were the most prominent lines of the absorption spectrum. This issue could be fixed by making use of the larger telescopes available within the software.

# References

- [1] E. Gregersen. (2024, November) spectrum. [Accessed 3 February 2025]. [Online]. Available: <https://www.britannica.com/science/spectrum>
- [2] NASA. (2022, September) The Electromagnetic Spectrum. [Accessed 3 February 2025]. [Online]. Available: <https://hubblesite.org/contents/articles/the-electromagnetic-spectrum>
- [3] L. Hustak. (2021, July) Spectra Picture Credit. [Accessed 2 February 2025]. [Online]. Available: <https://webbtelescope.org/contents/media/images/01F8GF8DK2PRY4FP9DA2XPQC8S>
- [4] Anon. (2020, January) SPECTROMETRY AND SPECTROSCOPY: WHAT'S THE DIFFERENCE? [Accessed 3 February 2025]. [Online]. Available: <https://www.atascientific.com.au/spectrometry/>
- [5] —. (n.d.) Absorption Line. [Accessed 3 February 2025]. [Online]. Available: <https://astronomy.swin.edu.au/cosmos/a/absorption+line>
- [6] OpenStax. (n.d.) Blackbody Radiation. [Accessed 1 March 2025]. [Online]. Available: [https://phys.libretexts.org/Bookshelves/University\\_Physics/University\\_Physics\\_\(OpenStax\)/University\\_Physics\\_III\\_-\\_Optics\\_and\\_Modern\\_Physics\\_\(OpenStax\)/06%3A\\_Photons\\_and\\_Matter\\_Waves/6.02%3A\\_Blackbody\\_Radiation](https://phys.libretexts.org/Bookshelves/University_Physics/University_Physics_(OpenStax)/University_Physics_III_-_Optics_and_Modern_Physics_(OpenStax)/06%3A_Photons_and_Matter_Waves/6.02%3A_Blackbody_Radiation)
- [7] Anon. (2004-2005) Blackbody Radiation. [Accessed 1 March 2025]. [Online]. Available: <https://maths.ucd.ie/met/msc/fezzik/Phys-Met/Ch04-2-Slides.pdf>
- [8] ESA. (2019, September) BLACKBODY RADIATION. [Accessed 1 March 2025]. [Online]. Available: <https://sci.esa.int/web/education/-/48986-blackbody-radiation>
- [9] P. S. Gabriel. (n.d.) Derivation and Application of Wien's Displacement Law. Department of Physics, Columbia University, New York, NY 10027. [Accessed 1 March 2025]. [Online]. Available: [https://www.spacefirst.org/uploads/7/9/1/7/79174470/derivation\\_and\\_applications\\_of\\_wien\\_s\\_displacement\\_law\\_\\_2\\_.pdf](https://www.spacefirst.org/uploads/7/9/1/7/79174470/derivation_and_applications_of_wien_s_displacement_law__2_.pdf)
- [10] E. Gregersen. (2018, April) magnitude. [Accessed 4 February 2025]. [Online]. Available: <https://www.britannica.com/science/magnitude-astronomy>
- [11] Anon. (n.d.) Apparent Magnitude. [Accessed 4 February 2025]. [Online]. Available: <https://lco.global/spacebook/distance/what-apparent-magnitude/>
- [12] —. (n.d.) Absolute Magnitude. [Accessed 4 February 2025]. [Online]. Available: <https://lco.global/spacebook/distance/what-absolute-magnitude/>
- [13] —. (n.d.) Absolute Magnitude. [Accessed 4 February 2025]. [Online]. Available: <https://astronomy.swin.edu.au/cosmos/a/Absolute+Magnitude>
- [14] —. (1924) A notes on the spectral atlas and spectral classification. [Accessed 15 February 2025]. [Online]. Available: <https://lweb.cfa.harvard.edu/~pberlind/atlas/htmls/note.html>
- [15] —. (n.d.) Stellar classification. [Accessed 15 February 2025]. [Online]. Available: [https://en.wikipedia.org/wiki/Stellar\\_classification](https://en.wikipedia.org/wiki/Stellar_classification)
- [16] —. (n.d.) Spectral type. [Accessed 15 February 2025]. [Online]. Available: <https://astronomy.swin.edu.au/cosmos/S/Spectral+Type>
- [17] —. (n.d.) Morgan-Keenan Luminosity Class. [Accessed 1 March 2025]. [Online]. Available: <https://astronomy.swin.edu.au/cosmos/m/morgan-keenana+luminosity+class>
- [18] —. (n.d.) Stellar Evolution. [Accessed 15 February 2025]. [Online]. Available: <https://www.schoolobservatory.org/learn/space/stars/evolution>
- [19] C. Nettles and K. Chamberlain. (2023, November) Main Sequence Star — Definition, Chart, Characteristics. [Accessed 15 February 2025]. [Online]. Available: <https://study.com/academy/lesson/main-sequence-star-definition-facts-quiz.html>
- [20] NASA. (n.d.) Types of Stars. [Accessed 15 February 2025]. [Online]. Available: <https://science.nasa.gov/universe/stars/types/>
- [21] P. Riaud and S. Jean, "Improving Earth-like planets' detection with an ELT: The differential radial velocity experiment," *Astronomy and Astrophysics - ASTRON ASTROPHYS*, vol. 469, pp. 355–361, 07 2007, [Accessed 1 March 2025]. [Online]. Available: [https://www.researchgate.net/figure/The-figure-shows-various-stellar-spectra-for-the-main-sequence-stars-A5-to-M0-Below\\_fig1\\_248851913](https://www.researchgate.net/figure/The-figure-shows-various-stellar-spectra-for-the-main-sequence-stars-A5-to-M0-Below_fig1_248851913)

- [22] CLEA. (n.d.) The Classification of Stellar Spectra. [Accessed 2 March 2025]. [Online]. Available: <http://public.gettysburg.edu/~marschal/clea/speclab.html>
- [23] UCD. (n.d.) The Classification of Stellar Spectra. [Accessed 2 March 2025]. [Online]. Available: <https://brightspace.ucd.ie/d2l/le/content/300670/viewContent/3521485/View>
- [24] CDS. (2000-2025) SIMBAD Astronomical Database. [Accessed 2 March 2025]. [Online]. Available: <https://simbad.cds.unistra.fr/simbad/>
- [25] Jacoby, Hunter, and Christian. (1984) Jacoby-Hunter-Christian Atlas. [Accessed 2 March 2025]. [Online]. Available: <https://www.stsci.edu/hst/instrumentation/reference-data-for-calibration-and-tools/astronomical-catalogs/jacoby-hunter-christian-atlas>
- [26] TESS. (2019-2023) TIC 95431305. [Accessed 3 March 2025]. [Online]. Available: <https://exofop.ipac.caltech.edu/tess/target.php?id=95431305>
- [27] M. Templeton. (n.d.) UV Ceti and the flare stars. [Accessed 3 March 2025]. [Online]. Available: <https://www.aavso.org/vsots-uvcet>
- [28] CDS. (2000-2025) SIMBAD: Spectral Type Coding. [Accessed 2 March 2025]. [Online]. Available: <http://simbad.cds.unistra.fr/Pages/guide/chD.htx>

## List of Figures

1	Types of spectra: continuous, emission, absorption [3] . . . . .	2
2	Diagram of blackbody radiation curves for stars of 10 000K, 6 000K, 3 500K, with the visible light spectrum shown on the x-axis [8]. . . . .	3
3	Chart for classifying of the main star types as per the Harvard classification [15].	5
4	Stellar spectra of the main sequence (V) stars (A5 to M0) [21]. . . . .	6
5	Screenshot of the Stellar Spectra CLEA program. . . . .	7
6	Image of the grayscale spectrum for star SAO 81292. . . . .	9
7	Lightcurve graph plot of the SAO 81292 star, TESS-SPOC data (2023) [26]. . .	10

## List of Tables

1	Table of the MK Luminosity Classes [17]. . . . .	5
2	Table of the estimated spectral class for the CLEA star list, with reasons listed.	8
3	Table of the actual spectral types of the CLEA star list as per SIMBAD. . . . .	9
4	Table of the actual spectral types of the CLEA star list as per the JHC Atlas. .	9
5	Table of the estimated spectral types for self-selected 1 dim and 2 bright stars. .	10
6	Stars found through telescope cross-matched on SIMBAD to a radius of 5 arc min.	11
7	Estimated spectral types of stars as per the SIMBAD approximate coordinates (non-laboratory rendition). . . . .	11

# Appendix

## Code

```
import matplotlib.pyplot as plt
import lightkurve as lk
%matplotlib inline # thank you to https://www.youtube.com/watch?v=fhQjDv_vuQU

data = lk.search_lightcurve("SAO 81292", mission = "TESS")
data

lc = data.download_all()
lc[6].plot() # from data table, [6]: exptime 20s, year 2023, author SPOC
```


Code for generating the lightcurve graph in figure 7.

#	mission	year	author	exptime s	target_name	distance arcsec
0	TESS Sector 45	2021	<a href="#">SPOC</a>	20	95431305	0.0
1	TESS Sector 46	2021	<a href="#">SPOC</a>	20	95431305	0.0
2	TESS Sector 45	2021	<a href="#">SPOC</a>	120	95431305	0.0
3	TESS Sector 46	2021	<a href="#">SPOC</a>	120	95431305	0.0
4	TESS Sector 48	2022	<a href="#">SPOC</a>	20	95431305	0.0
5	TESS Sector 48	2022	<a href="#">SPOC</a>	120	95431305	0.0
6	TESS Sector 72	2023	<a href="#">SPOC</a>	20	95431305	0.0
7	TESS Sector 72	2023	<a href="#">SPOC</a>	120	95431305	0.0
8	TESS Sector 45	2021	<a href="#">TESS-SPOC</a>	600	95431305	0.0
9	TESS Sector 46	2021	<a href="#">TESS-SPOC</a>	600	95431305	0.0
10	TESS Sector 48	2022	<a href="#">TESS-SPOC</a>	600	95431305	0.0
11	TESS Sector 72	2023	<a href="#">TESS-SPOC</a>	200	95431305	0.0
12	TESS Sector 45	2021	<a href="#">QLP</a>	600	95431305	0.0
13	TESS Sector 46	2021	<a href="#">QLP</a>	600	95431305	0.0
14	TESS Sector 48	2022	<a href="#">QLP</a>	600	95431305	0.0
15	TESS Sector 72	2023	<a href="#">QLP</a>	200	95431305	0.0


Data table sourced with the code from TESS.




# CLEA Atlas of Standard Spectra




Atlas of Standard Spectra: **O5**



Atlas of Standard Spectra: **B0**




Atlas of Standard Spectra: **B6**



Atlas of Standard Spectra: **A1**




Atlas of Standard Spectra: **A5**



Atlas of Standard Spectra: **F0**




Atlas of Standard Spectra: **F5**




Atlas of Standard Spectra: **G0**




Atlas of Standard Spectra: **G6**




Atlas of Standard Spectra: **K0**



Atlas of Standard Spectra: **K5**



Atlas of Standard Spectra: **M0**



Atlas of Standard Spectra: **M5**