



UCD School of Physics

PHYC30170 Physics Astronomy and Space Lab I

Compton Scattering

Joana C.C. Adao

Student No.: 23311051

September 20, 2026

Abstract

This experiment sought to investigate the fundamental aspects of the Compton scattering phenomenon using a Cs-137 gamma-ray source (0.6617 MeV) and a NaI(Tl) scintillation detector. The energy and differential cross-section of the scattered photons were measured at angular range of 20° - 90° and compared to the theoretical predictions to validate the relativistic kinematics and the Klein-Nishina formula. A linear regression of $1/E_{\gamma'}$ against $(1 - \cos \theta)$ yielded an experimental slope of 2.030 ± 0.012 and y-intercept of 1.58 ± 0.007 , with percentage errors of **3.77%** and **4.68%** respectively relative to theoretical calculations of **1.956** and **1.51**. While the plot of the differential cross-section showed a similar trend to the Klein-Nishina theoretical trend, the experimental values remained systematically lower than expected, suggesting systematic errors in the measurements.

Contents

1	Introduction	1
1.1	Kinematics	1
1.2	Differential Cross-Section	1
2	Methodology	1
2.1	Calibration of the Software	2
2.2	Measurement Acquisition	2
3	Results	2
3.1	Kinematics	2
3.2	Differential Cross-Section	3
4	Analysis and Discussion	3
5	Conclusion	4
	Appendix	6
	Tables	6
	Code	6

1. Introduction

Compton scattering describes the inelastic scattering of a photon from a nearly free electron, resulting in a change in the photon's energy, wavelength, and momentum, providing valuable information about how light interacts with matter.




Figure 1: Geometry of Compton Scattering. [1]

The incident (gamma-ray) photon with energy $E_\gamma = h\nu$ collides with an electron of energy $E_{\gamma'} = h\nu'$ assumed to be initially at rest, and the scattered photon deflects at an angle θ with respect to the original direction, illustrated in Figure 1. [2–5]

1.1. Kinematics

The behaviour of the scattered photon follows the energy and momentum conservation laws, which can be expressed as [5]:

$$E_\gamma = E_{\gamma'} + E_e \quad (1)$$

$$(\text{x-dir.}) \quad \frac{h\nu}{c} = \frac{h\nu'}{c} \cos \theta + mv \cos \phi \quad (2)$$

$$(\text{y-dir.}) \quad 0 = \frac{h\nu'}{c} \sin \theta - mv \sin \phi \quad (3)$$

With the use of these conservations of energy and momentum, the energy of the scattered photon can be derived to show [2–6]:

$$E_{\gamma'} = \frac{E_\gamma}{1 + \frac{E_\gamma}{m_0 c^2} (1 - \cos \theta)} \quad (4)$$

where $m_0 c^2$ is the energy of the rest electron (0.511 MeV). This can then be expressed in terms of wavelength shift as [4, 7]:

$$\Delta\lambda = \lambda' - \lambda = \frac{h}{m_0 c} (1 - \cos \theta) \quad (5)$$

where $h/m_0 c$ is the Compton wavelength of the electron (2.426×10^{-10} cm). The maximum wavelength shift thus occurs at $\theta = 180^\circ$ or π radians, and the minimum shift occurs through forward scattering at $\theta = 0$ (0° or radians). [4]

1.2. Differential Cross-Section

The probability of scattering at a given angle is described by the electronic differential cross-section, defined using relativistic quantum mechanics, and given by the Klein-Nishina formula [2–7]:

$$\frac{d\sigma}{d\Omega} = \frac{r_0^2}{2} \left\{ \frac{1 + \cos^2 \theta}{[1 + \alpha(1 - \cos \theta)]^2} \right\} \times \left\{ 1 + \frac{\alpha^2(1 - \cos \theta)^2}{(1 + \cos^2 \theta)(1 + \alpha[1 - \cos \theta])} \right\} \quad (6)$$

where $d\sigma/d\Omega$ is the differential cross-section, r_0 is the classical electron radius (2.8×10^{-13} cm), $\alpha = E_\gamma/m_0 c^2$. As energy increases, the scattering cross-section decreases, indicating that high-energy photons are less likely to scatter than low-energy photons. For low-energy photons ($\alpha \ll 1$) the formula reduces to the non-relativistic Thomson scattering cross-section, which is independent of energy [3, 4, 6, 7]:

$$\frac{d\sigma}{d\Omega} = \frac{r_0^2}{2} (1 + \cos^2 \theta) \quad (7)$$

Additionally, the probability of scattering depends on the number of electrons available in the target material, and thus increases linearly with the atomic number Z as ${}_a\sigma_c = Z {}_e\sigma_c$, where ${}_a\sigma_c$ is the *atomic* Compton scattering cross-section and ${}_e\sigma_c$ is the *electronic* Compton scattering cross-section. [2, 4]

2. Methodology

The experimental setup for Compton scattering, illustrated in Figure 2, used a high activity Cs-137 gamma-ray source (1) housed within lead bricks to minimise background radiation, and a 2"x2" NaI(Tl) scintillation detector (4) was mounted on a pivoting arm to measure the scattered photons at various angles between 20° and 90° . A collimator (3) was used to narrow the gamma-ray beam for detection, and a target of the scattering sample (2) was placed at the pivot point.




Figure 2: Illustration of the experimental setup for Compton scattering with a 2"x2" NaI(Tl) scintillation detector; 1. *Cs-137* source, 2. *Target*, 3. *Collimator*, 4. *Photomultiplier tube (PMT)*, *amplifier*, and *scintillation detector*

The incoming scattered gamma-ray photons were amplified and measured at the PMT once they had interacted with the scintillation detector, and the electrical signals were processed in the *Maestro-32* software to obtain the energy spectra and photopeaks.

2.1. Calibration of the Software

To calibrate the software, a sample of Am-241 and Cs-137 were used with their known photopeak energies of 59.5 keV and 662 keV, respectively, and comparing to the channel numbers of the measured photopeaks. The software created its own calibration curve within it, which was used for all further measurements.

Tweezers were used to handle the portable samples and the samples were placed directly in front of the detector and measured until a peak was seen to form. When not in use, all radioactive samples were stored in a locker in accordance with safety protocols and regulations.

2.2. Measurement Acquisition

To begin measurements, the source aperture was opened, and the detector positioned at the desired angle. The spectrum was recorded until a net count of at minimum 10,000 counts were obtained under the photopeak using the software. Measurements were taken at angle increments of 5°, and the respective net counts, count uncertainties (given by software), peak energies (in keV), and time taken for each measurement were recorded in a table.

Plots for the energy of the scattered photon and the differential cross-section, both as a function of the scattering angle, were plotted and compared graphically to theoretical predictions from Equation 4 and Equation 6 respectively.

3. Results

When calibrating the software, the obtained channels for the Am-241 and Cs-137 photopeaks were **89** and **850** respectively, and inputted into a window within the software to implicitly create a calibration curve. Time taken, scattering angle, and net counts and uncertainties were unnecessary for the calibration and thus not recorded.

3.1. Kinematics

The scattering angles were converted into radians and the energies into MeV for comparison to the theoretical predictions. A plot of $1/E_{\gamma'}$ against $(1 - \cos \theta)$ was made and fitted with a linear regression, shown in Figure 3.




Figure 3: Plot of $1/E_{\gamma'}$ against $(1 - \cos \theta)$ in radians with linear regression fit.

A value for the slope and y-intercept were obtained from the linear fit line, **2.030 ± 0.012** and **1.58 ± 0.007** respectively. Comparing to the theoretical predictions of **1.956** for the slope and **1.51** for the y-intercept, the experimental percentage errors were calculated to be **3.77%** and **4.68%** respectively.

The error on the scattering angle was estimated to be $\pm 2^\circ$ due to obstruction of the angle markings by the pivot arm and difficulty in moving the detector in small, precise increments. The error on the energy was not given by the software and thus not included in the error analysis.

3.2. Differential Cross-Section

From the values obtained for the net counts and time taken for each measurement, a series of calculations were performed to obtain the experimental values for the differential cross-section, which were plotted against the scattering angle and compared to the Klein-Nishina theoretical prediction, shown in Figure 4.




Figure 4: Plot of $d\sigma/d\Omega$ against scattering angle θ in degrees, with theoretical prediction from the Klein-Nishina formula.

The solid angle $\Delta\Omega$ of the detector was calculated using $\Delta\Omega = A/R_2^2$, with A as the area of the detector (5 cm) and R_2 as the distance from the detector to the target (26 cm). The number of electrons in the target was calculated using $N_e = Z_{Al}N_aM/A_{Al}$, where Z_{Al} is the atomic number of aluminium (13), N_a is Avogadro's number, M is the mass of the target (79.3 g), and A_{Al} is the molar mass of aluminium (26.9815 g/mol). The intrinsic efficiency of the detector was calculated using $E_{\text{eff}} = 0.1522(E)^{-1.1325}$, with E as the energy of the scattered photon (MeV). The count rate R was calculated using $R = N/t$, where N is the net counts and t is the time taken per measurement. The cross-section was then calculated using $(\partial\sigma/\partial\Omega) = R/N\Delta\Omega E_{\text{eff}}$.

The experimental values were scaled to the Klein-Nishina prediction at $\theta = 20^\circ$. The shape of the experimental plot follows the same trend as the theoretical prediction but with lower values, which may indicate to systematic errors in the measurements.

4. Analysis and Discussion

The linear relationship observed in Figure 3 is consistent with the theoretical prediction from Equation 4 using the Cs-137 gamma-ray source

(0.6617 MeV). The results obtained for the slope of 2.030 ± 0.012 and y-intercept of 1.58 ± 0.007 are relatively close to the theoretical predictions of 1.956 and 1.51 respectively, with percentage errors of 3.77% and 4.68% respectively.

The likely sources of error in the measurements include the estimated error on the scattering angle of $\pm 2^\circ$ due to mechanical limitations of the pivot arm in moving it in small increments and the obstruction of the angle markings. Errors when calibrating the software may have also contributed to the errors observed, which could have arisen from misplacing the samples in front of the detector or not measuring for long enough to obtain a clear photopeak. The experimental fit is observed to diverge more from the theoretical prediction at higher scattering angles, which may have been mitigated by taking more counts at all angles to reduce the statistical error.

The plot of the differential cross-section in Figure 4 shows a similar trend to the theoretical prediction obtained from the Klein-Nishina formula by following the expected behaviour of the probability of scattering decreasing with the increasing scattering angle. However, the experimental values were consistently lower than the theoretical line, even after scaling the data to the Klein-Nishina prediction at $\theta = 20^\circ$, which suggests presence of systematic errors in the measurements rather than random errors or fluctuations.

The likely sources of error could be partly due to the Cs-137 source, which was obtained in the 1970s and thus may have had a reduced activity due to decay, thus necessitating longer measurement times to obtain more counts to reduce the statistical error, and could have influenced signal-to-noise ratio in the software and the accuracy of the net counts obtained. Additionally, the NaI(Tl) scintillation device efficiency is influenced by age and temperature most notably, and thus the accuracy of the general efficiency formula used may have led to the underestimation of the experimental values. Another source of error could be the estimation of the solid angle of the detector, which generally assumes a point source and flat detector surface, for which the target may have caused angular smearing by measuring over a range of angles rather than a single angle.

Despite these discrepancies, the results obtained for both the kinematics and the differential cross-section show the expected respective trends and

are relatively close to the theoretical predictions, even if not in perfect agreement.

5. Conclusion

The results of this experiment provide evidence for the Compton scattering phenomenon and strong support for the theoretical predictions of the energy of the scattered gamma-ray photon and the differential cross-section as a function of the scattering angle. The linear relationship observed in the plot of $1/E_{\gamma'}$ against $(1 - \cos \theta)$ in Figure 3 is consistent with the expected behaviour from the Compton scattering formula. The small percentage errors obtained for the slope and y-intercept values demonstrate that the experimental setup was capable of accuracy despite mechanical limitations, particularly in the measurement and alignment of the scattering angles.

The plot of the differential cross-section in Figure 4 also shows a similar trend as to expectations with the theoretical prediction from the Klein-Nishina formula, though the underestimation of the experimental values showed the difference between the idealised theoretical model and the realistic experimental setup with its various sources of systematic errors.

Despite the errors present in the measurements, the results obtained provide strong evidence and support for the theory that underlines the Compton scattering phenomenon, and the experiment was successful in demonstrating the expected behaviours and providing understanding on quantum electrodynamics and the interaction of light with matter.

References

- [1] V. Venugopal and P. Bhagdikar, “de broglie wavelength and frequency of scattered electrons in the compton effect,” *Physics Education*, vol. 29, p. 35, 03 2013.
- [2] G. F. Knoll, *Radiation detection and measurement*. John Wiley & Sons, 2010.
- [3] C.-K. Qiao, J.-W. Wei, and L. Chen, “An overview of the compton scattering calculation,” *Crystals*, vol. 11, no. 5, 2021. [Online]. Available: <https://www.mdpi.com/2073-4352/11/5/525>
- [4] University College Dublin, “Interaction of ionizing radiation with matter, part 3: Gamma rays, x-rays and neutrons,” Lecture Notes, PHYC30090, 2026, [Accessed 31-March-2026].
- [5] —, “Compton Scattering Experiment,” Laboratory Manual, n.d., [Accessed 31-March-2026].
- [6] R. Pratt, L. LaJohn, V. Florescu, T. Surić, B. Chatterjee, and S. Roy, “Compton scattering revisited,” *Radiation Physics and Chemistry*, vol. 79, no. 2, pp. 124–131, 2010, special Issue on Future Directions in Atomic and Condensed Matter Research and Applications. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S0969806X09003491>
- [7] G. B. Rybicki and A. P. Lightman, *Radiative processes in astrophysics*. John Wiley & Sons, 2024.

Appendix

Tables

Table 1: Table of the raw data obtained for the experiment.

Angle (°)	NET count	Count Unc.	Energy (keV)	Live Time (s)
20	10852	358	591.67	123.48
25	11112	399	569.62	153.36
30	11156	516	539.86	275.78
35	12553	478	514.52	256.28
40	14329	585	485.86	401.84
45	10583	465	457.75	295.86
50	11493	501	429.49	379.7
55	10800	479	409.15	386.8
60	15131	537	383.57	557.98
65	10310	421	363.01	370.54
70	10420	389	343.42	374.12
75	10540	414	324.33	445.22
80	11269	388	306.35	424.02
85	10068	352	289.84	377.9
90	10104	305	279.4	317.9

compton

April 1, 2026

```
[1]: import numpy as np
import matplotlib.pyplot as plt
import pandas as pd
from scipy.optimize import curve_fit

plt.rcParams['figure.figsize'] = [10, 6]

[2]: compton = pd.read_csv("Compton - material wtv.csv")

scat_angle = compton['angle']
cnt = compton['NET count']
uncnt = compton['cnt uncertainty']
energy = compton['energy (keV)']
time = compton['live time (s)']
```

1 Part 1

```
[3]: angle = np.radians(scat_angle)
angle_err = np.radians(2) # it was impossible to see if i was directly on the
↪ correct angle

inv_energy = 1 / energy * 1e+3 # keV -> MeV
plot_angle = 1 - np.cos(angle)

plot_angle_err = np.sin(angle) * angle_err

def fit(m,x,c):
    return m*x+c

popt, pcov = curve_fit(fit, plot_angle, inv_energy)
m, c = popt
fit_err = np.sqrt(np.diag(pcov))

m_th = 1.956
c_th = 1.51

ang_range = np.linspace(plot_angle.min(), plot_angle.max(), 100)
```

```

fit_th = m_th * ang_range + c_th

slope_err = np.abs((m - m_th) / m_th) * 100
intcpt_err = np.abs((c - c_th) / c_th) * 100

```

```


[4]: plt.errorbar(plot_angle, inv_energy, xerr=plot_angle_err, fmt='o', color='tab:
      ↪orange', ecolord='tab:red', capsize=2, label='Experimental Data')
plt.plot(plot_angle, fit(plot_angle, *popt), '-', color='tab:blue',
      ↪label='Linear Fit')
plt.plot(ang_range, fit_th, '--', color='tab:blue', alpha=0.5,
      ↪label='Theoretical Line')

plt.xlabel(r'1-$\cos(\theta)$ (rad)')
plt.ylabel(r'1/$E_{\gamma}$ (MeV)') # since theoretical used MeV values
plt.minorticks_on()
plt.grid(alpha=0.3)
plt.legend(fontsize=13)

plt.show()

print(f'Slope: {m:.3f} ± {fit_err[0]:.4f}')
print(f'Intercept: {c:.2f} ± {fit_err[1]:.3f}')
print(f'Slope Percentage Error: {slope_err:.2f}%')
print(f'Intercept Percentage Error: {intcpt_err:.2f}%')

```



Slope: 2.030 ± 0.0120
Intercept: 1.58 ± 0.007
Slope Percentage Error: 3.77%
Intercept Percentage Error: 4.68%

2 Part 2

```
[43]: r_det = 2.5 # cm
dist = 26 # cm
M_rod = 79.3 # g
Z_Al = 13
A_Al = 26.9815
Na = 6.02e+23

domega = (np.pi * r_det**2) / dist**2 # solid angle (steradians)
N = (M_rod * Z_Al * Na) / A_Al # no. of electrons in scattering sample

def int_energy(energy):
    energy_mev = energy * 1e-3
    return 0.1522 * (energy_mev**(-1.1325))

eff = int_energy(energy)
cnt_rate = cnt / time
crosssect = cnt_rate / (N * domega * eff)

crosssect_unc = crosssect * (unccnt / cnt)
theta_err = 2

def KN(theta, energy_Cs=661.7e-3): # putting it immediately in MeV
    alpha = energy_Cs / 0.511
    thetar = np.radians(theta)
    cos = np.cos(thetar)

    p1 = (1 + cos**2) / (1 + alpha * (1 - cos)**2)
    p2 = 1 + ((alpha**2 * (1 - cos)**2) / ((1 + cos**2) * (1 + alpha * (1 -
↵cos)))))

    r0 = 2.82e-13 # cm
    return (r0**2) / 2 * p1 * p2

range_theta = np.linspace(0, 100, 100)
th_val = [KN(i) for i in range_theta]

scale = KN(20) / crosssect[0]
norm_data = crosssect * scale # normalised (scaled to KN)
norm_err = crosssect_unc * scale
```

```
[44]: plt.errorbar(scatter_angle, norm_data, yerr=norm_err, xerr=theta_err, fmt='o',
    color='tab:green', ecolor='tab:red', capsize=2, label='Scaled Experimental
    Data')
plt.plot(range_theta, th_val, color='tab:blue', label='Klein-Nishina Theory')

plt.xlabel(r'Scattering Angle (deg)')
plt.ylabel(r' $d\sigma/d\Omega$  (cm2/sr)')
plt.minorticks_on()
plt.grid(alpha=0.3)
plt.legend(fontsize=13)

plt.show()
```

