

# Assignment 3: Data Exploration

Isaac Benaka

## OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Exploration.

## Directions

1. Rename this file `<FirstLast>_A03_DataExploration.Rmd` (replacing `<FirstLast>` with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.
6. After Knitting, submit the completed exercise (PDF file) to the dropbox in Sakai.

The completed exercise is due on Sept 30th.

## Set up your R session

1. Check your working directory, load necessary packages (tidyverse), and upload two datasets: the ECOTOX neonicotinoid dataset (ECOTOX\_Neonicotinoids\_Insects\_raw.csv) and the Niwot Ridge NEON dataset for litter and woody debris (NEON\_NIWO\_Litter\_massdata\_2018-08\_raw.csv). Name these datasets “Neonics” and “Litter”, respectively. Be sure to include the subcommand to read strings in as factors.

```
library(tidyverse)
library(lubridate)
setwd("~/Desktop/Fall 2022/872 - Data Analytics/EDA-Fall2022")
Neonics <- read.csv("../Data/Raw/ECOTOX_Neonicotinoids_Insects_raw.csv", stringsAsFactors = TRUE)
Litter <- read.csv("../Data/Raw/NEON_NIWO_Litter_massdata_2018-08_raw.csv", stringsAsFactors = TRUE)
```

## Learn about your system

2. The neonicotinoid dataset was collected from the Environmental Protection Agency’s ECOTOX Knowledgebase, a database for ecotoxicology research. Neonicotinoids are a class of insecticides used widely in agriculture. The dataset that has been pulled includes all studies published on insects. Why might we be interested in the ecotoxicology of neonicotinoids on insects? Feel free to do a brief internet search if you feel you need more background information.

Answer: We are interested in neonicotinoid impacts on insects because the pesticides may be impacting other insect species than the ones they are targeting on crops. Neonicotinoids may also trickle into the wider food web by predators eating the insects they act on.

3. The Niwot Ridge litter and woody debris dataset was collected from the National Ecological Observatory Network, which collectively includes 81 aquatic and terrestrial sites across 20 ecoclimatic domains. 32 of these sites sample forest litter and woody debris, and we will focus on the Niwot Ridge long-term ecological research (LTER) station in Colorado. Why might we be interested in studying litter and woody debris that falls to the ground in forests? Feel free to do a brief internet search if you feel you need more background information.

Answer: We might be interested in studying woody debris data because forest litter that falls onto the ground contributes to decomposition and nutrient cycling.

4. How is litter and woody debris sampled as part of the NEON network? Read the `NEON_Litterfall_UserGuide.pdf` document to learn more. List three pieces of salient information about the sampling methods here:

Answer: 1. 0.5 m<sup>2</sup> mesh traps, elevated 80 cm above the ground, were used to collect litter and woody debris. 2. In forested tower airsheds, sampling took place over 20 40m x 40m plots. 3. Ground traps were sampled once per year.

## Obtain basic summaries of your data (Neonics)

5. What are the dimensions of the dataset?

```
dim(Neonics)
```

```
## [1] 4623 30
```

6. Using the `summary` function on the “Effect” column, determine the most common effects that are studied. Why might these effects specifically be of interest?

```
summary(Neonics$Effect)
```

```
##      Accumulation      Avoidance      Behavior      Biochemistry
##           12           102           360           11
##      Cell(s)      Development      Enzyme(s)      Feeding behavior
##           9           136           62           255
##      Genetics      Growth      Histology      Hormone(s)
##          82           38           5           1
##      Immunological      Intoxication      Morphology      Mortality
##          16           12           22           1493
##      Physiology      Population      Reproduction
##           7           1803           197
```

Answer: Based on the summary, mortality and population were the most common effects that were studied. These effects were likely studied the most because they are the most important factors when it comes to population modeling.

7. Using the `summary` function, determine the six most commonly studied species in the dataset (common name). What do these species have in common, and why might they be of interest over other insects? Feel free to do a brief internet search for more information if needed.

```
summary(Neonics$Species.Common.Name)
```

##	Honey Bee	Parasitic Wasp
##	667	285
##	Buff Tailed Bumblebee	Carniolan Honey Bee
##	183	152
##	Bumble Bee	Italian Honeybee
##	140	113
##	Japanese Beetle	Asian Lady Beetle
##	94	76
##	Euonymus Scale	Wireworm
##	75	69
##	European Dark Bee	Minute Pirate Bug
##	66	62
##	Asian Citrus Psyllid	Parastic Wasp
##	60	58
##	Colorado Potato Beetle	Parasitoid Wasp
##	57	51
##	Erythrina Gall Wasp	Beetle Order
##	49	47
##	Snout Beetle Family, Weevil	Sevenspotted Lady Beetle
##	47	46
##	True Bug Order	Buff-tailed Bumblebee
##	45	39
##	Aphid Family	Cabbage Looper
##	38	38
##	Sweetpotato Whitefly	Braconid Wasp
##	37	33
##	Cotton Aphid	Predatory Mite
##	33	33
##	Ladybird Beetle Family	Parasitoid
##	30	30
##	Scarab Beetle	Spring Tiphia
##	29	29
##	Thrip Order	Ground Beetle Family
##	29	27
##	Rove Beetle Family	Tobacco Aphid
##	27	27
##	Chalcid Wasp	Convergent Lady Beetle
##	25	25
##	Stingless Bee	Spider/Mite Class
##	25	24
##	Tobacco Flea Beetle	Citrus Leafminer
##	24	23
##	Ladybird Beetle	Mason Bee
##	23	22
##	Mosquito	Argentine Ant
##	22	21
##	Beetle	Flatheaded Appletree Borer
##	21	20
##	Horned Oak Gall Wasp	Leaf Beetle Family
##	20	20
##	Potato Leafhopper	Tooth-necked Fungus Beetle

##	20	20
##	Codling Moth	Black-spotted Lady Beetle
##	19	18
##	Calico Scale	Fairyfly Parasitoid
##	18	18
##	Lady Beetle	Minute Parasitic Wasps
##	18	18
##	Mirid Bug	Mulberry Pyralid
##	18	18
##	Silkworm	Vedalia Beetle
##	18	18
##	Araneoid Spider Order	Bee Order
##	17	17
##	Egg Parasitoid	Insect Class
##	17	17
##	Moth And Butterfly Order	Oystershell Scale Parasitoid
##	17	17
##	Hemlock Woolly Adelgid Lady Beetle	Hemlock Woolly Adelgid
##	16	16
##	Mite	Onion Thrip
##	16	16
##	Western Flower Thrips	Corn Earworm
##	15	14
##	Green Peach Aphid	House Fly
##	14	14
##	Ox Beetle	Red Scale Parasite
##	14	14
##	Spined Soldier Bug	Armoured Scale Family
##	14	13
##	Diamondback Moth	Eulophid Wasp
##	13	13
##	Monarch Butterfly	Predatory Bug
##	13	13
##	Yellow Fever Mosquito	Braconid Parasitoid
##	13	12
##	Common Thrip	Eastern Subterranean Termite
##	12	12
##	Jassid	Mite Order
##	12	12
##	Pea Aphid	Pond Wolf Spider
##	12	12
##	Spotless Ladybird Beetle	Glasshouse Potato Wasp
##	11	10
##	Lacewing	Southern House Mosquito
##	10	10
##	Two Spotted Lady Beetle	Ant Family
##	10	9
##	Apple Maggot	(Other)
##	9	670

Answer: The six most commonly studied species are Honey bee, Parasitic wasp, Buff tailed bumblebee, Carniolan honey bee, Bumble bee, and Italian honey bee. All these species are key pollinators, meaning their ecosystem impacts and benefits are likely superior over other insect species.

8. Concentrations are always a numeric value. What is the class of `Conc.1..Author.` in the dataset, and why is it not numeric?

```
class(Neonics$Conc.1..Author.)
```

```
## [1] "factor"
```


Answer: The class of `Conc.1..Author.` is a factor. This class is a factor instead of numeric because these are categorical values assigned to each author.

## Explore your data graphically (Neonics)

9. Using `geom_freqpoly`, generate a plot of the number of studies conducted by publication year.

```
ggplot(Neonics) +  
  geom_freqpoly(aes(x = Publication.Year))
```

```
## 'stat_bin()' using 'bins = 30'. Pick better value with 'binwidth'.
```




10. Reproduce the same graph but now add a color aesthetic so that different `Test.Location` are displayed as different colors.

```
class(Neonics$Test.Location)
```

```
## [1] "factor"
```

```
ggplot(Neonics) +  
  geom_freqpoly(aes(color = Test.Location, x = Publication.Year)) +  
  theme(legend.position = "top")
```

```
## 'stat_bin()' using 'bins = 30'. Pick better value with 'binwidth'.
```




Interpret this graph. What are the most common test locations, and do they differ over time?

Answer: The most common test locations are “Lab” and “Field natural.” Starting in 2010, “Lab” test locations increased as “Field natural” locations decreased.

11. Create a bar graph of Endpoint counts. What are the two most common end points, and how are they defined? Consult the ECOTOX\_CodeAppendix for more information.

```
ggplot(Neonics, aes(x = Endpoint)) +  
  geom_bar()
```



Answer: LOEL and NOEL are the two most common end points. LOEL stands for lowest-observable-effect-level and NOEL stands for no-observable-effect-level.

## Explore your data (Litter)

12. Determine the class of `collectDate`. Is it a date? If not, change to a date and confirm the new class of the variable. Using the `unique` function, determine which dates litter was sampled in August 2018.

```
class(Litter$collectDate) #This class was originally a factor.
```

```
## [1] "factor"
```

```
Litter$collectDate <- ymd(Litter$collectDate)
class(Litter$collectDate) #Recheck the class of collectDate to view new class.
```

```
## [1] "Date"
```

```
unique(Litter$collectDate) #Litter was sampled on the 2nd and the 30th in August 2018.
```

```
## [1] "2018-08-02" "2018-08-30"
```

13. Using the `unique` function, determine how many plots were sampled at Niwot Ridge. How is the information obtained from `unique` different from that obtained from `summary`?

```
unique(Litter$plotID)
```

```
## [1] NIWO_061 NIWO_064 NIWO_067 NIWO_040 NIWO_041 NIWO_063 NIWO_047 NIWO_051  
## [9] NIWO_058 NIWO_046 NIWO_062 NIWO_057  
## 12 Levels: NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 ... NIWO_067
```


```
summary(Litter$plotID)
```

```
## NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 NIWO_058 NIWO_061  
##      20      19      18      15      14       8      16      17  
## NIWO_062 NIWO_063 NIWO_064 NIWO_067  
##      14      14      16      17
```

Answer: the 'unique' function only returns the names of the different plots, whereas the 'summary' function returns how many samples correspond with each plot.

14. Create a bar graph of functionalGroup counts. This shows you what type of litter is collected at the Niwot Ridge sites. Notice that litter types are fairly equally distributed across the Niwot Ridge sites.


```
ggplot(Litter, aes(x = functionalGroup)) +  
  geom_bar()
```




15. Using `geom_boxplot` and `geom_violin`, create a boxplot and a violin plot of `dryMass` by `functionalGroup`.



```
#Boxplot
ggplot(Litter) +
  geom_boxplot(aes(x = functionalGroup, y = dryMass))
```



```
#Violin plot
ggplot(Litter) +
  geom_violin(aes(x = functionalGroup, y = dryMass))
```



Why is the boxplot a more effective visualization option than the violin plot in this case?

Answer:

What type(s) of litter tend to have the highest biomass at these sites?

Answer: Needles tended to compose the largest amount of biomass at the sites.