

Homotopy theory

Ikhan Choi

September 19, 2022

Contents

I	Fiber bundles	2
1	Principal bundles	3
1.1	Category of bundles	3
1.2	Classifying spaces	3
1.3	Čech cohomology	3
2	Vector bundles	5
2.1	5
3	Characteristic classes	6
II	K-theory	7
III		8
IV		9

Part I

Fiber bundles

Chapter 1

Principal bundles

1.1 Category of bundles

1.1 (Pullback and restricted bundles).

1.2 (Product of bundles). the *Fiber product* or the *Whitney sum*

1.2 Classifying spaces

1.3 Čech cohomology

1.3. Let $\{U_\alpha\}_\alpha$ be an open cover of a topological space X . We say a presheaf \mathcal{F} on X is sheaf if the sequence

$$\mathcal{F}(U) \longrightarrow \prod_\alpha \mathcal{F}(U_\alpha) \xrightarrow[\text{res}_\beta]{\text{res}_\alpha} \prod_{\alpha, \beta} \mathcal{F}(U_\alpha \cap U_\beta)$$

is an equalizer.

1.4. Let \mathcal{F} be a preseah of groups and let $\mathcal{U} = \{U_\alpha\}_\alpha$ be an ordered open cover of a topological space X .

$$\prod_\alpha \mathcal{F}(U_\alpha) \xrightarrow[\text{res}_\beta]{\text{res}_\alpha} \prod_{\alpha < \beta} \mathcal{F}(U_\alpha \cap U_\beta) \xrightarrow[\text{res}_\gamma]{\text{res}_\alpha} \prod_{\alpha < \beta < \gamma} \mathcal{F}(U_\alpha \cap U_\beta \cap U_\gamma)$$

$$C^0(\mathcal{U}, \mathcal{F}) \xrightarrow{\delta} C^1(\mathcal{U}, \mathcal{F}) \xrightarrow{\delta} C^2(\mathcal{U}, \mathcal{F}) \xrightarrow{\delta} \dots$$

$$f = \{f_\alpha : U_\alpha \rightarrow \text{GL}(k, \mathbb{R})\}_\alpha \in \prod_\alpha \mathcal{F}(U_\alpha) = C^0(\mathcal{U}, \mathcal{F})$$

1.5. Let G be a sheaf of groups. Then, we have a natural one-to-one correspondence

$$\left\{ \begin{array}{l} \text{isomorphism classes of} \\ \text{principal } G\text{-bundles} \end{array} \right\} \xrightarrow{\sim} H^1(B, G).$$

Proof. (Injectivity) We show the correspondence is left invertible. Let $p : E \rightarrow B$ be a principal G -bundle. Define $p' : E' \rightarrow B$ by

$$E' := \left(\bigsqcup_\alpha \{\alpha\} \times U_\alpha \times \mathbb{R}^k \right) / \sim,$$

where the equivalence relation \sim denotes $(\alpha, b, g_{\alpha\beta}(v)) \sim (\beta, b, v)$ with $b \in U_\alpha \cap U_\beta$ for some α, β , and define $p' : E' \rightarrow B$, which is well-defined bundle.

We first check that p' is a principal G -bundle.

(Surjectivity)

□

Chapter 2

Vector bundles

2.1

2.1 (Vector space structure on total spaces).

2.2 (Vector bundle maps). Let $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ be vector bundles.

- (a) A vector bundle map u over B is vector bundle isomorphism if and only if it is a fiberwise linear isomorphism.

2.3 (Tautological bundles).

2.4 (Homotopy properties). Let $p : E \rightarrow B$ be a vector bundle

If $p_1 : E_1 \rightarrow B \times [0, \frac{1}{2}]$ and $p_2 : E_2 \rightarrow B \times [\frac{1}{2}, 1]$ are trivial, then

- (a)

2.5 (Principal bundle over the general linear group).

Chapter 3

Characteristic classes

Part II

K-theory

Part III

Part IV