

Probability Theory

Ikhan Choi

June 18, 2022

Contents

I	Probability distributions	2
1	Random variables	3
1.1	Sample spaces and distributions	3
1.2	Discrete probability distributions	3
1.3	Continuous probability distributions	3
1.4	Independence	3
2	Conditional probability	4
3	Convergence of probability measures	5
3.1	Weak convergence in \mathbb{R}	5
3.2	Weak topology in the space of probability measures	5
3.3	Characteristic functions	7
3.4	Moments	8
II	Discrete stochastic process	9
4	Limit theorems	10
4.1	Laws of large numbers	10
4.2	Renewal theory	11
4.3	Central limit theorems	11
5	Martingales	13
5.1	Submartingales	13
5.2	Martingale convergence theorem	13
5.3	Convergence in L^p and uniform integrability	13
5.4	Optional stopping theorem	13
6	Markov chains	14
III	Continuous stochastic processes	15
7	Brownian motion	16
7.1	Kolomogorov extension	16
IV	Stochastic calculus	17

Part I

Probability distributions

Chapter 1

Random variables

1.1 Sample spaces and distributions

sample space of an "experiment" random variables distributions expectation, moments, inequalities
equally likely outcomes coin toss dice roll ball drawing number permutation life time of a light bulb
joint distribution transformation of distributions distribution computations

1.2 Discrete probability distributions

1.3 Continuous probability distributions

1.4 Independence

1.1 (Dynkin's π - λ lemma). Let \mathcal{P} be a π -system and \mathcal{L} a λ -system respectively. Denote by $\ell(\mathcal{P})$ the smallest λ -system containing \mathcal{P} .

- (a) If $A \in \ell(\mathcal{P})$, then $\mathcal{G}_A := \{B : A \cap B \in \ell(\mathcal{P})\}$ is a λ -system.
- (b) $\ell(\mathcal{P})$ is a π -system.
- (c) If a λ -system is a π -system, then it is a σ -algebra.
- (d) If $\mathcal{P} \subset \mathcal{L}$, then $\sigma(\mathcal{P}) \subset \mathcal{L}$.

1.2 (Monotone class lemma).

Chapter 2

Conditional probability

2.1 (Monty Hall problem). Suppose you're on a game show, and you're given the choice of three doors A , B , and C . Behind one door is a car; behind the others, goats. You pick a door, say A , and the host, who knows what's behind the doors, opens another door, say B , which has a goat. He then says to you, "Do you want to pick door C ?" Is it to your advantage to switch your choice?

Proof. Let A , B , and C be the events that a car is behind the doors A , B , and C , respectively. Let X be the event that the challenger picked A , and Y the event that the game host opened B . Note $\{A, B, C\}$ is a partition of the sample space Ω , and X is independent to A , B , and C . Then, $P(A) = P(B) = P(C) = P(X) = 1/3$, and

$$P(Y|X, A) = \frac{1}{2}, \quad P(Y|X, B) = 0, \quad P(Y|X, C) = 1.$$

Therefore,

$$\begin{aligned} P(C|X, Y) &= \frac{P(X \cap Y \cap C)}{P(X \cap Y)} \\ &= \frac{P(Y|X, C)P(X \cap C)}{P(Y|X, A)P(X \cap A) + P(Y|X, B)P(X \cap B) + P(Y|X, C)P(X \cap C)} \\ &= \frac{1 \cdot \frac{1}{9}}{\frac{1}{2} \cdot \frac{1}{9} + 0 \cdot \frac{1}{9} + 1 \cdot \frac{1}{9}} = \frac{2}{3}. \end{aligned}$$

Similarly, $P(A|X, Y) = \frac{1}{3}$ and $P(B|X, Y) = 0$. □

Chapter 3

Convergence of probability measures

3.1 Weak convergence in \mathbb{R}

3.1 (Portemanteau theorem). Let F_n and F be distribution functions $\mathbb{R} \rightarrow [0, 1]$. We will define the *weak convergence* as follows: F_n converges weakly to F if $F_n(x) \rightarrow F(x)$ for every continuity point x of $F(x)$.

(a) $F_n(x) \rightarrow F(x)$ for all continuity points x of F .

3.2 (Skorokhod representation theorem).

3.3 (Continuous mapping theorem).

3.4 (Slutsky's theorem).

3.5 (Helly's selection theorem). (a) Monotonically increasing functions $F_n : \mathbb{R} \rightarrow [0, 1]$ has a point-wise convergent subsequence.

(b) If $(F_n)_n$ is tight, then

3.6 (Properties of probability Borel measures). Let S be a topological space.

(a) Every single probability Borel measure is regular if S is perfectly normal.

(b) Every single probability Borel measure is tight if S is Polish.

3.2 Weak topology in the space of probability measures

3.7 (Local limit theorems). Suppose f_n and f are density functions.

(a) If $f_n \rightarrow f$ a.s., then $f_n \rightarrow f$ in L^1 .

(Scheffé's theorem)

(b) $f_n \rightarrow f$ in L^1 if and only if in total variation.

(c) If $f_n \rightarrow f$ in total variation, then $f_n \rightarrow f$ weakly.

3.8 (Portmanteau theorem). Let S be a normal space and, μ_α be a net in $\text{Prob}(S)$. We define the *weak convergence* as follows: μ_α converges weakly to μ if

$$\int f d\mu_\alpha \rightarrow \int f d\mu$$

for every $f \in C_b(S)$. The following statements are all equivalent.

- (a) $\mu_\alpha \Rightarrow \mu$
- (b) $\mu_\alpha(g) \rightarrow \mu(g)$ for every uniformly continuous $g \in C_b(S)$.
- (c) $\limsup_\alpha \mu_\alpha(F) \leq \mu(F)$ for every closed F .
- (d) $\liminf_\alpha \mu_\alpha(U) \geq \mu(U)$ for every open U .
- (e) $\lim_\alpha \mu_\alpha(A) = \mu(A)$ for every Borel A such that $\mu(\partial A) = 0$.

Proof. (a) \Rightarrow (b) Clear.

(b) \Rightarrow (c) Let U be an open set such that $F \subset U$. There is uniformly continuous $g \in C_b(S)$ such that $\mathbf{1}_F \leq g \leq \mathbf{1}_U$. Therefore,

$$\limsup_\alpha \mu_\alpha(F) \leq \limsup_\alpha \mu_\alpha(g) = \mu(g) \leq \mu(U).$$

By the outer regularity of μ , we obtain $\limsup_\alpha \mu_\alpha(F) \leq \mu(F)$.

(c) \Leftrightarrow (d) Clear.

(c)+(d) \Rightarrow (e) It easily follows from

$$\limsup_\alpha \mu_\alpha(\bar{A}) \leq \mu(\bar{A}) = \mu(A) = \mu(A^\circ) \leq \liminf_\alpha \mu_\alpha(A^\circ).$$

(e) \Rightarrow (a) Let $g \in C_b(S)$ and $\varepsilon > 0$. Since the pushforward measure $g_*\mu$ has at most countably many mass points, there is a partition $(t_i)_{i=0}^n$ of an interval containing $[-\|g\|, \|g\|]$ such that $|t_{i+1} - t_i| < \varepsilon$ and $\mu(\{x : g(x) = t_i\}) = 0$ for each i . Let $(A_i)_{i=0}^{n-1}$ be a Borel decomposition of S given by $A_i := g^{-1}([t_i, t_{i+1}))$, and define $f_\varepsilon := \sum_{i=0}^{n-1} t_i \mathbf{1}_{A_i}$ so that we have $\sup_{x \in S} |g_\varepsilon(x) - g(x)| \leq \varepsilon$. From

$$\begin{aligned} |\mu_\alpha(g) - \mu(g)| &\leq |\mu_\alpha(g - g_\varepsilon)| + |\mu_\alpha(g_\varepsilon) - \mu(g_\varepsilon)| + |\mu(g_\varepsilon) - \mu(g)| \\ &\leq \varepsilon + \sum_{i=0}^{n-1} |t_i| |\mu_\alpha(A_i) - \mu(A_i)| + \varepsilon, \end{aligned}$$

we get

$$\limsup_\alpha |\mu_\alpha(g) - \mu(g)| < 2\varepsilon.$$

Since ε is arbitrary, we are done. □

3.9 (Dirac measures). Let S be a normal space.

- (a) $S \rightarrow \text{Prob}(S)$ is an embedding.
- (b) $S \subset \text{Prob}(S)$ is sequentially closed.

Proof. (a) It uses Urysohn.

(b) It uses (b) \Rightarrow (c) of Portmanteau. □

3.10 (Lévy-Prokhorov metric). Let S be a metric space, and $\text{Prob}(S)$ be the set of probability (regular) Borel measures on S . Define $\pi : \text{Prob}(S) \times \text{Prob}(S) \rightarrow [0, \infty)$ such that

$$\pi(\mu, \nu) := \inf\{\alpha > 0 : \mu(A) \leq \nu(A^\alpha) + \alpha, \nu(A) \leq \mu(A^\alpha) + \alpha, \forall A \in \mathcal{B}(S)\},$$

where A^α is the α -neighborhood of A .

- (a) π is a metric.
- (b) $\mu_n \rightarrow \mu$ in π implies $\mu_n \Rightarrow \mu$.
- (c) $\mu_\alpha \Rightarrow \mu$ implies $\mu_\alpha \rightarrow \mu$ in π , if S is separable.
- (d) (S, d) is separable if and only if $(\text{Prob}(S), \pi)$ is separable.

- (e) (S, d) is compact if and only if $(\text{Prob}(S), \pi)$ is compact
- (f) (S, d) is complete if and only if $(\text{Prob}(S), \pi)$ is complete.

Proof. (c) □

3.11 (Prokhorov's theorem). Let S be a normal space. Let $\text{Prob}(S)$ be the space of probability measures on S endowed with the topology of weak convergence.

- (a) The relative compactness implies the tightness if S is Polish.
- (b) The tightness implies the relative compactness.

Proof. (a) Fix a complete metric d on S . For each entourage E , we want to find a finite union of closed E -neighborhood that is arbitrarily full. And we require the countable fundamental system of uniformity.

(b) □

3.3 Characteristic functions

3.12 (Characteristic functions). Let μ be a probability measure on \mathbb{R} . Then, the *characteristic function* of μ is defined by

$$\varphi(t) := Ee^{itX} = \int e^{itx} d\mu(x).$$

Note that $\varphi(t) = \hat{\mu}(-t)$ where $\hat{\mu}$ is the Fourier transform of $\mu \in \mathcal{S}'(\mathbb{R})$.

- (a) $\varphi \in C_b(\mathbb{R})$.

3.13 (Inversion formula). Let μ be a probability measure on \mathbb{R} and φ its characteristic function.

- (a) For $a < b$, we have

$$\mu((a, b)) + \frac{1}{2}\mu(\{a, b\}) = \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-ita} - e^{-itb}}{it} \varphi(t) dt.$$

- (b) For $a \in \mathbb{R}$, we have

$$\mu(\{a\}) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T e^{-ita} \varphi(t) dt$$

- (c) If $\varphi \in L^1(\mathbb{R})$, then μ has density

$$f(x) = \frac{1}{2\pi} \int e^{-itx} \varphi(t) dt$$

in $C_0(\mathbb{R}) \cap L^1(\mathbb{R})$.

3.14 (Lévy's continuity theorem). The continuity theorem provides with a tool to verify the weak convergence in terms of characteristic functions. Let μ_n and μ be probability distributions on \mathbb{R} with characteristic functions φ_n and φ .

- (a) If $\mu_n \rightarrow \mu$ weakly, then $\varphi_n \rightarrow \varphi$ pointwise.
- (b) If $\varphi_n \rightarrow \varphi$ pointwise and φ is continuous at zero, then $(\mu_n)_n$ is tight and $\mu_n \rightarrow \mu$ weakly.

Proof. (a) For each t ,

$$\varphi_n(t) = \int e^{itx} d\mu_n(x) \rightarrow \int e^{itx} d\mu(x) = \varphi(t)$$

because $e^{itx} \in C_b(\mathbb{R})$.

(b) □

3.15 (Criteria for characteristic functions). Bochner's theorem and Polya's criterion

There are two ways to represent a measure: A measure μ is absolutely continuous iff its distribution F is absolutely continuous iff its density f is integrable. So, the fourier transform of an absolutely continuous measure is just the fourier transform of L^1 functions.

3.4 Moments

moment problem

moment generating function defined on $|t| < \delta$

Exercises

3.16. Let φ_n be characteristic functions of probability measures μ_n on \mathbb{R} . If there is a continuous function φ such that $\varphi_n = \varphi$ on $n^{-1}\mathbb{Z}$, then μ_n converges weakly.

3.17 (Convergence determining class).

3.18 (Vague convergence). Let S be a locally compact Hausdorff space.

- (a) $\mu_\alpha \rightarrow \mu$ vaguely if and only if $\int g d\mu_\alpha \rightarrow \int g d\mu$ for all $g \in C_c(S)$.
- (b) $\mu_\alpha \rightarrow \mu$ weakly if and only if vaguely.
- (c) $\delta_n \rightarrow 0$ vaguely but not weakly. (escaping to infinity)

Proof.

□

Part II

Discrete stochastic process

Chapter 4

Limit theorems

4.1 Laws of large numbers

Our purpose is to find appropriate a_n and slowly growing b_n such that $(S_n - a_n)/b_n \rightarrow 0$ in probability or almost surely.

4.1 (Truncation method). Let $X_{n,i} : \Omega \rightarrow \mathbb{R}$ be uncorrelated random variables (with respect to i for each n) and $S_n := X_{n,1} + \cdots + X_{n,n}$. For a positive sequence $(c_n)_{n=1}^\infty$, let $Y_{n,i} := X_{n,i} \mathbf{1}_{|X_{n,i}| \leq c_n}$ be truncated random variables and $T_n := Y_{n,1} + \cdots + Y_{n,n}$. Suppose that the truncation level c_n satisfies the approximation condition

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n P(|X_{n,i}| > c_n) = 0.$$

- (a) If $(T_n - ET_n)/b_n \rightarrow 0$ in probability, then $(S_n - ET_n)/b_n \rightarrow 0$ in probability.
- (b) If $(T_n - ET_n)/b_n \rightarrow Z$ in distribution, then $(S_n - ET_n)/b_n \rightarrow Z$ in distribution.

Proof. (a) Write

$$P\left(\left|\frac{S_n - ET_n}{b_n}\right| > \varepsilon\right) \leq P(S_n \neq T_n) + P\left(\left|\frac{T_n - ET_n}{b_n}\right| > \varepsilon\right) \rightarrow 0$$

since

$$P(S_n \neq T_n) \leq \sum_{i=1}^n P(X_{n,i} \neq Y_{n,i}) = \sum_{i=1}^n P(|X_{n,i}| > c_n) \rightarrow 0$$

as $n \rightarrow \infty$.

(b) By the Slutsky theorem. □

4.2 (Weak laws of large numbers). Let $X_{n,i} : \Omega \rightarrow \mathbb{R}$ be uncorrelated random variables and $S_n := X_{n,1} + \cdots + X_{n,n}$. For a positive sequence $(c_n)_{n=1}^\infty$, let $Y_{n,i} := X_{n,i} \mathbf{1}_{|X_{n,i}| \leq c_n}$ be truncated random variables and $T_n := Y_{n,1} + \cdots + Y_{n,n}$.

(a) If

$$b_n^2 \gg \sum_{i=1}^n E|Y_{n,i} - EY_{n,i}|^2,$$

then $(T_n - ET_n)/b_n \rightarrow 0$ in probability.

(b) Take slow c_n as possible such that

$$1 \gg \sum_{i=1}^n P(|X_{n,i}| > c_n).$$

Take slow b_n as possible such that

$$b_n^2 \gg \sum_{i=1}^n E|Y_{n,i}|^2.$$

then $(S_n - ET_n)/b_n \rightarrow 0$.

(c) If

$$\lim_{x \rightarrow \infty} \sup_i xP(|X_i| > x) = 0,$$

then $(S_n - ET_n)/n \rightarrow 0$ in probability. This is called the *Kolmogorov-Feller condition*.

Proof. (a) Since X_n are uncorrelated, we have for any $\varepsilon > 0$ that

$$P\left(\left|\frac{S_n - ES_n}{c_n}\right| > \varepsilon\right) \leq \frac{1}{\varepsilon^2 c_n^2} VS_n \rightarrow 0$$

as $n \rightarrow \infty$.

(c) Write $g(x) := \sup_i xP(|X_i| > x)$. Then, the truncation condition for $b_n = n$ is satisfied as

$$\sum_{i=1}^n P(|X_i| > n) \leq \sum_{i=1}^n \frac{1}{n} g(n) = g(n) \rightarrow 0$$

as $n \rightarrow \infty$.

On the other hand,

$$\begin{aligned} \frac{1}{n^2} \sum_{i=1}^n E|Y_i|^2 &= \frac{1}{n^2} \sum_{i=1}^n \int_0^\infty 2xP(|Y_i| > x) dx = \frac{1}{n^2} \sum_{i=1}^n \int_0^n 2xP(|X_i| > x) dx \\ &\leq \frac{2}{n} \int_0^n g(x) dx = 2 \int_0^1 g(nx) dx. \end{aligned}$$

Since $g(x) \leq x$ and $g(x) \rightarrow 0$ as $x \rightarrow \infty$, g is bounded so that the bounded convergence theorem implies $\int_0^1 g(nx) dx \rightarrow 0$ as $n \rightarrow \infty$.

Therefore, $(T_n - ET_n)/n \rightarrow 0$ in probability. By the truncation □

4.3 (Borel-Cantelli lemmas).

4.4 (Strong laws of large numbers). Proof by Etemadi

Random series

4.2 Renewal theory

4.3 Central limit theorems

4.5 (Lyapunov central limit theorem). Let $X_n : \Omega \rightarrow \mathbb{R}$ be independent random variables with $EX_i = \mu_i$ and $VS_i = \sigma_i^2$. If there is $\delta > 0$ such that the *Lyapunov condition*

$$\lim_{n \rightarrow \infty} \frac{1}{s_n^{2+\delta}} \sum_{i=1}^n E|X_i - \mu_i|^{2+\delta} = 0$$

is satisfied, then

$$\frac{S_n - ES_n}{s_n} \rightarrow N(0, 1)$$

weakly, where $S_n := \sum_{i=1}^n X_i$ and $s_n^2 := VS_n$.

4.6 (Lindeberg-Feller central limit theorem). Let $X_{i,n} : \Omega \rightarrow \mathbb{R}$ be independent random variables and $S_n := X_1 + \dots + X_n$.

(a) If

$$s_n^2 \sim \sum_{i=1}^n E|X_i - EX_i|^2,$$

and if for every $\varepsilon > 0$ we have

$$s_n^2 \gg \sum_{i=1}^n E|X_i - EX_i|^2 \mathbf{1}_{|X_i - EX_i| > \varepsilon s_n},$$

then $(S_n - ES_n)/s_n \rightarrow N(0, 1)$ in distribution. This is called the *Lindeberg condition*.

(b)

Berry-Esseen inequality

Exercises

4.7 (Bernstein polynomial). Let $X_n \sim \text{Bern}(x)$ be i.i.d. random variables. Since $S_n \sim \text{Binom}(n, x)$, $E(S_n/n) = x$, $V(S_n/n) = x(1-x)/n$. The L^2 law of large numbers implies $E(|S_n/n - x|^2) \rightarrow 0$. Define $f_n(x) := E(f(S_n/n))$. Then, by the uniform continuity $|x - y| < \delta$ implies $|f(x) - f(y)| < \varepsilon$,

$$|f_n(x) - f(x)| \leq E(|f(S_n/n) - f(x)|) \leq \varepsilon + 2\|f\|P(|S_n/n - x| \geq \delta) \rightarrow \varepsilon.$$

4.8 (High-dimensional cube is almost a sphere). Let $X_n \sim \text{Unif}(-1, 1)$ be i.i.d. random variables and $Y_n := X_n^2$. Then, $E(Y_n) = \frac{1}{3}$ and $V(Y_n) \leq 1$.

4.9 (Coupon collector's problem). $T_n := \inf\{t : |\{X_i\}_i| = n\}$ Since $X_{n,k} \sim \text{Geo}(1 - \frac{k-1}{n})$, $E(X_{n,k}) = (1 - \frac{k-1}{n})^{-1}$, $V(X_{n,k}) \leq (1 - \frac{k-1}{n})^{-2}$. $E(T_n) \sim n \log n$

4.10 (An occupancy problem).

4.11 (The St. Petersburg paradox).

4.12. Find the probability that arbitrarily chosen positive integers are coprime.

Poisson convergence, law of rare events, or weak law of small numbers (a single sample makes a significant attribution)

Chapter 5

Martingales

5.1 Submartingales

5.2 Martingale convergence theorem

5.1 (Doob's upcrossing inequality). (a)

5.2 (Martingale convergence theorems). (a)

5.3. (a)

5.3 Convergence in L^p and uniform integrability

5.4 Optional stopping theorem

Chapter 6

Markov chains

Part III

Continuous stochastic processes

Chapter 7

Brownian motion

7.1 Kolmogorov extension

7.1 (Kolmogorov extension theorem). A *rectangle* is a finite product $\prod_{i=1}^n A_i \subset \mathbb{R}^n$ of measurable $A_i \subset \mathbb{R}$, and *cylinder* is a product $A^* \times \mathbb{R}^{\mathbb{N}}$ where A^* is a rectangle. Let \mathcal{A} be the semi-algebra containing \emptyset and all cylinders in $\mathbb{R}^{\mathbb{N}}$. Let $(\mu_n)_n$ be a sequence of probability measures on \mathbb{R}^n that satisfies *consistency condition*

$$\mu_{n+1}(A^* \times \mathbb{R}) = \mu_n(A^*)$$

for any rectangles $A^* \subset \mathbb{R}^n$, and define a set function $\mu_0 : \mathcal{A} \rightarrow [0, \infty]$ by $\mu_0(A) = \mu_n(A^*)$ and $\mu_0(\emptyset) = 0$.

- (a) μ_0 is well-defined.
- (b) μ_0 is finitely additive.
- (c) μ_0 is countably additive if $\mu_0(B_n) \rightarrow 0$ for cylinders $B_n \downarrow \emptyset$ as $n \rightarrow \infty$.
- (d) If $\mu_0(B_n) \geq \delta$, then we can find decreasing $D_n \subset B_n$ such that $\mu_0(D_n) \geq \frac{\delta}{2}$ and $D_n = D_n^* \times \mathbb{R}^{\mathbb{N}}$ for a compact rectangle D_n^* .
- (e) If $\mu_0(B_n) \geq \delta$, then $\bigcap_{i=1}^{\infty} B_i$ is non-empty.

Proof. (d) Let $B_n = B_n^* \times \mathbb{R}^{\mathbb{N}}$ for a rectangle $B_n^* \subset \mathbb{R}^n$. By the inner regularity of $\mu_{r(n)}$, there is a compact rectangle $C_n^* \subset B_n^*$ such that

$$\mu_0(B_n \setminus C_n) = \mu_{r(n)}(B_n^* \setminus C_n^*) < \frac{\delta}{2^{n+1}}.$$

Let $C_n := C_n^* \times \mathbb{R}^{\mathbb{N}}$ and define $D_n := \bigcap_{i=1}^n C_i = D_n^* \times \mathbb{R}^{\mathbb{N}}$. Then,

$$\mu_0(B_n \setminus D_n) \leq \mu_0\left(\bigcup_{i=1}^n B_n \setminus C_i\right) \leq \mu_0\left(\bigcup_{i=1}^n B_i \setminus C_i\right) < \frac{\delta}{2},$$

which implies $\mu_0(D_n) \geq \frac{\delta}{2}$.

(e) Take any sequence $(\omega_n)_n$ in $\mathbb{R}^{\mathbb{N}}$ such that $\omega_n \in D_n$. Since each $D_n^* \subset \mathbb{R}^n$ is compact and non-empty, by diagonal argument, we have a subsequence $(\omega_k)_k$ such that ω_k is pointwise convergent, and its limit is contained in $\bigcap_{i=1}^{\infty} D_i \subset \bigcap_{i=1}^{\infty} B_i = \emptyset$, which is a contradiction that leads $\mu_0(B_n) \rightarrow 0$. \square

Part IV

Stochastic calculus