

Harmonic Functions Vanishing at Infinity on the Punctured Domain

Ikhan Choi

September 25, 2021.

1 Kelvin transform and decay rates

2 Potential field from a source

Theorem. *Let $d \geq 3$. A distribution $u \in \mathcal{D}'(\mathbb{R}^d)$ is a harmonic function on $\mathbb{R}^d \setminus \{0\}$ and vanishes at infinity if and only if there is a distribution $\rho \in \mathcal{D}'(\mathbb{R}^d)$ such that $u = \Phi * \rho$ and $\text{supp}(\rho) \subset \{0\}$, where Φ denotes the fundamental solution of Laplace's equation.*

Proof. (\Rightarrow) Define a distribution ρ by

$$\langle \rho, \varphi \rangle := -\langle u, \Delta \varphi \rangle$$

for $\varphi \in C_c^\infty(\mathbb{R}^d)$. In other words, $\rho = -\Delta u$ in distributional sense. Then, ρ has the support contained in $\{0\}$ because if $\varphi \in C_c^\infty(\mathbb{R}^d \setminus \{0\})$ then

$$\langle \rho, \varphi \rangle = -\langle u, \Delta \varphi \rangle = -\int u(x) \Delta \varphi(x) dx = -\int \Delta u(x) \varphi(x) dx = 0.$$

Therefore, we only need to verify $u = \Phi * \rho$ to complete the proof.

Let $\varphi \in C_c^\infty(\mathbb{R}^d)$. Be cautious that the argument

$$\langle \Phi * \rho, \varphi \rangle = \langle \rho, \Phi * \varphi \rangle = -\langle u, \Delta(\Phi * \varphi) \rangle = \langle u, \varphi \rangle$$

fails to provide a proof because the function $\Phi * \rho$ is not compactly supported so that we cannot deduce $\langle \rho, \Phi * \varphi \rangle = -\langle u, \Delta(\Phi * \varphi) \rangle$, and here we use the condition that

u vanishes at infinity to justify the equality. Define a cutoff function $\chi \in C_c^\infty(\mathbb{R}^d)$ such that

$$\chi(x) = \begin{cases} 1 & \text{if } |x| \leq \frac{5}{4} \\ 0 & \text{if } |x| \geq \frac{7}{4} \end{cases}.$$

If we denote $\chi_r(x) := \chi(\frac{x}{r})$, then we have

$$\langle \rho, (\Phi \chi_r) * \varphi \rangle = -\langle u, \Delta((\Phi \chi_r) * \varphi) \rangle$$

by the definition of ρ . We have the limit of the left-hand side

$$\lim_{r \rightarrow \infty} \langle \rho, (\Phi \chi_r) * \varphi \rangle = \langle \rho, \Phi * \varphi \rangle$$

because

$$\begin{aligned} \text{supp}((\Phi(1 - \chi_r) * \varphi)) &\subset \text{supp}(\Phi(1 - \chi_r)) + \text{supp}(\varphi) \\ &\subset \mathbb{R}^d \setminus B(0, 2R) + \overline{B}(0, R) = \mathbb{R}^d \setminus B(0, R) \end{aligned}$$

for all $r > 2R$ so that the supports of $\Phi(1 - \chi_r) * \varphi$ and ρ are disjoint, where we define $R := \sup_{x \in \text{supp}(\varphi)} |x|$. However, the right-hand limit

$$-\lim_{r \rightarrow \infty} \langle u, \Delta((\Phi \chi_r) * \varphi) \rangle = -\langle u, \Delta(\Phi * \varphi) \rangle$$

is not a trivial result.

Assuming $\chi(x) = \chi(-x)$ without loss of generality, we have

$$\langle u, \Delta(\Phi(1 - \chi_r) * \varphi) \rangle = \langle u * \Delta(\Phi(1 - \chi_r)), \varphi \rangle.$$

Because

$$\Delta_y \left[\Phi(x - y) \left(1 - \chi\left(\frac{x-y}{r}\right) \right) \right] = 0$$

for $|y| < R$ and $x \in \text{supp}(\varphi)$ if $r > 2R$, we can write

$$\langle u * \Delta(\Phi(1 - \chi_r)), \varphi \rangle = \int \varphi(x) \int u(y) \Delta_y \left[\Phi(x - y) \left(1 - \chi\left(\frac{x-y}{r}\right) \right) \right] dy dx.$$

We compute

$$\begin{aligned} \Delta_y \left[\Phi(x - y) \left(1 - \chi\left(\frac{x-y}{r}\right) \right) \right] &= 2 \nabla \Phi(x - y) \cdot \frac{1}{r} \nabla \chi\left(\frac{x-y}{r}\right) - \Phi(x - y) \frac{1}{r^2} \Delta \chi\left(\frac{x-y}{r}\right) \\ &= -\frac{2}{\omega_d} \frac{x - y}{|x - y|^d} \cdot \frac{1}{r} \nabla \chi\left(\frac{x-y}{r}\right) - \frac{1}{(d-2)\omega_d} \frac{1}{|x - y|^{d-2}} \frac{1}{r^2} \Delta \chi\left(\frac{x-y}{r}\right). \end{aligned}$$

Then, since $\frac{5}{4}r \leq |x - y| \leq \frac{7}{4}r$ if $\nabla \chi(\frac{x-y}{r}) \neq 0$ and $\Delta \chi(\frac{x-y}{r}) \neq 0$, we obtain

$$\left| \Delta_y \left[\Phi(x-y) \left(1 - \chi\left(\frac{x-y}{r}\right) \right) \right] \right| \leq C \frac{1}{r^d} \psi\left(\frac{x-y}{r}\right)$$

for some constant $C > 0$, where

$$\psi(y) := |\nabla \chi(y)| + |\Delta \chi(y)|.$$

For each $x \in \text{supp}(\varphi)$, since we have $\frac{5}{4}r \leq |x - y| \leq \frac{7}{4}r$ implies $r \leq |y| \leq 2r$ if $r > 4R$, it follows that

$$\begin{aligned} \left| \int u(y) \Delta_y \left[\Phi(x-y) \left(1 - \chi\left(\frac{x-y}{r}\right) \right) \right] dy \right| &\leq C \int |u(y)| \frac{1}{r^d} \psi\left(\frac{x-y}{r}\right) dy \\ &\leq C \max_{r \leq |y| \leq 2r} u(y) \end{aligned}$$

converges to zero as $r \rightarrow \infty$. By the bounded convergence theorem, we can deduce

$$\lim_{r \rightarrow \infty} \int \varphi(x) \int u(y) \Delta_y \left[\Phi(x-y) \left(1 - \chi\left(\frac{x-y}{r}\right) \right) \right] dy dx = 0,$$

so we are done.

(\Leftarrow) Let $\varphi \in C_c^\infty(\mathbb{R}^d \setminus \{0\})$. Since

$$\langle \Phi * \rho, \Delta \varphi \rangle = \langle \rho, \Phi * (\Delta \varphi) \rangle = \langle \rho, \varphi \rangle = 0,$$

the distribution $\Phi * \rho$ on $\mathbb{R}^d \setminus \{0\}$ is weakly harmonic, and by Weyl's lemma for distributions, it is a smooth harmonic function on $\mathbb{R}^d \setminus \{0\}$.

Since ρ is supported at zero, we have a positive integer k and constants a_α such that

$$|\langle \rho, \varphi \rangle| \leq \sum_{|a| \leq k} |a_\alpha D^\alpha \varphi(0)|$$

for $\varphi \in C^\infty(\mathbb{R}^d)$. Then, for non-zero $x \in \mathbb{R}^d$, by taking a cutoff function $\chi \in C_c^\infty(\mathbb{R}^d)$ such that

$$\chi(y) = \begin{cases} 1 & \text{if } |y - x| \leq \frac{1}{3}|x| \\ 0 & \text{if } |y| \leq \frac{1}{3}|x| \end{cases},$$

we have

$$|\Phi * \rho(x)| = |(\Phi \chi) * \rho(x)| = |\langle \rho(x-y), \Phi(y) \chi(y) \rangle_y| \leq \sum_{|a| \leq k} |a_\alpha D^\alpha \Phi(x)| = O(r^{2-d})$$

as $r \rightarrow \infty$. Therefore, $\Phi * \rho$ vanishes at infinity. \square

Lemma. Let ρ be a distribution on \mathbb{R}^d such that $\text{supp}(\rho) \subset \{0\}$. Then, there is a constant coefficient partial differential operator $P(D)$ such that $\rho = P(D)\delta$.

Corollary. Let $d \geq 3$. If a distribution $u \in \mathcal{D}'(\mathbb{R}^d)$ is a harmonic function on $\mathbb{R}^d \setminus \{0\}$ and vanishes at infinity, then there are an integer $k \geq 0$ and constants a_α such that

$$u(x) = \sum_{|\alpha| \leq k} a_\alpha D^\alpha \Phi(x)$$

for $x \neq 0$, where Φ denotes the fundamental solution of Laplace's equation.