

Differential Equations

Ikhan Choi

November 20, 2023

Contents

I	Linear ordinary differential equations	3
1	Initial value problems	4
1.1	Constant coefficient equations	4
1.2	Variable coefficient equations	4
2	Boundary value problems	5
2.1	Second order linear equations	5
2.2	Orthogonal polynomials	5
2.3	Sturm-Liouville theory	5
3	Inhomogeneous problems	6
3.1	Method of undetermined coefficients	6
3.2	Variation of parameters	6
3.3	Laplace transform	6
II	Nonlinear ordinary differential equations	7
4	First order nonlinear equations	8
4.1	Local existence theorems	8
4.2	Implicit equations	9
4.3	Global existence	9
5	Dynamical systems	10
5.1	Equilibrium and stability	10
5.2	Autonomous systems	10
5.3	Hamiltonian systems	10
5.4	Planar systems	10
6	Chaos	11
III	Linear partial differential equations	12
7	Laplace's equation	13
7.1	Harmonic functions	13
7.2	Poisson equation	13
7.3	Eigenvalue problems	14

8 Heat equation	15
8.1 Heat kernel	15
8.2 Separation of variables	15
9 Wave equation	16
9.1 First order partial differential equations	16
9.2 Initial value problems	16
9.3 Boundary value problems	16
9.4 Dispersive equations	16
IV Nonlinear partial differential equations	17
10 Geometric PDEs	18
11 Fluid dynamics	19
11.1 Conservation laws	19
11.2 Euler and Burger equation	19
11.3 Non-linear waves	19
11.4 Navier-Stokes equation	19

Part I

Linear ordinary differential equations

Chapter 1

Initial value problems

1.1 Constant coefficient equations

existence uniqueness system of equations characteristic equations complex roots repeated roots

1.2 Variable coefficient equations

existence uniqueness series solution Frobenius method Fuch's theorem

Chapter 2

Boundary value problems

2.1 Second order linear equations

Helmholtz Bessel Legendre Hermite Laguerre

2.2 Orthogonal polynomials

L^2 space

2.3 Sturm-Liouville theory

Eigenvalue problems boundary conditions

Exercises

2.1 (Rayleigh-Ritz principle).

Chapter 3

Inhomogeneous problems

3.1 Method of undetermined coefficients

3.2 Variation of parameters

3.3 Laplace transform

discontinuous data gluing

Exercises

3.1 (Damped oscillation).

Part II

Nonlinear ordinary differential equations

Chapter 4

First order nonlinear equations

4.1 Local existence theorems

4.1 (Picard-Lindelöf theorem). Consider the following initial value problem:

$$x'(t) = f(t, x(t)), \quad x(0) = x_0.$$

Construct an approximate solution $(x_n)_{n=0}^\infty$ defined inductively such that $x_0(t) \equiv x_0$ and

$$x'_{n+1}(t) = f(t, x_n(t)), \quad x_{n+1}(0) = x_0.$$

Suppose f satisfies

$$|f(t, x)| \leq \frac{R}{T}, \quad |f(t, x) - f(t, y)| \lesssim |x - y|$$

on the cylinder $[0, T] \times \overline{B(x_0, R)}$.

- (a) x_n is in $C^1([0, T], \overline{B(x_0, R)})$.
- (b) x_n is Cauchy in $C^1([0, T], \overline{B(x_0, R)})$.
- (c) The equation has a unique solution in $C^1([0, T], \overline{B(x_0, R)})$.

Proof. (a) It clearly follows from the explicit formula

$$x_{n+1}(t) = x_0 + \int_0^t f(s, x_n(s)) ds.$$

(b) Since

$$|x_1(t) - x_0(t)| \leq \int_0^t |f(s, x_0)| ds \leq Mt$$

and

$$\begin{aligned} |x_{n+1}(t) - x_n(t)| &\leq \int_0^t |f(s, x_n(s)) - f(s, x_{n-1}(s))| ds \\ &\leq K \int_0^t |x_n(s) - x_{n-1}(s)| ds \\ &\leq MK^n \int_0^t \frac{s^n}{n!} ds \\ &= MK^n \frac{t^{n+1}}{(n+1)!}, \end{aligned}$$

we have the convergent series

$$\sum_{n=0}^{\infty} \|x_{n+1} - x_n\|_{\infty} \leq TM \frac{e^{KT} - 1}{KT}.$$

Also,

$$|x'_{n+1}(t) - x'_n(t)| \leq |f(t, x_n(t)) - f(t, x_{n-1}(t))| \leq K|x_n(t) - x_{n-1}(t)| \leq MK^{n+1} \frac{t^{n+1}}{(n+1)!}.$$

(c) Limiting check.

□

4.2 (Cauchy-Peano theorem).

4.3 (Carathéodory existence theorem).

4.2 Implicit equations

integrating factor, separable equations, exact equations

4.3 Global existence

4.4 (Gronwall's inequality).

4.5 (A priori estimate).

Chapter 5

Dynamical systems

5.1 Equilibrium and stability

Bifurcations

Stability theory Lyapunov, invariant set

5.2 Autonomous systems

5.3 Hamiltonian systems

5.4 Planar systems

periodic orbit

5.1 (Poincaré-Bendixon).

Exercises

5.2 (Undamped pendulum).

$$x''(t) + \sin x(t) = 0$$

5.3 (Approximated pendulum).

$$x''(t) + x(t) - \frac{1}{6}x(t)^3 = \alpha$$

5.4 (Van der Pol oscillator).

$$x''(t) - \mu(1 - x(t)^2)x'(t) + x(t) = 0$$

5.5 (Lotka-Volterra model). Also known as predator-prey equations.

Chapter 6

Chaos

Attractors

Part III

Linear partial differential equations

Chapter 7

Laplace's equation

7.1 Harmonic functions

7.1 (Mean value property).

7.2 (Maximum principle).

7.3 (Newtonian potential).

7.4 (Dirichlet problem for half space).

7.5 (Dirichlet problem for open ball).

7.2 Poisson equation

7.6 (Weak derivative).

7.7 (Dirac delta function). Let Ω be an open subset of \mathbb{R}^d . The *Dirac delta function* is a linear functional $\delta : C_c^\infty(\Omega) \rightarrow \mathbb{R}$ defined by $\delta(\varphi) := \varphi(0)$. We conventionally use the function-like notation $\delta(x)$ to denote $\varphi(0)$ by

$$\int \delta(x) \varphi(x) dx.$$

7.8 (Fundamental solution of the Laplace equation). Let $d \geq 2$. The *Fundamental solution of the Laplace equation* is a function $\Phi : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}$ that solves the boundary value problem

$$\begin{cases} -\Delta \Phi(x) = \delta(x) & \text{in } \mathbb{R}^d, \\ \Phi(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty. \end{cases}$$

(a) The fundamental solution is given by

$$\Phi(x) := \begin{cases} -\frac{1}{2\pi} \log |x| & \text{if } d = 2 \\ \frac{1}{(d-2)\omega_d} \frac{1}{|x|^{d-2}} & \text{if } d \geq 3 \end{cases}.$$

In particular, Φ and $\nabla \Phi$ are locally integrable on \mathbb{R}^d but $\nabla^2 \Phi$ is not.

(b) For $u \in C_0^2(\mathbb{R}^d)$,

$$u(x) = - \int \Phi(x-y) \Delta u(y) dy.$$

Proof. Note that $\nabla\Phi(y) \cdot \nabla u(x-y)$ is integrable in y . Then,

$$\begin{aligned} -\int \Phi(y)\Delta u(x-y)dy &= -\int \nabla\Phi(y) \cdot \nabla u(x-y)dy \\ &= -\lim_{\varepsilon \rightarrow \infty} \int_{|y| \geq \varepsilon} \nabla\Phi(y) \cdot \nabla u(x-y)dy \\ &= -\lim_{\varepsilon \rightarrow \infty} \int_{|y|=\varepsilon} \nabla\Phi(y)u(x-y) \cdot \nu dS. \end{aligned}$$

Since

$$\nabla\Phi(x) = -\frac{1}{\omega_d} \frac{x}{|x|^d}, \quad \nu = \frac{x}{|x|},$$

we get

$$-\int \Phi(y)\Delta u(x-y)dy = \lim_{\varepsilon \rightarrow \infty} \frac{1}{\omega_d \varepsilon^{d-1}} \int_{|y|=\varepsilon} u(x-y) dS_y = u(x).$$

□

7.9 (Green's function of the Poisson equation). Let Ω be a bounded open subset of \mathbb{R}^d for $d \geq 2$. *Green's function of the Poisson equation* is a function $G : \Omega^2 \setminus \{(x, x) \in \Omega\} \rightarrow \mathbb{R}$ that solves the boundary value problem

$$\begin{cases} -\Delta_y G(x, y) = \delta(x-y) & \text{in } y \in \Omega \setminus \{x\}, \\ G(x, y) = 0 & \text{on } y \in \partial\Omega. \end{cases}$$

for each $x \in \Omega$.

Define $\phi : \Omega^2 \rightarrow \mathbb{R}$ to be a function that solves the boundary value problem

$$\begin{cases} -\Delta_y \phi(x, y) = 0 & \text{in } y \in \Omega, \\ \phi(x, y) = \Phi(x-y) & \text{on } y \in \partial\Omega. \end{cases}$$

for each $x \in \Omega$. Assume for the domain Ω that there exists a unique ϕ .

(a) Green's function is given by

$$G(x, y) = \Phi(x-y) - \phi(x, y),$$

where Φ is the fundamental solution of the Laplace equation. Physically, $y \mapsto -\phi(x, y)$ has a meaning of the electric potential generated by the induced surface charge of a grounded conductor provided a point charge is at x .

(b) The *Green representation formula* holds: for $u \in C^2(\Omega) \cap C(\overline{\Omega})$,

$$u(x) = -\int_{\Omega} G(x, y)\Delta u(y)dy - \int_{\partial\Omega} u(y)\nabla_y G(x, y) \cdot \nu dS_y.$$

7.10 (Existence and uniqueness of Poisson equation). representation formulas describe the solution assuming

7.3 Eigenvalue problems

Chapter 8

Heat equation

8.1 Heat kernel

Duhamel's principle

8.2 Separation of variables

Chapter 9

Wave equation

9.1 First order partial differential equations

9.2 Initial value problems

d'Alembert

Kirchhoff

odd reflection

9.3 Boundary value problems

Dirichlet, Neumann, Mixed

9.4 Dispersive equations

Part IV

Nonlinear partial differential equations

Chapter 10

Geometric PDEs

gradient flow curvature flow

Chapter 11

Fluid dynamics

11.1 Conservation laws

11.2 Euler and Burger equation

11.3 Non-linear waves

Nonlinear diffusion?

11.4 Navier-Stokes equation