

Geometry

Ikhan Choi

April 23, 2024

Contents

I	Classical geometry	3
1	Euclidean geometry	4
1.1	Plane geometry	4
1.2	Solid geometry	4
1.3	Axiomatization	4
2	Non-Euclidean geometry	5
2.1	Absolute geometry	5
2.2	Spherical and elliptic geometry	5
2.3	Hyperbolic geometry	5
3	Non-metric geometry	6
3.1	Ordered and incidence geometry	6
3.2	Affine and projective geometry	6
3.3	Conformal and inversive geometry	6
II	Smooth surfaces	7
4	Smooth manifolds	8
4.1	Local coordinates	8
4.2	Space curves	8
4.3	Space surfaces	8
5	Fundamental forms	10
5.1	Riemannian metrics	10
5.2	Gaussian curvatures	10
6		13
III	Riemann surfaces	14
7	Riemann-Roch theorem	15
8	Algebraic curves	17
8.1	17
8.2	17
9	Uniformization	18

IV Topological surfaces	19
10 Fundamental groups	20
10.1 Homotopy	20
10.2 Covering spaces	20
11 Homology groups	21
11.1 Singular homology	21
11.2 Simplicial homology	21
11.3 Cellular homology	21
12 Classification of surfaces	22
12.1 Combinatorial surfaces	22

Part I

Classical geometry

Chapter 1

Euclidean geometry

1.1 Plane geometry

1.2 Solid geometry

1.3 Axiomatization

Chapter 2

Non-Euclidean geometry

2.1 Absolute geometry

axioms 1 to 4

2.2 Spherical and elliptic geometry

axioms 2 and 4

2.3 Hyperbolic geometry

axiomes 1 to 4

Models of hyperbolic geometry (metric description) Elementary figures Isometries Length, volume, angle

Chapter 3

Non-metric geometry

3.1 Ordered and incidence geometry

axioms 1 and 2

3.2 Affine and projective geometry

axioms 1,2,5

3.3 Conformal and inversive geometry

Part II

Smooth surfaces

Chapter 4

Smooth manifolds

4.1 Local coordinates

4.2 Space curves

4.3 Space surfaces

Reparametrizations

Theorem 4.3.1. *Let S be a regular surface. Let v, w be linearly independent tangent vectors in $T_p S$ for a point $p \in S$. Then, S admits a parametrization α such that $\alpha_x|_p = v$ and $\alpha_y|_p = w$.*

Theorem 4.3.2. *Let X, Y be linearly independent tangent vector fields on a regular surface S . Then, S admits a parametrization α such that $\alpha_x|_p$ and $\alpha_y|_p$ are parallel to $X|_p, Y|_p$ respectively for each $p \in S$.*

Theorem 4.3.3. *Let X, Y be linearly independent tangent vector fields on a regular surface S . If $\partial_X Y = \partial_Y X$, then S admits a parametrization α such that $\alpha_x|_p = X|_p$ and $\alpha_y|_p = Y|_p$ for each $p \in S$.*

Let S be a regular surface embedded in \mathbb{R}^3 . The inner product on $T_p S$ induced from the standard inner product of \mathbb{R}^3 can be represented not only as a matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

in the basis $\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\} \subset \mathbb{R}^3$, but also as a matrix

$$\begin{pmatrix} \langle \alpha_x, \alpha_x \rangle & \langle \alpha_x, \alpha_y \rangle \\ \langle \alpha_y, \alpha_x \rangle & \langle \alpha_y, \alpha_y \rangle \end{pmatrix}$$

in the basis $\{\alpha_x|_p, \alpha_y|_p\} \subset T_p S$.

Definition 4.3.4. *Metric coefficients*

$$\begin{aligned} \langle \alpha_x, \alpha_x \rangle &=: g_{11} & \langle \alpha_x, \alpha_y \rangle &=: g_{12} \\ \langle \alpha_y, \alpha_x \rangle &=: g_{21} & \langle \alpha_y, \alpha_y \rangle &=: g_{22} \end{aligned}$$

Theorem 4.3.5 (Normal coordinates). ...?

Differentiation of tangent vectors

Definition 4.3.6. Let $\alpha : U \rightarrow \mathbb{R}^3$ be a regular surface. The *Gauss map* or *normal unit vector* $\nu : U \rightarrow \mathbb{R}^3$ is a vector field on α defined by:

$$\nu(x, y) := \frac{\alpha_x \times \alpha_y}{\|\alpha_x \times \alpha_y\|}(x, y).$$

The set of vector fields $\{\alpha_x|_p, \alpha_y|_p, \nu|_p\}$ forms a basis of $T_p\mathbb{R}^3$ at each point p on α . The Gauss map is uniquely determined up to sign as α changes.

Definition 4.3.7 (Gauss formula, Γ_{ij}^k, L_{ij}). Let $\alpha : U \rightarrow \mathbb{R}^3$ be a regular surface. Define indexed families of smooth functions $\{\Gamma_{ij}^k\}_{i,j,k=1}^2$ and $\{L_{ij}\}_{i,j=1}^2$ by the Gauss formula

$$\begin{aligned} \alpha_{xx} &= \Gamma_{11}^1 \alpha_x + \Gamma_{11}^2 \alpha_y + L_{11} \nu, & \alpha_{xy} &= \Gamma_{12}^1 \alpha_x + \Gamma_{12}^2 \alpha_y + L_{12} \nu, \\ \alpha_{yx} &= \Gamma_{21}^1 \alpha_x + \Gamma_{21}^2 \alpha_y + L_{21} \nu, & \alpha_{yy} &= \Gamma_{22}^1 \alpha_x + \Gamma_{22}^2 \alpha_y + L_{22} \nu. \end{aligned}$$

The *Christoffel symbols* refer to eight functions $\{\Gamma_{ij}^k\}_{i,j,k=1}^2$. The Christoffel symbols and L_{ij} do depend on α .

We can easily check the symmetry $\Gamma_{ij}^k = \Gamma_{ji}^k$ and $L_{ij} = L_{ji}$. Also,

$$\begin{aligned} \partial_X Y &= X^i \partial_i (Y^j \alpha_j) \\ &= X^i (\partial_i Y^j) \alpha_j + X^i Y^j \partial_i \alpha_j \\ &= (X^i \partial_i Y^j + X^i Y^j \Gamma_{ij}^k) \alpha_k + X^i Y^j L_{ij} \nu. \end{aligned}$$

Differentiation of normal vector

The partial derivative $\partial_X \nu$ is a tangent vector field since

$$\langle \partial_X \nu, \nu \rangle = \frac{1}{2} \partial_X \langle \nu, \nu \rangle = 0.$$

Therefore, we can define the following useful operator.

Definition 4.3.8. Let S be a regular surface embedded in \mathbb{R}^3 . The *shape operator* is $S : \mathfrak{X}(S) \rightarrow \mathfrak{X}(S)$ defined as

$$S(X) := -\partial_X \nu.$$

Proposition 4.3.9. The shape operator is self-adjoint, i.e. symmetric.

Proof. Recall that $\partial_X Y - \partial_Y X$ is a tangent vector field. Then,

$$\langle X, S(Y) \rangle = \langle X, -\partial_Y \nu \rangle = \langle \partial_Y X, \nu \rangle = \langle \partial_X Y, \nu \rangle = \langle S(X), Y \rangle. \quad \square$$

Theorem 4.3.10. Let $\alpha : U \rightarrow \mathbb{R}^3$ be a regular surface and S be the shape operator. Then S has the coordinate representation

$$S = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix}^{-1} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix}$$

with respect to the frame $\{\alpha_x, \alpha_y\}$ for tangent spaces. In other words, if we let $X = X^i \alpha_i$ and $S(X) = S(X)^j \alpha_j$, then

$$\begin{pmatrix} S(X)^1 \\ S(X)^2 \end{pmatrix} = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix}^{-1} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} \begin{pmatrix} X^1 \\ X^2 \end{pmatrix}.$$

Proof. Let $S(X)^j = S_i^j X^i$. Then,

$$g_{ik} X^i S_j^k Y^j = \langle X, S(Y) \rangle = \langle \partial_X Y, \nu \rangle = X^i Y^j L_{ij}$$

implies $g_{ik} S_j^k = L_{ij}$. \square

Chapter 5

Fundamental forms

5.1 Riemannian metrics

5.2 Gaussian curvatures

Theorema egregium surfaces of constant gaussian curvature

Definition 5.2.1. Let $\alpha : U \rightarrow \mathbb{R}^3$ be a regular surface.

$$\begin{aligned} E &:= \langle \alpha_x, \alpha_x \rangle = g_{11}, & F &:= \langle \alpha_x, \alpha_y \rangle = g_{12}, & G &:= \langle \alpha_y, \alpha_y \rangle = g_{22}, \\ L &:= \langle \alpha_{xx}, \nu \rangle = L_{11}, & M &:= \langle \alpha_{xy}, \nu \rangle = L_{12}, & N &:= \langle \alpha_{yy}, \nu \rangle = L_{22}. \end{aligned}$$

Corollary 5.2.2. We have $GM - FN = EM - FL$, and the Weingarten equations:

$$\begin{aligned} \nu_x &= \frac{FM - GL}{EG - F^2} \alpha_x + \frac{FL - EM}{EG - F^2} \alpha_y, \\ \nu_y &= \frac{FN - GM}{EG - F^2} \alpha_x + \frac{FM - EN}{EG - F^2} \alpha_y. \end{aligned}$$

Theorem 5.2.3.

$$\Gamma_{ij}^l = \frac{1}{2} g^{kl} (g_{ik,j} - g_{ij,k} + g_{kj,i}).$$

$$\frac{1}{2} (\log g)_x = \Gamma_{11}^1.$$

$$\nu_x \times \nu_y = K \sqrt{\det g} \, \nu.$$

$$\alpha_x \times \alpha_y = \sqrt{\det g} \, \nu$$

$$\langle \nu_x \times \nu_y, \alpha_x \times \alpha_y \rangle = \det \begin{pmatrix} \langle \nu_x, \alpha_x \rangle & \langle \nu_x, \alpha_y \rangle \\ \langle \nu_y, \alpha_x \rangle & \langle \nu_y, \alpha_y \rangle \end{pmatrix} = \det \begin{pmatrix} -L & -M \\ -M & -N \end{pmatrix} = K \det g$$

5.1 (Gaussian curvature formula). (a) In general,

$$K = \frac{LN - M^2}{EG - F^2}.$$

(b) For orthogonal coordinates such that $F \equiv 0$,

$$K = -\frac{1}{2\sqrt{\det g}} \left(\left(\frac{1}{\sqrt{\det g}} E_y \right)_y + \left(\frac{1}{\sqrt{\det g}} G_x \right)_x \right).$$

(c) For $f(x, y, z) = 0$,

$$K = -\frac{1}{|\nabla f|^4} \begin{vmatrix} 0 & \nabla f \\ \nabla f^T & \text{Hess}(f) \end{vmatrix},$$

where ∇f denotes the gradient $\nabla f = (f_x, f_y, f_z)$.

(d) (Beltrami-Enneper) If τ is the torsion of an asymptotic curve, then

$$K = -\tau^2.$$

(e) (Brioschi) E, F, G describes K .

Proof. (a) Clear.

(b) We have $GM = EM$ and

$$\begin{aligned} v_x &= -\frac{L}{E}\alpha_x - \frac{M}{G}\alpha_y, & v_y &= -\frac{M}{E}\alpha_x - \frac{N}{G}\alpha_y. \\ v_x \times v_y &= \frac{LN - M^2}{EG}\alpha_x \times \alpha_y \end{aligned}$$

After curvature tensors...

□

5.2 (Computation of Gaussian curvatures). (a) (Monge's patch) For $(x, y, f(x, y))$,

$$K = \frac{f_{xx}f_{yy} - f_{xy}^2}{(1 + f_x^2 + f_y^2)^2}.$$

(b) (Surface of revolution). Let $\gamma(t) = (r(t), z(t))$ be a plane curve with $r(t) > 0$. If $t \mapsto (r(t), z(t))$ is a unit-speed curve, then

$$K = -\frac{r''}{r}.$$

(c) (Models of hyperbolic planes)

Proof. (b) Let

$$\alpha(\theta, t) = (r(t)\cos\theta, r(t)\sin\theta, z(t))$$

be a parametrization of a surface of revolution. Then,

$$\begin{aligned} \alpha_\theta &= (-r(t)\sin\theta, r(t)\cos\theta, 0) \\ \alpha_t &= (r'(t)\cos\theta, r'(t)\sin\theta, z'(t)) \\ v &= \frac{1}{\sqrt{r'(t)^2 + z'(t)^2}}(z'(t)\cos\theta, z'(t)\sin\theta, -r'(t)), \end{aligned}$$

and

$$\begin{aligned} \alpha_{\theta\theta} &= (-r(t)\cos\theta, -r(t)\sin\theta, 0) \\ \alpha_{\theta t} &= (-r'(t)\sin\theta, r'(t)\cos\theta, 0) \\ \alpha_{tt} &= (r''(t)\cos\theta, r''(t)\sin\theta, z''(t)). \end{aligned}$$

Thus we have

$$E = r(t)^2, \quad F = 0, \quad G = r'(t)^2 + z'(t)^2,$$

and

$$L = -\frac{r(t)z'(t)}{\sqrt{r'(t)^2 + z'(t)^2}}, \quad M = 0, \quad N = \frac{r''(t)z'(t) - r'(t)z''(t)}{\sqrt{r'(t)^2 + z'(t)^2}}.$$

Therefore,

$$K = \frac{LN - M^2}{EG - F^2} = \frac{z'(r'z'' - r''z')}{r(r'^2 + z'^2)^2}.$$

In particular, if $t \mapsto (r(t), z(t))$ is a unit-speed curve, then

$$K = -\frac{r''}{r}.$$

□

5.3 (Local isomorphism). Surfaces of the same constant Gaussian curvature are locally isomorphic.

Proof. Let

$$\begin{pmatrix} \|\alpha_r\|^2 & \langle \alpha_r, \alpha_t \rangle \\ \langle \alpha_t, \alpha_r \rangle & \|\alpha_t\|^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & h(r, t)^2 \end{pmatrix}$$

be the first fundamental form for a geodesic coordinate chart along a geodesic curve so that α_{tt} and α_{rr} are normal to the surface. Then,

$$K = -\frac{h_{rr}}{h}$$

is constant. Also, since

$$\frac{1}{2}(h^2)_r + \langle \alpha_r, \alpha_{tt} \rangle = \langle \alpha_{rt}, \alpha_t \rangle + \langle \alpha_r, \alpha_{tt} \rangle = \langle \alpha_r, \alpha_t \rangle_t = 0$$

implies $h_r = 0$ at $r = 0$, the function $f : r \mapsto h(r, t)$ satisfies the following initial value problem

$$f_{rr} = -Kf, \quad f(0) = 1, \quad f'(0) = 0.$$

Therefore, h is uniquely determined by K .

□

Chapter 6

Part III

Riemann surfaces

Chapter 7

Riemann-Roch theorem

Let X be a compact Riemann surface. Consider a vector space $\mathcal{M}^\times(X) \cup \{0\}$.

$$L(D) := H^0(X, \mathcal{O}(D)) = \{f \in \mathcal{M}^\times(X) : (f) + D \geq 0\} \cup \{0\}.$$

$$\text{Div}(X) = H^0(X, \mathcal{M}^\times / \mathcal{O}^\times) = \Gamma(\mathcal{M}^\times / \mathcal{O}^\times).$$

$$\text{Pic}(X) = H^1(X, \mathcal{O}^\times).$$

First Chern class $H^1(X, \mathcal{O}^\times) \rightarrow H^2(X, \mathbb{Z})$.

7.1. Let X be a compact Riemann surface. A *Weil divisor* D on X is an element of the free abelian group $\text{Div}(X)$ generated by points of X . By compactness of X , a meromorphic function $f \in \mathcal{M}(X)$ gives rise to a divisor $(f) := \sum_{p \in X} \text{ord}_p(f)p$. Such a divisor is called a *principal divisor*.

Let $D = \sum n_i p_i$ on X be a Weil divisor on X . Each point $P \in X$ has a meromorphic function f on an open neighborhood U of P such that $(f) = D$ on U . It implies that there is a collection $\{f_\alpha\}$ of meromorphic functions f_α defined on U_α , where $\{U_\alpha\}$ is an open cover of X , such that f_α/f_β is a well-defined holomorphic functions on $U_\alpha \cap U_\beta$. The collection $\{f_\alpha\}$ is called a *Cartier divisor*.

A Cartier divisor defines a line bundle.

7.2. Given $\{p_i\}_{i=1}^n$ points and $\{f_i\}_{i=1}^n$ principal parts, there is a meromorphic function f with pre-described principal parts if and only if for every holomorphic 1-form ω we have $\sum_{i=1}^n \text{Res}(f_i \omega, p_i) = 0$.

7.3.

$$l(D) - l(K - D) = \deg(D) + 1 - g.$$

The genus can be defined by $g = h^0(X, \Omega^1)$. For algebraic curves, it can be proved as follows: Assuming the Serre duality, we have $\chi(D) = h^0(D) - h^1(D) = l(D) - l(K - D)$ and $\chi(0) = h^0(0) - h^1(0) = 1 - g$. Then, the Riemann-Roch is boiled down to $\chi(D) = \deg(D) + \chi(0)$, which can be shown inductively.

However, we want to prove a compact Riemann surface is projective as an application of the Riemann-Roch theorem, we need to prove the Riemann-Roch theorem without theory of algebraic curves.

(a) If $\deg D < 0$, then $l(D) = 0$.

Proof. (a) Let $f \in L(D) \setminus \{0\}$. Then, $(f) + D \geq 0$ and $\deg(f) = 0$ imply $\deg D \geq 0$, which is a contradiction.

(b) Let $D = 0$. Then, it follows from $l(K) = g$ and $l(0) = 1$.

Let $D > 0$. We may assume $D = \sum_{i=1}^n n_i p_i$ with $n_i > 0$. (why?) Let

$$V_i := \left\{ \sum_{k=-n_i}^{-1} c_k (z - p_i)^k : c_k \in \mathbb{C} \right\}$$

and $V := \bigoplus_{i=1}^n V_i$. (how can we define the principal part of f on Riemann surface?) Then, $\dim V = \deg D$. Define $L(D) \rightarrow V$ by principal part at each point p_i . □

7.4 (Embedding theorem). Let X be a compact Riemann surface. The *complete linear system* of a divisor D on X is

$$|D| := \{(f) + D : f \in \mathcal{O}(X)\}.$$

Then, $|D|$ can be identified with the projective space $(L(D) \setminus \{0\})/\mathbb{C}^\times = \mathbb{CP}^{l(D)-1}$. Let $(f_i)_{i=0}^{l(D)-1}$ be a basis of $L(D)$.

For a linear system Δ of projective dimension $n-1$, we can take (how?) a basis $(f_i)_{i=0}^{n-1}$ such that the following map is well-defined:

$$X \setminus \text{Bl}(\Delta) \rightarrow \mathbb{CP}^{n-1} : p \mapsto (f_0 : \cdots : f_{n-1}).$$

Chapter 8

Algebraic curves

8.1

multiplicities, Bezout theorem

8.2

divisors, line bundles euler characteristic (tangent line bundle degree $2-2g$, canonical line bundle $2g-2$)

$$L(D) := \Gamma(X, \mathcal{O}(D)) = H^0(X, \mathcal{O}(D))$$

Jacobian variety (moduli spaces....)

8.1 (Chow theorem). A complex submanifold of a projective space is algebraic.

Chapter 9

Uniformization

The uniformization theorem provides one philosophy to classify compact Riemann surfaces. The universal covering is one of the three: the Riemann sphere, the complex plane, and the open unit disk. Each compact Riemann surface is realized as a quotient of these model space with a properly discontinuous action.

- $g = 0$: Riemann sphere (spherical) \rightarrow Riemann sphere itself
- $g = 1$: complex plane (Euclidean) \rightarrow elliptic curves
- $g \geq 2$: open unit disk (hyperbolic) \rightarrow hyperbolic surfaces, classified by Fuchsian groups (with which properties?)

Part IV

Topological surfaces

Chapter 10

Fundamental groups

10.1 Homotopy

10.1. A *homotopy of paths* is a continuous map $h : I \times I \rightarrow X$ such that $h(0, \cdot) = x_0$ and

- (a) linear homotopy
- (b) reparametrization

10.2. The fundamental group is a group composition

10.3 (Van Kampen theorem).

10.2 Covering spaces

10.4 (Path lifting property). Let $p : Y \rightarrow X$ be a covering map. For a path $\gamma : [0, 1] \rightarrow X$ and a point $y_0 \in Y$ such that $p(y_0) = \gamma(0)$, there is a unique lift $\tilde{\gamma} : I \rightarrow Y$ of γ such that $\tilde{\gamma}(0) = y_0$.

As a corollary, if γ_0 and γ_1 are end-fixing homotopic and have lifts $\tilde{\gamma}_0$ and $\tilde{\gamma}_1$ such that $\tilde{\gamma}_0(0) = \tilde{\gamma}_1(0)$, then $\tilde{\gamma}_0$ and $\tilde{\gamma}_1$ are basepoint-preserving homotopic.

As a corollary, for $p(y_0) = x_0$, the induced map $p_* : \pi_1(Y, y_0) \rightarrow \pi_1(X, x_0)$ is injective.

Proof. (Uniqueness) The interval $[0, 1]$ can be replaced to any connected set.

(Existence) By the compactness of $[0, 1]$, there is an increasing finite sequence $(t_i)_{i=0}^n$ such that

$$t_0 = 0, \quad t_n = 1, \quad [t_i, t_{i+1}] \subset \gamma^{-1}(U_i), \quad 0 \leq i < n,$$

where U_i is trivializing p .

□

10.5 (Universal covering). connected, locally path connected, semi-locally simply connected

10.6 (Classification of covering spaces). connected, locally path connected, semi-locally simply connected $\pi_1(X, x_0)/p_*(\pi_1(Y, y_0)) \rightarrow p^{-1}(x_0)$ is always injective, and bijective if Y is path connected.

Chapter 11

Homology groups

11.1 Singular homology

11.2 Simplicial homology

11.3 Cellular homology

Chapter 12

Classification of surfaces

12.1 Combinatorial surfaces

triangulation orientability euler characteristic genus connected sum