

Three-dimensional Topology

Ikhan Choi

Lectured by Takahiro Kitayama
University of Tokyo, Spring 2023

June 20, 2023

Contents

1	Day 1: April 11	2
1.1	Fundamental groups of higher dimensional manifolds	2
1.2	Surfaces and their groups	3
2	Day 2: April 18	3
2.1	Spherical manifolds	4
2.2	Fibered manifolds	4
2.3	Heegaard decomposition	5
2.4	Dehn surgery	5
3	Day 3: April 25	7
3.1	Alexander's theorem	7
3.2	Irreducible manifolds	7
3.3	Normal surfaces	8
4	Day 4: May 9	9
3.4	Prime decomposition theorem	9
4.1	Loop theorem	10
5	Day 5: May 16	11
6	Day 6: May 23	13
5.1	Essential surfaces	13
5.2	Haken hierarchy	14
7	Day 7: June 6	15
6.1	Seifert fibrations	15
8	Day 8: June 13	17
6.2	Classifications	17
9	Day 9: June 20	19
7.1	Existence	19
7.2	Uniqueness	19

1 Day 1: April 11

Plan:

1. Fundamental groups of manifolds
2. Examples and constructions
3. Prime decomposition
4. Loop and sphere theorems
5. Haken manifolds
6. Seifert manifolds
7. JSJ composition
8. Geometrization
9. Virtually special theorems

References:

1. J. Hempel, 3-manifolds
2. W. Jaco, Lectures on three-manifold topology
3. B. Martelli, An introduction to geometric topology
4. Morimoto, An introduction to three-dimensional manifolds (Japanses)

Grading: Submit a report for any three among the exercises given in the lecture (ITC-LMS Kadai).

Cancellation of class: 5/2, 7/11(maybe)

Convention:

- manifold = connected compact orientable smooth manifold
- surface = connected compact orientable smooth 2-dimensional manifold
- tub nbd, isotopy extension, transversality, triangulation,...

1. Fundamental group

1.1 Fundamental groups of higher dimensional manifolds

Proposition 1.1. *Let π be a finitely presented group. Then, for every $d \geq 4$ there is a d -manifold X such that $\pi_1(X) \cong \pi$.*

Proof. Let $\pi = \langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ be given. If $Y = (S^1 \times S^{d-1})^{\#n}$, then $\pi_1(Y) \cong \langle x_1, \dots, x_n \rangle$ by the van Kampen theorem. Let

$$Z = Y \setminus \left(\bigcup_{i=1}^m \nu(l_i) \right),$$

where $l_i \subset Y$ is embedded loops representing r_i and ν denotes the open tubular neighborhood. Then, $\partial Z = \bigcup_{i=1}^m l_i \times S^{d-2}$. Since $d \geq 4$, any loops and disks can be pushed off l_1, \dots, l_n , we have an isomorphism $\pi_1(Z) \rightarrow \pi_1(Y)$. Then, if we let

$$X = Z \cup_{\partial} \left(\bigcup_{i=1}^m D^2 \times S^{d-2} \right),$$

then $l_i \times * = \partial(D^2 \times *)$, we have $\pi_1(X) \cong \pi_1(Y) / \langle [l_1], \dots, [l_m] \rangle \cong \pi$. □

1.2 Surfaces and their groups

Theorem 1.2 (Radó, Whitehead). *Every topological surface admits a unique smooth and PL structure.*

Theorem 1.3. *Every surface is diffeomorphic to only one of $\Sigma_{g,b}$, where $\Sigma_{g,b} = (T^2)^{\#g} \# (D^2)^{\#b}$.*

Corollary 1.4. *$S = S^2, T^2, D^2$ are prime, that is, $S = S_1 \# S_2$ implies $S_i \approx S^2$ for $i = 1$ or $i = 2$.*

Remark 1.5. If the orientability is reduced out, then \mathbb{RP}^2 is prime. Also note that $T^2 \# \mathbb{RP}^2 \approx (\mathbb{RP}^2)^{\#3}$.

Theorem 1.6 (Uniformization). *For every surface $S \neq D^2$, its interior admits a complete Riemannian metric of constant curvature*

$$\begin{cases} 1, & \chi(S) > 0 \\ 0, & \chi(S) = 0 \\ -1, & \chi(S) < 0 \end{cases}$$

with universal covering $S^2, \mathbb{R}^2, \mathbb{H}^2$, respectively.

The hyperbolic plane is $\mathbb{H}^2 = \{(x, y) \in \mathbb{R}^2 : y > 0\}$ with the Riemannian metric $ds^2 = (dx^2 + dy^2)/y^2$, and $\text{Isom}^+(\mathbb{H}^2) = \text{PSL}_2(\mathbb{R})$.

Proposition 1.7. *If a surface S has $\chi(S) < 0$, then there is a discrete group $\Gamma \leq \text{PSL}_2(\mathbb{R})$ such that $S \approx \mathbb{H}^2/\Gamma$. In particular, $\pi_1(S)$ is isomorphic to Γ .*

We have

$$\pi_1(\Sigma_{g,b}) \cong F_{2g+b-1} \quad \text{and} \quad \pi_1(\Sigma_g) \cong \langle a_1, b_1, \dots, a_g, b_g \mid [a_1, b_1], \dots, [a_g, b_g] \rangle.$$

Proposition 1.8. *$\pi_1(\Sigma_g)$ is torsion free.*

Exercise 1. Prove Proposition 1.7.

Theorem 1.9 (Newman). *$\pi_1(\Sigma_{g,b})$ is linear over \mathbb{Z} , that is, is isomorphic to a subgroup of $\text{GL}_n(\mathbb{R})$. For example, we can check $F_n \hookrightarrow F_2 \hookrightarrow \text{SL}_2(\mathbb{Z})$ according to the pingpong lemma.*

Over \mathbb{R} , we may embed $\pi_1(S) \hookrightarrow \text{PSL}_2(\mathbb{Z}) \cong \text{SO}_{1,2}^+(\mathbb{R}) < \text{GL}_3(\mathbb{R})$ if $\chi(S) < 0$,

Definition. A group π is called residually finite (RF) if for every $1 \neq \gamma \in \pi$ there is a group homomorphism $\varphi : \pi \rightarrow G$ to a finite group G such that $\varphi(\gamma) \neq 1$. A subgroup $\pi' < \pi$ is called separable if there is a group homomorphism $\varphi : \pi \rightarrow G$ to a finite group G such that $\varphi(\gamma) \notin \varphi(\pi')$. In particular, π is residually finite if the trivial subgroup is separable in π . A group π is called locally extended residually finite (LERF) if every finitely generated subgroup of π is separable.

Theorem 1.10 (Mal'cev). *Every finitely generated linear group over a field is residually finite.*

Over \mathbb{Z} , if $1 \neq (a_{ij}) \in \pi < \text{GL}_n(\mathbb{Z})$ given, then for $m > \max_{i,j} |a_{ij}|$ if we let $\varphi_m : \pi \hookrightarrow \text{GL}_n(\mathbb{Z}) \rightarrow \text{GL}_n(\mathbb{Z}/m\mathbb{Z})$, then $\varphi_m((a_{ij})) \neq 1$.

Theorem 1.11 (Scott). *$\pi_1(\Sigma_{g,b})$ is LERF*

2 Day 2: April 18

2. Examples and constructions of 3-manifolds

Theorem 2.1 (Moise). *Every topological 3-manifold (not necessarily compact, connected, orientable) admits a unique smooth and PL structure.*

2.1 Spherical manifolds

Recall

$$\begin{aligned} S^3 &:= \{x \in \mathbb{R}^4 : |x| = 1\} \\ &= \{(z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 = 1\} \\ &= \{a + bi + cj + dk \in \mathbb{H} : a^2 + b^2 + c^2 + d^2 = 1\}. \end{aligned}$$

Lens spaces

Let $p, q \in \mathbb{Z}$, $p > 0$, $(p, q) = 1$. Then, $\mathbb{Z}/p\mathbb{Z} = \langle \zeta = \exp(2\pi\sqrt{-1}/p) \rangle$ acts on S^3 such that $\zeta \cdot (z, w) = (\zeta z, \zeta^q w)$. Then, the Lens spaces are defined as

$$L(p, q) := S^3 / (\mathbb{Z}/p\mathbb{Z}) \quad \text{with} \quad \pi_1(L(p, q)) = \mathbb{Z}/p\mathbb{Z}.$$

For example, $L(1, 1) = S^3$ and $L(2, 1) = \mathbb{RP}^3$.

Theorem 2.2 (Reidemeister).

- (a) $L(p, q) \simeq L(p, q')$ (homotopy equiv) if and only if there is $a \in \mathbb{Z}$ such that $qq' \equiv \pm a^2 \pmod{p}$.
- (b) $L(p, q) \approx L(p, q')$ (diffeo) if and only if $q' \equiv \pm q^{\pm 1} \pmod{p}$.

For example, $L(7, 1) \simeq L(7, 2)$ since $1 \cdot 2 \equiv 3^2 \pmod{7}$, but $L(7, 1) \not\approx L(7, 2)$ since $2 \not\equiv \pm 1 \pmod{7}$.

Sketch of (\Leftarrow). Direct construction. (a) With the linking form $H_1(L) \times H_1(L) \rightarrow \mathbb{Q}/\mathbb{Z}$. (b) Reidemeister torsion. \square

General quotients

A spherical manifold is the orbit space S^3/Γ , where Γ is a finite subgroup of $\text{SO}(4)$ and $\Gamma \curvearrowright S^3$ freely.

Example. With an action $\langle -1, i, j, k \rangle \curvearrowright S^3$, we obtain the prism manifold.

Example. With an action of the binary icosahedral group $\Gamma = \mathbb{Z}/2\mathbb{Z} \rtimes A_5$ on S^3 , we obtain the Poincaré sphere. We have $H_*(S^3/\Gamma) \cong H_*(S^3)$. If we take $3/10$ turn instead of $1/10$, we have the Seifert-Weber space.

2.2 Fibered manifolds

Twisted bundles

$$N_{g,b} = (\mathbb{RP}^2)^{\#g} \# (D^2)^{\#b}.$$

Let D be a polygon with oriented sides $a_1, a'_1, a_2, a'_2, \dots, a_g, a'_g$.

$$N_g \widetilde{\times} [0, 1] := D \times [0, 1] / \sim,$$

where $(x, t) \sim (x', 1 - t)$ for $x \in a_i$, $x' \in a'_i$, $t \in [0, 1]$ with $[x] = [x'] \in N_g$, and it is orientable.

$$N_g \widetilde{\times} S^1 := N_g \widetilde{\times} [0, 1] / (x, 0) \sim (x, 1), \quad x \in N_g$$

$$N_{g,b} \widetilde{\times} S^1 := N_g \widetilde{\times} S^1 \setminus \nu(b \text{ fibers}).$$

Exercise 2. Show the following:

- (a) $\mathbb{RP}^2 \widetilde{\times} [0, 1] \approx \mathbb{RP}^3 \setminus (\text{open ball})$.
- (b) $\mathbb{RP}^2 \widetilde{\times} S^1 \approx \mathbb{RP}^3 \# \mathbb{RP}^3$.
- (c) $N_{1,1} \widetilde{\times} S^1 \approx N_2 \widetilde{\times} [0, 1]$.

Mapping tori

The mapping class group is

$$\mathcal{M}_{g,b} := \text{Diff}^+(\Sigma_{g,b}, \partial \Sigma_{g,b}) / \text{isotopy relative to } \partial.$$

Theorem 2.3 (Dehn, Lickorish). $\mathcal{M}_{g,b}$ is finitely generated by Dehn twists.

For examples, $\mathcal{M}_0 = \mathcal{M}_{0,1} = 1$ by the Alexander trick, and $\mathcal{M}_{0,2} \cong \mathbb{Z}$, $\mathcal{M}_1 = \mathcal{M}_{1,1} \cong \text{SL}_2(\mathbb{Z})$.

Let $\varphi \in \mathcal{M}_{g,b}$. Then, a mapping torus is defined by

$$M_\varphi := \Sigma_{g,b} \times [0, 1] / (\varphi(x), 0) \sim (x, 1).$$

2.3 Heegaard decomposition

A manifold with a boundary

$$H_g := D^3 \cup (D^2 \times [0, 1])^{\cup g}$$

is called the handle body with genus g . Then, $\pi_1(H_g) \cong F_g$ and $\partial H_g \approx \Sigma_g$. Let $\varphi : \partial H_g \rightarrow \partial H_g$ be an orientation-preserving diffeomorphism, i.e. an element of the mapping class group \mathcal{M}_g . If a 3-manifold M satisfies

$$M \approx H_g \approx H_g,$$

then the right-hand side is called the Heegaard decomposition (splitting) of M .

Proposition 2.4. Every closed 3-manifold admitting a Heegaard decomposition of genus 0 is diffeomorphic to S^3 .

Proof. $\mathcal{M}_0 = 1$. □

Proposition 2.5. Every closed 3-manifold admitting a Heegaard decomposition of genus 1 is diffeomorphic to S^3 , $S^2 \times S^1$, or $L(p, q)$.

Exercise 3. Prove the above proposition.

Theorem 2.6. Every closed 3-manifold M admits a Heegaard decomposition along some Σ_g .

Proof. Pick a triangulation T of M . Then, $H := \overline{\nu(T^{(1)})}$ is a handlebody. Then, $H' := M \setminus \text{Int} H$ is also a handlebody. Since M is orientable, so are H and H' , thus we are done. □

There is another proof using Morse theory.

Corollary 2.7. The fundamental group of every closed 3-manifold admits a finite presentation of deficiency 0, i.e. the number of generators is equal to the number of relations.

Proof. Apply the van Kampen theorem to

$$M = H_g \cup H_g = H_g \cup (D^2 \times [-\varepsilon, \varepsilon])^{\cup g} \cup D^3. \quad \square$$

2.4 Dehn surgery

Let L be a link. The link exterior is the set $E_L = S^3 \setminus \nu(L)$.

Proposition 2.8. Let M be a 3-manifold, and $T \subset \partial M$ a torus component. Let $h : \partial(D^2 \times S^1) \rightarrow T$ be a diffeomorphism. Then, $M \cup_h (D^2 \times S^1)$ is determined only by $\pm[h(\partial D^2 \times *)] \in H_1(T)$.

Proof. Write

$$M \cup_h (D^2 \times S^1) = M \cup_h (D^2 \times (-\varepsilon, \varepsilon)) \cup D^3. \quad \square$$

For a knot K , there are two generators μ and λ , called the meridian and the longitude, of $H_1(\partial E_K)$ such that $\ker(H_1(\partial E_K) \rightarrow H_1(E_K) = \mathbb{Z}\mu)$ is generated by λ .

Exercise 4. Show that $L(p, q)$ with $p \neq 0$ and $S^2 \times S^1$ are obtained by the p/q -Dehn surgery along the unknot.

Theorem 2.9 (Lickorish-Wallace). *Every closed 3-manifold can be obtained by an (integral)-Dehn surgery along some link in S^3 .*

Sketch. Heegaard decomposition and $\mathcal{M}_g = \langle \text{Dehn twists} \rangle$. Each Dehn twist realizes the Dehn surgery steps. □

3 Day 3: April 25

3. Prime decomposition

3.1 Alexander's theorem

Theorem 3.1. Every (smooth) embedding $S^2 \subset \mathbb{R}^3$ bounds some (smooth) embedding $D^3 \subset \mathbb{R}^3$.

Remark. The above theorem does not hold in the category of topological spaces. Alexander's horned sphere is one of the counterexamples.

If \mathbb{R}^d for $d \geq 5$, then more complicated result such as h-cobordism theorem must be used to obtain the same conclusion.

Sketch. Isotope such a sphere S so that the coordinate $z : S \rightarrow \mathbb{R}$ is a Morse function. Assume that for all $p \neq q \in \text{Crit}(z)$, then $z(p) \neq z(q)$. We use induction on (m, n) , where m is the number of saddles and

$$n := \min\{\#\pi_0(S \cap z^{-1}(r)) : r \text{ is a regular value s.t. } z(p) < r < z(q) \text{ for some saddles } p, q\}.$$

Note that $\#(\text{minima of } z) - m + \#(\text{maxima of } z) = \chi(S) = 2$.

For the case $m = 0$ so that there are only one minimum and maximum, then we can construct a ball by applying the Jordan-Schönflies theorem to each level.

For the case $m = 1$, then only four types appear: a jelly bean, a red blood cell, and their upside down versions. Apply the Jordan Schönflies again.

For the case $m \geq 2$, let r be a regular value realizing the value of n . Let D be union of the closure of the interior of the innermost circles of $S \cap z^{-1}(r)$. Replace S by $(S \setminus \partial D \times (-\varepsilon, \varepsilon)) \cup (D \times \{-\varepsilon, \varepsilon\})$. Then, each connected component has lower (m, n) so that it bounds a ball. Attaching all balls bounded by the components with balls $D \times [-\varepsilon, \varepsilon]$, S also bounds a ball. \square

3.2 Irreducible manifolds

The connected sum is defined as

$$M \# N := (M \setminus (\text{open ball})) \cup_{\partial} (N \setminus (\text{open ball})).$$

Proposition 3.2.

- (a) $M \# N \approx N \# M$.
- (b) $(M_1 \# M_2) \# M_3 \approx M_1 \# (M_2 \# M_3)$.
- (c) $M \# S^3 \approx M$.

We say a manifold M is *prime* if $M = N_1 \# N_2$ implies $N_1 \approx S^3$ or $N_2 \approx S^3$. We say a 3-manifold M is *irreducible* if every embedding $S^2 \subset M$ bounds some embedding $D^3 \subset M$. In other words, in a prime manifold every separating sphere bounds a ball, in an irreducible manifold every sphere bounds a ball.

Corollary 3.3. Every irreducible 3-manifold is prime.

Corollary 3.4. By Theorem 3.1, S^3 is irreducible.

Theorem 3.5.

- (a) $S^2 \times S^1$ is prime, but is not irreducible.
- (b) Every closed prime 3-manifold which is not irreducible is diffeomorphic to $S^2 \times S^1$.

Proof. (a) A sphere $S^2 \times *$ cannot bound any $D^3 \subset S^2 \times S^1$ because $[S^2 \times *] \neq 0 \in H_2(S^2 \times S^1)$, so $S^2 \times S^1$ is not irreducible.

Suppose $S^2 \times S^1 = N_1 \# N_2$. Since $\pi_1(N_1) * \pi_1(N_2) \cong \mathbb{Z}$, one of $\pi_1(N_1)$ or $\pi_1(N_2)$ is trivial. Assume $\pi_1(N_1)$ is trivial and let $B := N_1 \setminus (\text{open ball})$. Since B is also simply connected, it lifts diffeomorphically into the universal cover $S^2 \times \mathbb{R}$ of $S^2 \times S^1$. Because $S^2 \times \mathbb{R} \approx \mathbb{R}^3 \setminus \{0\}$, we have an embedding $B \subset \mathbb{R}^3$. Because $\partial B \approx S^2$, by Theorem 3.1 we have $B \approx D^3$, so $N_1 \approx S^3$.

(b) If every sphere in M is separating, then it has to be irreducible since M is prime, so such M contains a nonseparating sphere S . Let γ be an arc connecting the inside and the outside of $\partial \nu(S)$. If we let $M' := \overline{\nu(S)} \cup \overline{\nu(\gamma)}$, then $\partial M' \approx S^2 \# (S^1 \times I) \# S^2 \approx S^2$ is a separating sphere. Since $M \setminus \text{Int} M' \approx D^3$ because M' is not simply connected and M is prime, M is diffeomorphic to $S^2 \times S^1$. \square

Proposition 3.6. *If a covering space \tilde{M} of M is irreducible, then so is M .*

Exercise 4. Prove Proposition 3.6.

Remark. The converse of Proposition 3.6 is also known to be true.

3.3 Normal surfaces

Fix a 3-manifold M and its triangulation T . A (possibly disconnected) subsurface $S \subset M$ is called a *normal surface* with respect to T if S is a union of *normal disks*, defined as seven types of disks in a given tetrahedron: four triangles and three quadrilaterals.

Proposition 3.7. *Every (possibly disconnected) subsurface $S \subset M$ becomes a normal surface with respect to T by isotopies and the following operations:*

- (i) *Replace S by $(S \setminus \partial D \times (-\varepsilon, \varepsilon)) \cup (D \times \{\pm \varepsilon\})$ for a disk D satisfying $D \cap (S \cup \partial M) = \partial D$.*
- (ii) *Remove a components of S contained in a ball in M .*

Proof. Isotope S so that $S \pitchfork T$. It is sufficient to realize the following:

- (a) For every tetrahedron $\Delta^{(3)} \subset T^{(3)}$, $S \cap \Delta^{(3)} = \coprod (\text{disks})$.
- (b) For every disk component D in (a) and for every edge $\Delta^{(1)} \in T^{(1)}$, $\#(D \cap \Delta^{(1)}) \leq 1$.
- (c) For every triangle $\Delta^{(2)} \subset T^{(2)}$, $S \cap \Delta^{(2)} = \coprod (\text{arcs})$.

For (a), if there is a non-disk connected component of $S \cap \Delta^{(3)}$, perform (i) along innermost loops in $S \cap \partial \Delta^{(3)}$. Then, perform (ii) for closed components of S contained in $\Delta^{(3)}$.

For (b), if there is a disk component of $S \cap \Delta^{(3)}$ which intersects an edge $\Delta^{(1)}$ more than twice, then for an inner most pair of two points connected by an arc in D , push the arc to the outside $\Delta^{(3)}$ with an ambient isotopy.

For (c)

\square

4 Day 4: May 9

3.4 Prime decomposition theorem

Proposition 3.8. *Let T be a triangulation of M and let $S \subset M$ be a normal surface with respect to T . If no component of $M \setminus \nu(S)$ is a trivial $[0, 1]$ -bundle, then*

$$\# \pi_0(S) \leq 6 \cdot \# T^{(3)} + \text{rk} H_2(M, \partial M; \mathbb{Z}/2\mathbb{Z}).$$

Proof. Let M_0 be a component of $M \setminus \nu(S)$ such that for each tetrahedron $\Delta^{(3)}$ the intersection $M_0 \cap \Delta^{(3)}$ is a trivial $[0, 1]$ -bundle over a normal disk $D_0(\Delta^{(3)}) \subset \Delta^{(3)}$. Let $S' = (S \setminus \sqcup \partial M_0) \sqcup (\sqcup D_0)$. Then,

$$\# \pi_0(S) = \# \pi_0(S') = \text{rk} H_2(S', \partial S'; \mathbb{Z}/2\mathbb{Z}),$$

and since

$$H_3(M, \partial M \cup S'; \mathbb{Z}/2\mathbb{Z}) \rightarrow H_2(S', \partial S'; \mathbb{Z}/2\mathbb{Z}) \rightarrow H_2(M, \partial M; \mathbb{Z}/2\mathbb{Z})$$

is exact, we have

$$\text{rk} H_2(S', \partial S'; \mathbb{Z}/2\mathbb{Z}) \leq \text{rk} H_3(M, \partial M \cup S'; \mathbb{Z}/2\mathbb{Z}) + \text{rk} H_2(M, \partial M; \mathbb{Z}/2\mathbb{Z}).$$

If we let $M' := (M \setminus \nu(S)) \setminus \sqcup M_0$, then by the excision theorem,

$$\text{rk} H_3(M, \partial M; \mathbb{Z}/2\mathbb{Z}) = \text{rk} H_3(M', \partial M'; \mathbb{Z}/2\mathbb{Z}) = \pi_0(M').$$

Every component of M' is not a trivial $[0, 1]$ -bundle over a normal disk in some tetrahedron in T . The number of such non-trivial components of the complement $\Delta^{(3)} \setminus \nu(S)$ is at most 6. \square

Theorem 3.9 (Prime decomposition theorem; Kneser, Milnor). *Every 3-manifold $M \neq S^3$ admits a decomposition*

$$M = N_1 \# \cdots \# N_n, \quad N_i : \text{prime} \neq S^3.$$

Moreover, such a decomposition is unique up to diffeomorphisms and permutations.

Proof. (Existence) If M contains a non-separating sphere, then there is M' such that $M = M' \# (S^2 \times S^1)$ as in the proof of Proposition 3.5. Since we have $H_1(M) = H_1(M') \oplus \mathbb{Z}$, the number of such $S^2 \times S^1$ factors is finite, so by cutting out all of them, we may assume that every sphere in M is separating.

Let S be an arbitrary union of spheres in M such that

$$\text{no components of } M \setminus \nu(S) \text{ is diffeomorphic to } S^3 \setminus (\text{balls}). \quad (\dagger)$$

It suffices to show that $\pi_0(S)$ is finite. Let T be a triangulation of M and $S' \subset M$ be a normal surface with respect to T obtained from S as Proposition 3.7. Suppose that a component $S_0 \subset S$ is contained in a ball. Then by the Alexander theorem such an innermost component would bound a ball, which contradicts to (\dagger) , hence the operation (ii) in Proposition 3.7 does not occur when we construct S' . Each operation (i) splits a component $S_0 \subset S$ into two spheres S_1 and S_2 , and one of them satisfies (\dagger) . Therefore, there exists a union of components $S'' \subset S'$ satisfying the same condition with S with $\pi_0(S'') = \pi_0(S)$, which is bounded by Proposition 3.8.

(Uniqueness) Suppose

$$P_1 \# \cdots \# P_m \# (S^2 \times S^1)^{\#k} = Q_1 \# \cdots \# Q_n \# (S^2 \times S^1)^{\#l},$$

where P_i and Q_j are irreducible but not S^3 . It suffices to find a union of spheres $T \subset M$ such that

$$(P) \quad M \setminus \nu(T) = \bigsqcup_{i=1}^m (P_i \setminus (\text{balls})) \sqcup (\bigsqcup S^3 \setminus (\text{balls})),$$

$$(Q) \quad M \setminus \nu(T) = \bigsqcup_{j=1}^n (Q_j \setminus (\text{balls})) \sqcup (\bigsqcup S^3 \setminus (\text{balls})).$$

Pick unions of spheres $R, S \subset M$ such that R satisfies (P), S satisfies (Q), and $R \pitchfork S$. If $R \cap S = \emptyset$, then let $T := R \sqcup S$. If not, pick components $R_0 \subset R$ and $S_0 \subset S$ such that $R_0 \cap S_0 \neq \emptyset$. Split R_0 along an innermost disk in S_0 into two spheres R_1 and R_2 , and replace R by $R' := (R \setminus R_0) \sqcup R_1 \sqcup R_2$. Then, R' satisfies (P) and $\# \pi_0(R' \cap S) < \# \pi_0(R \cap S)$ so that such operations reduce to the case of $R \cap S = \emptyset$. \square

Exercise 5. Show that the existence of the prime decomposition theorem follows from the Poincaré conjecture and Grushko's theorem: $\text{rk } G * H = \text{rk } G + \text{rk } H$, where the rank is defined to be the minimum number of generators.

Theorem 3.10 (Kneser conjecture, proved by Stallings). *Let M be an oriented 3-manifold such that $\pi_1(M) = G_1 * G_2$. Then, there are oriented 3-manifolds N_1 and N_2 such that $M = N_1 \# N_2$ and $\pi_1(N_i) \cong G_i$ for $i \in \{1, 2\}$.*

For this theorem, it suffices to study irreducible 3-manifolds.

4. Loop and sphere theorems

4.1 Loop theorem

Theorem 4.1 (Loop theorem). *For a subsurface $S \subset M$, there is a continuous embedding $f : (D^2, \partial D^2) \rightarrow (M, S)$ such that $f|_{\partial D^2} \neq \text{null homotopic in } S$. In particular, we may assume f is smooth.*

Remark. The loop theorem holds also for noncompact or nonorientable 3-manifolds.

Corollary 4.2 (Dehn's lemma). *Every loop in ∂M null homotopic in M bounds some embedded disk in M .*

Proof. Apply the loop theorem for the tubular neighborhood of the loop in ∂M . \square

Theorem 4.3. *Let $K \subset S^3$ be a knot and $E_K := S^3 \setminus \nu(K)$. Then, K is unknot if and only if $\pi_1(E_K) \cong \mathbb{Z}$.*

Proof. (\Rightarrow) Clear.

(\Leftarrow) Take a longitude $\lambda \in \pi(\partial E_K) = H_1(\partial E_K)$ such that

$$\lambda \in \ker(\pi_1(\partial E_K) \rightarrow H_1(E_K)) \subset \iota_*^{-1}([\pi_1(E_K), \pi_1(E_K)]),$$

where $\iota : \partial E_K \hookrightarrow E_K$. Then, $\pi_1(E_K) \cong \mathbb{Z}$ implies $\lambda = 1$, and apply Dehn's lemma. \square

Exercise 6. Let M be an irreducible 3-manifold and $T \subset M$ a torus. Show that the followings are equivalent:

- (i) $\pi_1(T) \rightarrow \pi_1(M)$ is not injective.
- (ii) T bounds some embedded solid torus $D^2 \times S^1$ in M or T is contained in some embedded ball in M .


5 Day 5: May 16

Theorem 4.1 (Loop theorem, Papakyriakopoulos). *For a subsurface S of 3-manifold M , if there is a continuous map $f : (D^2, \partial D^2) \rightarrow (M, S)$ such that $f|_{\partial D^2}$ is not null-homotopic, then there is a smooth embedding $g : (D^2, \partial D^2) \rightarrow (M, S)$ such that $g|_{\partial D^2}$ is not null-homotopic.*

Proof. Step 1: *Tower construction.* Pick triangulations of (M, S) and D^2 .

Let $f_0 : (D^2, \partial D^2) \rightarrow (M, S)$ be a simplicial map such that $f_0 \simeq f$. Define $V_0 \subset M$ to be a 3-submanifold deformation retracting to $f_0(D^2)$ and $S_0 \subset \partial V_0$ a component of $S \cap V_0$ containing $f_0(\partial D^2)$.

Given the i -th level, for any double covering $p_{i+1} : M_{i+1} \rightarrow V_i$, if it exists, let $f_{i+1} : (D^2, \partial D^2) \rightarrow (M_{i+1}, p_{i+1}^{-1}(S_i))$ be a lift of f_i . Similarly as the case $i+1 = 0$, define $V_{i+1} \subset M_{i+1}$ to be a 3-manifold deformation retracting to $f_{i+1}(D^2)$ and $S_{i+1} \subset \partial V_{i+1}$ the component of $p_{i+1}^{-1}(S_i) \cap V_{i+1}$ containing $f_{i+1}(\partial D^2)$.



Since

$$\# \text{ of simplices of } f_0(D^2) < \# \text{ of simplices of } f_1(D^2) < \dots \leq \# \text{ of simplices of } D^2,$$

there is n such that V_n has no double cover. If we let $N_i := \ker(\pi_1(S_i) \rightarrow \pi_1(S))$, then $f_{i*}([\partial D^2]) \notin N_i$ implies that $N_i \neq \pi_1(S_i)$ for each i .

Step 2: *Embedding on the top.* Consider the following exact sequence:

$$H_2(V_n, \partial V_n; \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(\partial V_n; \mathbb{Z}/2\mathbb{Z}) \rightarrow H_1(V_n; \mathbb{Z}/2\mathbb{Z}).$$

Since V_n has no double cover, the Poincaré duality writes

$$H_2(V_n, \partial V_n; \mathbb{Z}/2\mathbb{Z}) \cong H^1(V_n; \mathbb{Z}/2\mathbb{Z}) \cong \text{Hom}(\pi_1(V_n), \mathbb{Z}/2\mathbb{Z}) = 0.$$

Also we have

$$H_1(V_n; \mathbb{Z}/2\mathbb{Z}) \cong H_1(V_n) \otimes \mathbb{Z}/2\mathbb{Z} = 0,$$

which implies $H_1(\partial V_n; \mathbb{Z}/2\mathbb{Z}) = 0$ so that every component of ∂V_n is a sphere.

Note that $S_n \approx S^2 \setminus (\text{disks})$ and $\pi_1(S_n)$ is generated by components of ∂S_n . Since $N_n \neq \pi_1(S_n)$, there is a component $l \subset \partial S_n$ such that $[l] \notin N_n$. Pick

Step 3: *Descending the tower.* We shall show that given an embedding $g_i : (D^2, \partial D^2) \rightarrow (V_i, S_i)$ such that $g_{i*}([\partial D^2]) \notin N_i$, there is such an embedding g_{i-1} . Then, g_0 is the desired embedding.

Isotope $p_i \circ g_i$ so that self-intersections are simple double curves. These are modified by the following operations which give g_{i-1} .

Case 2 double arcs These are cancelled in two ways and at least one of the results satisfies $g_{i-1}([\partial D^2]) \notin N_{i-1}$. \square

Let X be a path-connected space and $x_0 \in X$. Then,

$$\pi_n(X) := \pi_n(X, x_0) = \{f : (S^n, *) \rightarrow (X, x_0)\} / \text{homotopy fixing base points}.$$

For a covering $p : \tilde{X} \rightarrow X$, the induced homomorphism $p_* : \pi_n(\tilde{X}) \rightarrow \pi_n(X)$ is an isomorphism for $n \geq 2$. We will use the following two theorems without proofs

Theorem (Whitehead theorem). Let X, Y be CW-complexes. If $\varphi : X \rightarrow Y$ satisfies $\varphi_* : \pi_n(X) \rightarrow \pi_n(Y)$ is an isomorphism for each n , then φ is a homotopy equivalence.

Theorem (Hurewicz theorem). For, $h_n : \pi_n(X) \rightarrow H_n(X) : [f] \mapsto f_*([S_n])$, h_1 is the abelianization, and h_{n+1} is an isomorphism if $\pi_i(X) = 0$ for $1 \leq i \leq n$.

Theorem 4.4 (Sphere theorem, Papakyriakopoulos). For a 3-manifold M with $\pi_2(M) \neq 0$, there exists an embedding $f : S^2 \rightarrow M$ such that $[f] \neq 0$.

Corollary 4.5. If M is irreducible, then $\pi_2(M) = 0$.

Theorem 4.6. Let M be an irreducible 3-manifold with infinite fundamental group.

- (a) M is aspherical, i.e. $\pi_n(M) = 0$ for $n \geq 2$.
- (b) $\pi_1(M)$ is torsion-free.

It is well-known that a manifold M is aspherical if and only if $M \simeq K(\pi_1(M), 1)$.

Lemma 4.7. Let $m \geq 2$. Just use the identity $H_*(\pi) = H_*(K(\pi, 1))$ as the definition of group homology. Then,

$$H_n(\mathbb{Z}/m\mathbb{Z}) = \begin{cases} \mathbb{Z} & n = 0, \\ \mathbb{Z}/m\mathbb{Z} & n : \text{odd}, \\ 0 & \text{otherwise.} \end{cases}$$

In fact, $K(\mathbb{Z}/m\mathbb{Z}, 1) \simeq S^\infty/(\mathbb{Z}/m\mathbb{Z})$.

Proof of Theorem 4.6. (a) By Corollary 4.5, we have $\pi_2(M) = 0$. If we let \tilde{M} be the universal cover of M , which is not compact because $\pi_1(M)$ is infinite, so that we have

$$\pi_3(M) = \pi_3(\tilde{M}) = H_3(\tilde{M}) = 0.$$

Inductively, and since $\dim M = 3$, we have for $n \geq 4$ that

$$\pi_n(M) = \pi_n(\tilde{M}) = H_n(\tilde{M}) = 0.$$

(b) Suppose that there is a non-trivial finite cyclic subgroup $C \leq \pi_1(M)$. If we let \hat{M} be the cover of M corresponding to C , then $\pi_1(\hat{M}) = C$. It follows from (i) that $\hat{M} = K(C, 1)$. By Lemma 4.7, we have $H_n(\hat{M}) \neq 0$ for odd n , but $\dim \hat{M} = 3$ implies that $H_n(\hat{M}) = 0$ for $n \geq 4$. \square

Theorem 4.8. The only abelian groups appearing as the fundamental group of closed (orientable) 3-manifolds are $1, \mathbb{Z}, \mathbb{Z}^3, \mathbb{Z}/n\mathbb{Z}$.

Remark. If we also consider non-orientable 3-manifolds, then $\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ also appears as $\pi_1(\mathbb{RP}^2 \times S^1)$.

Exercise 7. Show that the only abelian groups admitting presentations of deficiency zero are

$$1, \mathbb{Z}, \mathbb{Z}^2, \mathbb{Z}^3, \mathbb{Z}/n\mathbb{Z}, \mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}.$$

(Hint: use Hopf's formula $H_2(\pi) = R \cap [F, F]/[F, R]$, where $\pi = F/R$.)

Proof of Theorem 4.8. By Corollary 2.7 and Exercise 7, it suffices to show $\pi_1(M) \not\cong \mathbb{Z}^2, \mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$. By the prime decomposition theorem, we may assume that M is irreducible. By the part (b) of Theorem 4.6, $\pi_1(M) \not\cong \mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$. Now suppose that $\pi_1(M) \cong \mathbb{Z}^2$.

Since M is aspherical by Theorem 4.6, $M \cong K(\mathbb{Z}^2, 1) \simeq \mathbb{T}^2$. However, $\mathbb{Z} = H^3(M) \cong H^3(\mathbb{T}^2) = 0$ leads to a contradiction. \square

6 Day 6: May 23

5. Haken manifolds

In the following, we only consider *irreducible* 3-manifolds, i.e. every embedded sphere bounds a ball.

5.1 Essential surfaces

When we say subsurfaces S , we always assume $S \cap \partial M = \partial S$.

A subsurface $S \subset M$ which is not a sphere is called *incompressible* if there is no (non-trivial) compressing disk. A *compressing disk* of a subsurface S is a disk $D \subset M$ such that $D \cap S = \partial D$ and ∂D bounds no disk in S . The second condition intuitively says that the compression does not generate spheres so that the disk compresses S non-trivially.

A subsurface $S \subset M$ which is not a sphere is called *boundary incompressible* if there is no (non-trivial) boundary compressing disks. A *boundary compressing disk* of S is a disk D such that arcs $\gamma = D \cap S$ and $\gamma' = D \cap \partial M$ form the boundary $\partial D = \gamma \cup \gamma'$ and $\gamma \cup \gamma''$ does not bound a disk in S for any arc $\gamma'' \subset \partial S$.

A surface S is called *boundary parallel* if S can be isotoped into ∂M .

A (possibly disconnected) subsurface $S \subset M$ is called an *essential surface* if every component S_0 of S is incompressible, boundary incompressible, and not boundary parallel. (S_0 must not be a sphere)

Proposition 5.1. *For a subsurface $S \subset M$, S is incompressible if and only if $\pi_1(S) \rightarrow \pi_1(M)$ is injective.*

Proof. The loop theorem. □

Proposition 5.2. *For nonzero $\alpha \in H_2(M, \partial M)$, there is an essential surface $S \subset M$ with $[S] = \alpha$.*

Let S be a possibly disconnected subsurface of M . We define a complexity $c(S) := \sum_i (2 - \chi(S_i))^2$, where S_i are components of S .

Lemma 5.3. *If S' is obtained from S by the surgery along (boundary) compressing disks, then $c(S') < c(S)$.*

Proof. We may assume S is connected. Note that every component has $\chi < 2$ since we are not allowing spheres after surgery. If S' is connected, then since $\chi(S') = \chi(S) + 2$, we are done. If $S' = S_1 \sqcup S_2$, then since $\chi(S_1) + \chi(S_2) = \chi(S') = \chi(S) + 2$, we are done. □

Proof of Proposition 5.2. Pick a (possibly disconnected) subsurface $S_0 \subset M$ with $[S_0] = \alpha$. (Choose $f : M \rightarrow S^1$ so that $[f] \in [M, S^1] = H^1(M)$ is equal to the Poincaré dual of α and let $S_0 := f^{-1}(\text{a regular value})$) Perform the surgery S_0 along (boundary) compressing disks until every components of S_0 becomes (boundary) incompressible. By Lemma 5.3, this operation stops, and before and after of the compression are homologous. Removing all spheres and boundary parallel components of S_0 , now we have a desired subsurface S with $[S] = [S_0] = \alpha$. Since $\alpha \neq 0$, $S \neq \emptyset$. □

A 3-manifold is a *Haken manifold* if it admits an essential surface in M .

Corollary 5.4. *If $b_1(M) > 0$, then M is Haken.*

Proof. It follows from Proposition 5.2. □

Corollary 5.5. *If there is a non-sphere component of ∂M , then M is Haken.*

Proof. Since $H_1(\partial M)$ contains an element of infinite order, the exact sequence

$$H^1(M) \cong H_2(M, \partial M) \rightarrow H_1(\partial M) \rightarrow H_1(M)$$

implies $b_1(M) > 0$. □

Theorem 5.6 (Thurston). *For the figure eight knot, the Dehn filling by a rational number $r = p/q$ is Haken if and only if $r = 0, \pm 4$.*

5.2 Haken hierarchy

Proposition 5.7. *Let $S \subset M$ be an essential surface. Let $M' = M \setminus \nu(S)$. Then, (a) M' is irreducible and (b) every closed incompressible surface in M' is incompressible also in M .*

Proof. (a) Let $S' \subset M'$ be arbitrary sphere. Since M is irreducible, S' bounds a ball $B \subset M$. Then, we have $S \subset B$ or $S \cap B = \emptyset$. Since B contains no closed incompressible surface, $B \cap S = \emptyset$, so $B \subset M'$.

(b) Suppose that there is a closed incompressible surface $S' \subset M'$ not incompressible in M . Let $D \subset M$ be a compressing disk for S' such that $D \cap S$ and $\# \pi_0(D \cap S)$ is minimal. Since S is essential, an innermost loop in $S \cap D \subset S$ bounds a disk $D' \subset S$ (if we take any compressible disk of S in M whose boundary is $S \cap D$, then the assumption S is essential implies that the compression gives a sphere; we can take a disk D'). Since M is irreducible, $D \cup D'$ bounds a ball in M , so D can be isotoped such that $\# \pi_0(D \cap S)$ would reduce, which is a contradiction. \square

Proposition 5.8. *If $\partial M \neq \emptyset$ and M contains no closed incompressible surface, then M is a handlebody.*

Proof. Cut M along essential disks into $\coprod_i M_i$ until every component of ∂M_i is incompressible. Suppose that there is i_0 such that $M_{i_0} \not\approx D^3$. By Proposition 5.7(a), every component of ∂M_{i_0} is not a sphere. By Proposition 5.7(b), the components of ∂M_{i_0} are closed incompressible surfaces in M which contradicts the assumption. Therefore, for every i we have $M_i \approx D^3$, M is a handlebody. \square

Theorem 5.9 (Haken hierarchy). *For every Haken manifold M , there is a sequence of irreducible (possibly disconnected) 3-manifolds $M = M_0, M_1, \dots, M_n$ such that*

- (a) $M_{i+1} = M_i \setminus \nu(S_i)$ for some essential surface $S_i \subset M_i$
- (b) $M_n \approx \coprod D^3$.

The strongest feature of the Haken hierarchy is the possibility of using induction.

Theorem 5.10 (Waldhausen). *Two closed Haken manifolds with isomorphic fundamental groups are diffeomorphic.*

Proof. Induction on the Haken hierarchy. \square

Remark. When $\partial M \neq \emptyset$, the theorem also holds under a certain additional condition on isomorphisms on π_1 .

7 Day 7: June 6

Exercise 8. Show that every connected essential surface in $\Sigma_g \times [0, 1]$ ($g \geq 1$) is isotopic to an annulus $\gamma \times [0, 1]$ for some simple closed curve γ bounding no disk in Σ_g .

Lemma 5.11. Let M be an irreducible 3-manifold. Let $\partial M = \coprod_i R_i$, where R_i are components which are not diffeomorphic to sphere. Then, there is $\alpha \in H_2(M, \partial M)$ such that for every i we have $\partial \alpha|_{R_i} \neq 0 \in H_1(R_i)$.

Exercise 9. Prove Lemma 5.11.

Proof of Theorem 5.9. Note that the theorem holds for handlebodies H_g . We may assume $M \not\approx H_g$. By Proposition 5.8, M contains a union of closed incompressible surfaces S . By Proposition 3.7 and 3.8, S is isotopic to a normal surface with respect to some triangulation of M , and if no component of $M \setminus \nu(S)$ is a trivial $[0, 1]$ -bundle, then $\pi_0(S)$ is bounded.

Take $S_0 \subset M$ with maximal fundamental group. Let $M_1 = M \setminus \nu(S_0)$. Let R_i are components of ∂M_1 such that $R_i \not\approx S^2$. By Corollary 5.5 and Proposition 5.7(a), M_1 is Haken. By Lemma 5.11, pick $\alpha \in H_2(M, \partial M)$ such that $\partial \alpha|_{R_i} \neq 0 \in H_1(R_i)$. By Proposition 5.2, pick an essential surface $S_1 \subset M$ with $[S_1] = \alpha$. Since $\pi_0(S_0)$ is maximal, there is a component M' of $M \setminus (S_0 \sqcup S')$ is a trivial $[0, 1]$ -bundle. Since $\partial \alpha|_{R_i} = [S_1 \cap R_i] \neq 0$, some component of S_1 intersects only one component of $\partial M'$. However, there is not such an essential in $M' \approx S' \times [0, 1]$. Then, contradiction (Exercise 8). \square

6. Seifert manifolds

6.1 Seifert fibrations

Let S be a (possibly non-orientable) surface. Let $p_i, q_i \in \mathbb{Z}$, $p_i \geq 1$, $(p_i, q_i) = 1$, $1 \leq i \leq k$.

If orientable, then $\times^{(\sim)} = \times$, and if non-orientable, then $\times^{(\sim)} = \tilde{\times}$, the twisted direct product.

$$M(S; (p_1, q_1), \dots, (p_k, q_k)) := \left((S \setminus \coprod_{i=1}^k D_i) \times^{(\sim)} S^1 \right) \cup_{h_1, \dots, h_k} \coprod_{i=1}^k (D^2 \times S^1),$$

where $h_i : \partial(D^2 \times S^1) \rightarrow \partial(D_i \times S^1)$ is a diffeomorphism such that

$$[h(\partial D^2 \times \{*\})] = p_i[\partial D_i \times \{*\}] + q_i[\{*\} \times S^1] \in H_1(\partial D_i \times S^1).$$

For example,

- $M(S; (1, 0)) \approx S \times^{(\sim)} S^1$.
- $M(D^2; (p, q)) \approx D^2 \times S^1$.
- $M(S^2; (p, q)) \approx L(q, p)$.
- $M(S^2; (p_1, q_1), (p_2, q_2)) \approx L(p_1 q_2 + p_2 q_1, q_2 r + p_2 s)$, where $p_1 s - q_r = \pm 1$.
- $M(D^2; (p, q)) = D^2 \times [0, 1] / (r, \theta, 1) \sim (r, \theta + \frac{2\pi q}{p}, 0)$.

Note that we have $M(D^2; (p, q)) \rightarrow D^2 / (r, \theta) \sim (r, \theta + \frac{2\pi n}{p})$ for $n \in \mathbb{Z}$. As a generalization of this, we have the Seifert fibration

$$M(S; (p_1, q_1), \dots, (p_k, q_k)) \rightarrow S(p_1, \dots, p_k),$$

where $S(p_1, \dots, p_k)$ is an orbifold constructed from S with cone points of order p_1, \dots, p_k . A *singular fiber* is the fiber of a cone point of order ≥ 2 .

Remark. Every Seifert fibration without any singular fiber is a S^1 -bundle over S .

Proposition 6.1. Every S^1 -bundle over S with orientable total space M is a Seifert fibration.

Proof. If $\partial S \neq \emptyset$, then $M \cong S \times^{(\sim)} S^1$. If $\partial S = \emptyset$, then $M \setminus \nu(\text{fiber}) \cong (S \setminus (\text{open disk})) \times^{(\sim)} S^1$. Thus there is $e \in \mathbb{Z}$ such that $M \cong M(S; (1, e))$. \square

Exercise 10. Show that if $\partial S = \emptyset$ and $e, e' \in \mathbb{Z}$, then $M(S; (1, e)) \cong M(S; (1, e'))$ if and only if $e = \pm e'$.

Let $\pi : M(S; (p_1, q_1), \dots, (p_k, q_k)) \rightarrow S(p_1, \dots, p_k)$ be a Seifert fibration. The *Euler number* of π is

$$e(\pi) := \sum_{i=1}^k \frac{q_i}{p_i},$$

which is set to sit in \mathbb{Q} if $\partial S = \emptyset$, and in \mathbb{Q}/\mathbb{Z} if $\partial S \neq \emptyset$.

Theorem 6.2. *Two Seifert fibrations*

$$\pi : M(S; (p_1, q_1), \dots, (p_k, q_k)) \rightarrow S(p_1, \dots, p_k)$$

and

$$\pi' : M(S; (p_1, q'_1), \dots, (p_k, q'_k)) \rightarrow S(p_1, \dots, p_k)$$

with $p_i \geq 2$ for all i are isomorphic by an orientation preserving diffeomorphism if and only if the following two conditions hold:

- (i) $q_i \equiv q'_i \pmod{p_i}$ for all i ,
- (ii) $e(\pi) = e(\pi')$.

Remark. We write $M(S; (p_1, -q_1), \dots, (p_k, -q_k)) = -M(S; (p_1, q_1), \dots, (p_k, q_k))$.

This theorem follows from the following proposition.

Proposition 6.3. *Isomorphism by an orientation preserving diffeomorphism is obtained by a finite number of the following moves:*

- (i) $(p_i, q_i), (p_j, q_j) \leftrightarrow (p_i, q_i + p_i), (p_j, q_j - p_j)$,
- (ii) $(p_i, q_i) \leftrightarrow (p_i, q_i \pm p_i)$ if $\partial S \neq \emptyset$.
- (iii) $(1, 0) \leftrightarrow \emptyset$.

Proof. First we prove the moves generate orientation preserving diffeomorphisms. Let $A \subset M$ be a fibered annulus connecting $M(D^2; (p_i, q_i))$ and $M(D^2; (p_j, q_j))$ (or resp. ∂M). Then, moves (i) (and resp. (ii)) correspond to the 3-dimensional version of twist along A . The move (iii) inserts or eliminates $M(D^2, (1, 0)) \approx D^2 \times S^1$.

Now we show the converse. First we can reduce the regular fibers $p_i = 1$ with three kinds of moves.

Case (a): no singular fiber. If $\partial S = \emptyset$, then we have $M(S; (1, e))$ with the same e (Exercise 10). Otherwise $\partial S \neq \emptyset$, then we have $S \times^{(\sim)} S^1$.

Case (b): singular fiber. We have $p_i \geq 2$ for all i . Let $\gamma_1, \dots, \gamma_n \subset S$ disjoint arcs connecting cone points or ∂S , cutting S into a disk D . Note that

$$M(S; (p_1, q_1), \dots, (p_k, q_k)) = \coprod_{i=1}^k M(D^2; (p_i, q_i)) \cup \coprod_{j=1}^n \overline{\nu(A_j)} \cup D^2 \times S^1,$$

where $A_j := \gamma_j \times S^1$. Every isomorphism by an orientation preserving diffeomorphism is determined by mapping classes on $\overline{\nu(A_j)}$ since it sends singular fibers to themselves. Such a mapping class is some product of the twist along A_j , which corresponds the moves (i) or (ii). \square

8 Day 8: June 13

6.2 Classifications

Let O and \tilde{O} be 2-orbifolds with cone points. A map $p : \tilde{O} \rightarrow O$ is called a *covering* if for each $x \in O$ has an open neighborhood U such that $p^{-1}(U) = \coprod_i U_i$, where $p|_{U_i} : U_i \rightarrow U$ is diffeomorphically interpreted as $\mathbb{R}^2/(\mathbb{Z}/m\mathbb{Z}) \rightarrow \mathbb{R}^2/(\mathbb{Z}/mn\mathbb{Z})$ for some $m, n \geq 1$.

The Euler characteristic is defined by

$$\chi(S(p_1, \dots, p_k)) := \chi(S) - \sum_{i=1}^k \left(1 - \frac{1}{p_i}\right) \in \mathbb{Q}.$$

Then we have $\chi(\tilde{O}) = \deg p \chi(O)$, where $\deg p = \#p^{-1}(x)$ for a non-singular $x \in O$.

Lemma 6.4. *Every 2-orbifold except $S^2(p)$ and $S^2(p_1, p_2)$ with $p_1 \neq p_2$ is finitely covered by some (usual) surface.*

Proposition 6.5. *Every Seifert manifold M is finitely covered by S^3 , $S^2 \times S^1$, S^1 -bundles over Σ_g ($g \geq 1$), or $\Sigma_{g,b} \times S^1$ ($b \geq 1$).*

Proof. Let $M \rightarrow S$ be a Seifert fibration. If $S = S^2(p)$ or $S^2(p_1, p_2)$, then M is a lens space or diffeomorphic to $S^2 \times S^1$. Otherwise, by Lemma 6.4, S is finitely covered by some surface S' . Pulling back M along $S' \rightarrow S$, we have a S^1 -bundle $M' \rightarrow S'$. If $S' = S^2$, then M' is a lens space or diffeomorphic to $S^2 \times S^1$. Otherwise, M' is a S^1 -bundle over Σ_g with $g \geq 1$, or diffeomorphic to $\Sigma_{g,b} \tilde{\times} S^1$ with $b \neq 0$. \square

Proposition 6.6. *Every Seifert manifold finitely covered by $S^2 \times S^1$ is diffeomorphic to $S^2 \times S^1$ or $\mathbb{RP}^2 \tilde{\times} S^1$.*

Exercise 11. Prove Proposition 6.6.

Theorem 6.7. *Every Seifert manifold not diffeomorphic to $S^2 \times S^1$ nor $\mathbb{RP}^2 \tilde{\times} S^1$ is irreducible.*

Proof. By Proposition 6.5 and 6.6, the interior of such a 3-manifold has a universal cover diffeomorphic to S^3 or \mathbb{R}^3 . Therefore, the theorem follows from Theorem 3.1 and Proposition 3.6. \square

A subsurface Σ of a Seifert manifold M is called *horizontal* if every fiber transversally intersects Σ , and *vertical* if it is a union of regular fibers(?). Then, every horizontal subsurface is a finite cover of the base orbifold, and every vertical subsurface is diffeomorphic either to a torus or an annulus.

Proposition 6.8. *Every connected essential surface in an irreducible Seifert manifold is isotopic to a horizontal or vertical subsurface.*

Proof. Let $M \rightarrow S$ be a Seifert fibration with an irreducible M . Let C_1, \dots, C_k be singular fibers. If there is no, pick one regular fiber as C_1 . Let $\gamma_1, \dots, \gamma_n \subset S$ be distinct arcs connecting cone points or ∂S , and cutting S into a disk D . Let $A_j := \gamma_j \times S^1$ be annuli, then we have

$$M = \coprod_{i=1}^k \overline{\nu(C_i)} \cup \coprod_{j=1}^n \overline{\nu(A_j)} \cup (D \times S^1).$$

Let Σ be a connected essential surface and isotope Σ so that

- for each i we have $\Sigma \cap \overline{\nu(C_i)} = \coprod_p (\text{disk}) \times \{p\}$, $p \in C_i$,
- for each j we have $\Sigma \cap A_j$,
- among these satisfying the above two conditions, $(\sum_{i=1}^k \# \Sigma \cap C_i, \sum_{j=1}^n \# \pi_0(\Sigma \cap A_j))$ is minimal.

It suffices to show the following:

- (a) Every component of $\Sigma \cap D \times S^1$ is isotopic to a horizontal disk or a vertical annulus.
- (b) Every component of $\Sigma \cap A_j$ is a horizontal arc or a vertical loop.

(a) We can check that the desired statement holds if and only if every component of $\Sigma \cap D \times S^1$ is incompressible in $D \times S^1$. Suppose there is a compressing disk $D' \subset D \times S^1$ for $\Sigma \cap D \times S^1$. Since Σ is essential in M , $\partial D'$ bounds a disk $D'' \subset \Sigma$. Since M is irreducible, the sphere $D' \cup D''$ bounds some ball in M . We can isotope such a ball and it would reduce $\sum_{i=1}^k \# \Sigma \cap C_i$ or $\sum_{j=1}^n \# \pi_0(\Sigma \cap A_j)$, which is a contradiction, hence every component of $\Sigma \cap D \times S^1$ is incompressible.

(b) The same argument shows that for each j there is no loop in $\Sigma \cap A_j$ bounding a disk in A_j . If there is an arc γ in $\Sigma \cap A_j$ bounding a disk, the since Σ is boundary incompressible, $\partial \gamma$ is adjacent to some C_i . An isotopy pushing the arc toward C_i along such a disk would reduce $\sum_{i=1}^k \# \Sigma \cap C_i$, which is a contradiction, so the conclusion follows. \square

Proposition 6.9. *Let M is an irreducible Seifert manifold.*

- (a) *Every horizontal subsurface in M is essential.*
- (b) *A vertical subsurface in M is essential if its projection in the base orbifold is not the following:*
 - (i) *a loop bounding a disk with at most one cone point,*
 - (ii) *a boundary parallel curve,*

Sketch. (a) Let $\Sigma \subset M$ be a horizontal subsurface. It is easy to see that Σ is not boundary parallel. Since Σ is a finite cover of the base manifold S , then we have $\pi_1(\Sigma) \hookrightarrow \pi_1(S) \hookrightarrow \pi_1(M)$, so Σ is incompressible in M . The same argument shows that $\Sigma \cup_{\partial} \Sigma$ in a closed manifold $M \cup_{\partial} M$ is incompressible, which implies Σ is boundary incompressible.

(b) Let $\Sigma \subset M$ be a vertical subsurface satisfying the conditions. By the assumption (ii), Σ is not boundary parallel. Components of $M \setminus \nu(\Sigma)$ are Seifert manifolds, suppose there is an essential disk in $M \setminus \nu(\Sigma)$. By Proposition 6.8, $M \setminus \nu(\Sigma)$ contains a horizontal disk D , and D is a finite cover of the base orbifold, which is diffeomorphic to D^2 or $D^2(p)$. \square

Theorem 6.10. *Two Seifert manifolds are diffeomorphic if and only if their Seifert fibrations are isomorphic, except the following:*

- (i) $M(D^2; (p, q)) \approx D^2 \times S^1$.
- (ii) $M(D^2; (2, 1), (2, 1)) \approx N_{1,1} \tilde{\times} S^1$.
- (iii) $M(S^2; (p_1, q_1), (p_2, q_2)) \approx S^3, S^2 \times S^1$, or $L(p, q)$.
- (iv) $M(S^2; (p, q), (2, 1), (2, -1)) \approx M(\mathbb{RP}^2; (q, p))$.
- (v) $M(S^2; (2, 1), (2, 1), (2, -1), (2, -1)) \approx N_2 \tilde{\times} S^1$.

Theorem 6.11. *Two closed Seifert manifold with isomorphic fundamental groups are diffeomorphic or both are lens spaces.*

Theorem 6.12 (Casson-Junreis, Gabai). *For an irreducible manifold M with infinite $\pi_1(M)$. Then, M is Seifert if and only if $\pi_1(M)$ has a infinite cyclic normal subgroup.*

9 Day 9: June 20

7. JSJ decomposition

Today, we assume that M is an irreducible 3-manifold with $\partial M = \emptyset$ or $\coprod T^2$. We say M is *atoroidal* if there is no essential torus in M .

Theorem 7.1 (JSJ(Jaco-Shalen, Johanson) decomposition). *There is a (possibly empty) essential surface $S \subset M$ consisting of tori such that every component of $M \setminus \nu(S)$ is atoroidal or a Seifert manifold. Moreover, such S with minimal $\# \pi_0(S)$ is unique up to isotopy.*

7.1 Existence

Proposition 7.2. *There is a (possibly empty) essential surface $S \subset M$ consisting of tori such that every component of $M \setminus \nu(S)$ is atoroidal.*

Proof. If $S' \subset M$ is an essential surface consisting of tori such that no component of $M \setminus \nu(S')$ is a trivial $[0, 1]$ -bundle, then by Proposition 3.7, S' is isotopic to a normal surface with respect to some triangulation of M , and by Proposition 3.8, $\# \pi_0(S')$ is bounded.

Take $S \subset M$ to be such a surface S' conditioned as above with maximal $\# \pi_0(S)$. Suppose that there is a component of $M \setminus \nu(S)$ contains an essential torus T . Then, no component of $M \setminus \nu(S \cup T)$ is a trivial $[0, 1]$ -bundle and $\#(S \cup T) = \# \pi_0(S) + 1$, which is a contradiction. \square

7.2 Uniqueness

Lemma 7.3. *Every atoroidal manifold M containing an essential annulus is a Seifert manifold.*

Lemma 7.4. *Every annulus in a Seifert manifold M is isotopic to a vertical subsurface, if necessary after we change Seifert fibrations.*

Proof. Take $S \subset M$ to be such a surface S' condition as in the proof of existence, with maximal $\# \pi_0(S)$. Suppose that there is a component of $M \setminus \nu(S)$ contains an essential torus T . Then, no component of $M \setminus \nu(S \cap T)$ is a trivial $[0, 1]$ -bundle and $\# \pi_0(S \cup T) = \# \pi_0(S) + 1$. It is a contradiction. \square

Proof of uniqueness. Let $S, S' \subset M$ be such essential surfaces consisting of tori. Isotope S and S' so that they are transversal and $\#(S \cap S')$ is minimal. In the following we show:

- (a) $S \cap S' = \emptyset$,
- (b) there is a pair of parallel components in S and S' .

If so, cutting M along one of such parallel tori, we have the case with less $\# \pi_0(S)$. Therefore, the induction on $\# \pi_0(S)$ shows the uniqueness.

(a) Suppose that $S \cap S' \neq \emptyset$. Since M is irreducible, and since S and S' are essential, the standard argument shows that every component of $S \cap S'$ is a loop cutting torus components into annuli. Pick a torus component $T \subset S$ such that $T \cap S' \neq \emptyset$. Let N be the union of one or two components of $M \setminus \nu(S)$ attaching $\overline{\nu(T)}$. Every annulus $A \subset N$ which is a part of S' is essential. In particular, A is not boundary parallel. Otherwise, an isotopy would reduce $\# \pi_0(S \cap S')$. By Lemma 7.3, N is a Seifert manifold. By Lemma 7.4, we may assume that every annulus is a vertical subsurface. The fibration of A induces one of T , which agrees with that of N . Therefore, $N \cup \overline{\nu(T)}$ becomes a bigger Seifert manifold, so $S \setminus T$ would satisfy the condition.

(b) Suppose that no pair of torus components in S and S' is parallel. Then, every torus in S (resp. S') is contained in a Seifert component of $M \setminus \nu(S')$ (resp. $M \setminus \nu(S)$), and it is a vertical subsurface by Proposition 6.8. Every atoroidal component of $M \setminus \nu(S)$ (resp. $M \setminus \nu(S')$) must be contained in a Seifert

component of $M \setminus \nu(S')$ (resp. $M \setminus (S)$), and it is also a Seifert manifold. Then, there are components $N \subset M \setminus \nu(S)$ and $N' \subset M \setminus \nu(S')$ such that $N \cap N' \neq \emptyset$ and any Seifert fibrations of N and N' are different in $N \cap N'$, since all Seifert fibrations would agree and M would be a Seifert manifold otherwise. By Theorem 6.2 and 6.10, $N \cap N' \approx M(D^2; (2, 1), (2, 1)) \approx N_{1,1} \tilde{\times} S^1$. Since $\partial(N \cap N')$ is connected, we have $N = N \cap N' \subset N'$ or $N' = N \cap N' \subset N$, and this implies that the Seifert fibration of the small one could be replaced by that of the bigger one, a contradiction. \square

Proof of Lemma 7.3. Let M be atoroidal and $A \subset M$ be an essential annulus. We have the following cases:

- (a) ∂A lies in two tori $T_1, T_2 \subset \partial M$.
- (b) ∂A lies in one torus $T_0 \subset \partial M$: (b1) A is twisted, (b2) A is untwisted.

If we let $V := \nu(A \cup T_1 \cup T_2)$ or $\nu(A \cup T_0)$, then $V \approx \Sigma_{0,3} \times S^1$ for (a) and (b1), and $V \approx N_{1,2} \tilde{\times} S^1$ for (b2). Since M is atoroidal, $\partial V \setminus \partial M$ is compressible or boundary parallel. Such a compressible torus bounds $D^2 \times S^1$ (Exercise 6). Then, $M \approx V$ or $V \cup (\bigsqcup D^2 \times S^1)$. Since such $D^2 \times S^1$ is attached so that $\partial D \times \{*\}$ is not isotopic to a fiber of V because A is incompressible, we have M a Seifert manifold. \square

Sketch of Lemma 7.4. Let M be a Seifert manifold and $A \subset M$ be an essential annulus. By Proposition 6.8, A is isotopic to a horizontal or vertical subsurface. Suppose that A is a horizontal annulus. Then, A is a finite cover of the base manifold O . Since $\chi(O) = \chi(A) = 0$, $O \approx \Sigma_{0,2}$, $N_{1,1}$, or $D^2(2, 2)$, so $M \approx \Sigma_{0,2} \times S^1$, $N_{1,1} \tilde{\times} S^1$, or $M(D^2; (2, 1), (2, 1))$. Since we have $\Sigma_{0,2} \times S^1 \approx [0, 1] \times S^1 \times S^1$ and $N_{1,1} \tilde{\times} S^1 \approx M(D^2; (2, 1), (2, 1))$ with different Seifert fibrations, A becomes a vertical subsurface if we swap Seifert fibrations of M . \square