

Lebesgue Theory

Ikhan Choi

February 17, 2022

Contents

I	Measure theory	3
1	Measures and σ-algebras	4
1.1	Definition of measures	4
1.2	Carathéodory extension theorem	4
2	Measures on the real line	7
3	Measurable functions	8
3.1	Extended real numbers	8
3.2	Simple functions	8
II	Integration	9
4	Lebesgue integration	10
4.1	Definition of Lebesgue integration	10
4.2	Convergence theorems	10
4.3	Radon-Nikodym theorem	11
4.4	Modes of convergence	11
5	Product measures	13
5.1	Fubini-Tonelli theorem	13
5.2	Lebesgue measure on Euclidean spaces	13
6	Lebesgue spaces	14
6.1	L^p spaces	14
6.2	L^2 spaces	14
6.3	Dual spaces	14

III	Linear operators	15
7		16
8	Bounded linear operators	17
8.1	Continuity	17
8.2	Density arguments	17
8.3	Interpolation	17
9	Convergence of linear operators	18
9.1	Translation and multiplication operators	18
9.2	Convolution type operators	18
9.3	Computation of integral transforms	18
IV	Fundamental theorem of calculus	19
10	Weak derivatives	20
11	Absolutely continuity	21
12	Lebesgue differentiation theorem	22

Part I

Measure theory

Chapter 1

Measures and σ -algebras

1.1 Definition of measures

1.2 Carathéodory extension theorem

1.1 (Outer measures). Let Ω be a set. An *outer measure* on Ω is a function $\mu^* : \mathcal{P}(\Omega) \rightarrow [0, \infty]$ with $\mu^*(\emptyset) = 0$ such that

(i) if $E_1 \subset E_2$, then $\mu^*(E_1) \leq \mu^*(E_2)$, (monotonicity)

(ii) $\mu^*(\bigcup_{i=1}^{\infty} E_i) \leq \sum_{i=1}^{\infty} \mu^*(E_i)$, (countable subadditivity)

for any $\{E_i\}_{i=1}^{\infty} \subset \mathcal{P}(\Omega)$.

(a) A function $\mu^* : \mathcal{P}(\Omega) \rightarrow [0, \infty]$ with $\mu^*(\emptyset) = 0$ is an outer measure if and only if $\mu^*(E) \leq \sum_{i=1}^{\infty} \mu^*(E_i)$ whenever $E \subset \bigcup_{i=1}^{\infty} E_i$.

(b) Let $\mathcal{A} \subset \mathcal{P}(\Omega)$ with $\emptyset \in \mathcal{A}$. If a function $\rho : \mathcal{A} \rightarrow [0, \infty]$ satisfies $\rho(\emptyset) = 0$, then we can associate an outer measure $\mu^* : \mathcal{P}(\Omega) \rightarrow [0, \infty]$ by defining as

$$\mu^*(E) := \inf \left\{ \sum_{i=1}^{\infty} \rho(A_i) : E \subset \bigcup_{i=1}^{\infty} A_i, A_i \in \mathcal{A} \right\},$$

where we use the convention $\inf \emptyset = \infty$.

1.2 (Carathéodory measure). Let μ^* be an outer measure on a set Ω . A subset $A \subset \Omega$ is called *Carathéodory measurable* relative to μ^* if

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \setminus A)$$

for every subset $E \subset \Omega$. Let \mathcal{M} be the collection of all Carathéodory measurable subsets relative to μ^* .

- (a) \mathcal{M} is an algebra and μ^* is finitely additive on \mathcal{M} .
- (b) \mathcal{M} is a σ -algebra and μ^* is countably additive on \mathcal{M} .
- (c) The measure $\mu := \mu^*|_{\mathcal{M}} : \mathcal{M} \rightarrow [0, \infty]$ is complete. We call μ the *Carathéodory measure* constructed from ρ .

1.3 (Carathéodory extension theorem). Let $\mathcal{A} \subset \mathcal{P}(\Omega)$ with $\emptyset \in \mathcal{A}$. Let $\rho : \mathcal{A} \rightarrow [0, \infty]$ with $\rho(\emptyset) = 0$. Consider two conditions

- (i) $A \subset \bigcup_{i=1}^{\infty} A_i$ implies $\rho(A) \leq \sum_{i=1}^{\infty} \rho(A_i)$,
- (ii) for any $\varepsilon > 0$ and B, A there are A_1, A_2 such that $B \cap A \subset A_1$, $B \setminus A \subset A_2$ and $\rho(B) + \varepsilon > \rho(A_1) + \rho(A_2)$.

Let $\mu^* : \mathcal{P}(\Omega) \rightarrow [0, \infty]$ be the associated outer measure of ρ , and $\mu : \mathcal{M} \rightarrow [0, \infty]$ the measure defined by the restriction of μ^* on Carathéodory measurable subsets.

- (a) $\mu^*|_{\mathcal{A}} = \rho$ if (i) is satisfied.
- (b) $\mathcal{A} \subset \mathcal{M}$ if (ii) is satisfied.

Proof. (a) Clearly $\mu^*(A) \leq \rho(A)$ for $A \in \mathcal{A}$.

We may assume $\mu^*(A) < \infty$. For arbitrary $\varepsilon > 0$ there is $\{A_i\}_{i=1}^{\infty}$ such that $A \subset \bigcup_{i=1}^{\infty} A_i$ and

$$\mu^*(A) + \varepsilon > \sum_{i=1}^{\infty} \rho(A_i) \geq \rho(A).$$

(b) Let $E \in \mathcal{P}(\Omega)$ and $A \in \mathcal{A}$. Then, $E \subset \bigcup_{i=1}^{\infty} A_i$ and $A_i \cap A \subset A_{i,1}$ and $A_i \setminus A \subset A_{i,2}$ such that

$$\begin{aligned} \mu^*(E) + \varepsilon &> \sum_{i=1}^{\infty} \left(\rho(A_i) + \frac{\varepsilon}{2^{i+1}} \right) > \sum_{i=1}^{\infty} \rho(A_{i,1}) + \sum_{i=1}^{\infty} \rho(A_{i,2}) \\ &\geq \mu^*(E \cap A) + \mu^*(E \setminus A). \end{aligned}$$

□

1.4 (Carathéodory extension from semi-ring). Let $\mathcal{A} \subset \mathcal{P}(\Omega)$ be a semi-ring of sets on a set X . A function $\rho : \mathcal{A} \rightarrow [0, \infty]$ with $\rho(\emptyset) = 0$ is called a *pre-measure* if

- (i) $\rho(\bigsqcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \rho(A_i)$, (disjoint countable subadditivity)
- (ii) $\rho(\bigsqcup_{i=1}^n A_i) = \sum_{i=1}^n \rho(A_i)$, (finite additivity)

for any $\{A_i\}_{i=1}^\infty \subset \mathcal{A}$ with $\bigsqcup_{i=1}^\infty A_i \in \mathcal{A}$ and $n \in \mathbb{N}$.

Let $\mu^* : \mathcal{P}(\Omega) \rightarrow [0, \infty]$ be the associated outer measure of ρ , and $\mu : \mathcal{M} \rightarrow [0, \infty]$ the measure defined by the restriction of μ^* on Carathéodory measurable subsets.

(a) A pre-measure is a priori countably additive.

1.5 (Uniqueness of Carathéodory extensions). The Carathéodory extension theorem provides with a uniqueness theorem for measures.

Monotone class lemma: alternative direct proof method without using Carathéodory extension.

Chapter 2

Measures on the real line

distribution functions

Exercises

2.1. * A Lebesgue measurable set in \mathbb{R} with positive measure contains an arbitrarily long subsequence of an arithmetic progression.

Chapter 3

Measurable functions

3.1 Extended real numbers

3.2 Simple functions

Pointwise limit of simple functions is measurable.

Proof. Let $f(x) = \lim_{n \rightarrow \infty} s_n(x)$.

□

Every measurable extended real-valued function is a pointwise limit of simple functions.

Part II

Integration

Chapter 4

Lebesgue integration

4.1 Definition of Lebesgue integration

4.2 Convergence theorems

Stein: Egorov \rightarrow BCT \rightarrow Fatou \rightarrow MCT \rightarrow L1 is a measure

Stein: BCT + L1 is a measure \rightarrow DCT

Folland: MCT \rightarrow Fatou \rightarrow DCT \rightarrow BCT

4.1 (Egorov's theorem). Let Ω be a finite measure space. Let $(f_n : \Omega \rightarrow \mathbb{R})_n$ be a sequence of a.e. convergent measurable functions. For $\varepsilon > 0$, there exists a measurable $E_\varepsilon \subset \Omega$ such that $\mu(\Omega \setminus E_\varepsilon) < \varepsilon$ and f_n uniformly convergent on E_ε .

Proof. Assume $f_n \rightarrow 0$. The set of convergence is

$$\bigcap_{k>0} \bigcup_{n_0>0} \bigcap_{n \geq n_0} \{x : |f_n(x)| < \frac{1}{k}\},$$

which is a full set. We want to get rid of the dependence on the point x of n_0 in the union $\bigcup_{n_0>0}$. Since

$$\bigcap_{n \geq n_0} \{x : |f_n(x)| < \frac{1}{k}\}$$

is increasing as $n_0 \rightarrow \infty$ to a full set for each $k > 0$, we can find $n_0(k, \varepsilon)$ such that

$$\mu\left(\bigcap_{n \geq n_0} \{x : |f_n(x)| < \frac{1}{k}\}\right) > \mu(\Omega) - \frac{\varepsilon}{2^k}.$$

Then,

$$\mu\left(\bigcap_{k>0} \bigcap_{n \geq n_0} \{x : |f_n(x)| < \frac{1}{k}\}\right) > \mu(\Omega) - \varepsilon.$$

If we define

$$E_\varepsilon := \bigcap_{k>0} \bigcap_{n \geq n_0} \{x : |f_n(x)| < \frac{1}{k}\},$$

then for any $k > 0$ and $x \in E_\varepsilon$, and with the $n_0(k, \varepsilon)$ we have chosen, we have

$$n \geq n_0 \Rightarrow |f_n(x)| < \frac{1}{k}.$$

□

Since $\{f_n(x)\}_n$ diverges if and only if

$$\exists k > 0, \quad \forall n_0 > 0, \quad \exists n > n_0 : |f_n(x) - f(x)| > \frac{1}{k},$$

we have

$$\begin{aligned} \{x : \{f_n(x)\}_n \text{ diverges}\} &= \bigcup_{k>0} \bigcap_{n_0>0} \bigcup_{n>n_0} \{x : |f_n - f| > \frac{1}{k}\} \\ &= \bigcup_{k>0} \limsup_n \{x : |f_n - f| > \frac{1}{k}\}. \end{aligned}$$

Since for every k we have

$$\begin{aligned} \limsup_n \{x : |f_n - f| > \frac{1}{k}\} &\subset \limsup_{n>k} \{x : |f_n - f| > \frac{1}{n}\} \\ &= \limsup_n \{x : |f_n - f| > \frac{1}{n}\}, \end{aligned}$$

we have

$$\{x : \{f_n(x)\}_n \text{ diverges}\} \subset \limsup_n \{x : |f_n - f| > \frac{1}{n}\}.$$

4.3 Radon-Nikodym theorem

4.4 Modes of convergence

4.2 (Convergence in measure). Let (X, μ) be a measure space. Let f_n be a sequence of measurable functions. If f_n converges to f in measure, then f_n has a subsequence that converges to f μ -a.e.

Proof. We can extract a subsequence f_{n_k} such that

$$\mu(\{x : |f_{n_k} - f| > \frac{1}{k}\}) > \frac{1}{2^k}.$$

Since

$$\sum_{k=1}^{\infty} \mu(\{x : |f_{n_k} - f| > \frac{1}{k}\}) < \infty,$$

by the Borel-Canteli lemma, we get

$$\mu(\limsup_k \{x : |f_{n_k} - f| > \frac{1}{k}\}) = 0.$$

Therefore, f_{n_k} converges μ -a.e.

□

Chapter 5

Product measures

5.1 Fubini-Tonelli theorem

5.2 Lebesgue measure on Euclidean spaces

Chapter 6

Lebesgue spaces

6.1 L^p spaces

6.2 L^2 spaces

6.3 Dual spaces

riesz representations

Part III

Linear operators

Chapter 7

Chapter 8

Bounded linear operators

8.1 Continuity

Schur test

8.2 Density arguments

extension of operators

8.3 Interpolation

weak L_p , marcinkiewicz

Chapter 9

Convergence of linear operators

9.1 Translation and multiplication operators

9.2 Convolution type operators

approximation of identity

9.3 Computation of integral transforms

Part IV

Fundamental theorem of calculus

Chapter 10

Weak derivatives

The space of weakly differentiable functions with respect to all variables $= W_{\text{loc}}^{1,1}$.

10.1 (Product rule for weakly differentiable functions). We want to show that if u , v , and uv are weakly differentiable with respect to x_i , then $\partial_{x_i}(uv) = \partial_{x_i}u v + u \partial_{x_i}v$.

(a) If u is weakly differentiable with respect to x_i and $v \in C^1$, then $\partial_{x_i}(uv) = \partial_{x_i}u v + u \partial_{x_i}v$.

10.2 (Interchange of differentiation and integration). Let $f : \Omega \rightarrow \mathbb{R}$ such that $f(x, y)$ and $\partial_{x_i}f(x, y)$ are both locally integrable in x and integrable y . Then,

$$\partial_{x_i} \int f(x, y) dy = \int \partial_{x_i} f(x, y) dy$$

where ∂_{x_i} denotes the weak partial derivative.

Chapter 11

Absolutely continuity

- (a) f is Lip_{loc} iff f' is L_{loc}^{∞}
- (b) f is AC_{loc} iff f' is L_{loc}^1
- (a) f is Lip iff f' is L^{∞}
- (b) f is AC iff f' is L^1
- (c) f is BV iff f' is a finite regular Borel measure

Chapter 12

Lebesgue differentiation theorem