

# Foundations of Calculus

Ikhan Choi

May 8, 2022

# Contents

<b>I</b>	<b>Convergence</b>	<b>3</b>
<b>1</b>	<b>Sequences</b>	<b>4</b>
1.1	Control of the error . . . . .	4
1.2	Approximate sequences . . . . .	4
1.3	Bounded sequences . . . . .	4
1.4	Recursive sequences . . . . .	4
<b>2</b>	<b>Series</b>	<b>5</b>
2.1	Absolute convergence . . . . .	5
2.2	Convergence tests . . . . .	5
<b>3</b>	<b>Metrics and norms</b>	<b>7</b>
3.1	Metric spaces . . . . .	7
3.2	Normed spaces . . . . .	8
3.3	Open sets and closed sets . . . . .	8
3.4	Compact sets . . . . .	8
3.5	Connected sets . . . . .	8
<b>II</b>	<b>Real functions</b>	<b>9</b>
<b>4</b>	<b>Continuous functions</b>	<b>10</b>
4.1	Intermediate and extreme value theorems . . . . .	10
4.2	Uniform continuity . . . . .	10
4.3	Uniform convergence . . . . .	10
<b>5</b>	<b>Differentiable functions</b>	<b>11</b>
5.1	Monotonicity and convexity . . . . .	11
5.2	Mean value theorem . . . . .	11
5.3	Taylor's theorem . . . . .	11
5.4	Differentiable class . . . . .	11
<b>6</b>	<b>Analytic functions</b>	<b>12</b>
6.1	Power series . . . . .	12
6.2	Complex analytic functions . . . . .	12
6.3	Special functions . . . . .	12

<b>III</b>	<b>Integration</b>	<b>13</b>
<b>7</b>	<b>Riemann integral</b>	<b>14</b>
7.1	Riemann integral . . . . .	14
7.2	Henstock-Kurzweil intergral . . . . .	14
7.3	Improper integral . . . . .	14
7.4	Fundamental theorem of calculus for continuous functions . . . . .	14
<b>8</b>	<b>Integrable functions</b>	<b>15</b>
8.1	. . . . .	15
<b>9</b>		<b>16</b>
<b>IV</b>	<b>Multivariable Calculus</b>	<b>17</b>
<b>10</b>	<b>Fréchet derivatives</b>	<b>18</b>
10.1	Tangent space . . . . .	18
<b>11</b>	<b>Inverse function theorem</b>	<b>19</b>
11.1	Banach fixed point theorem . . . . .	19
11.2	Variations of the inverse function theorem . . . . .	19
<b>12</b>	<b>Differential forms</b>	<b>20</b>

**Part I**

**Convergence**

# Chapter 1

## Sequences

### 1.1 Control of the error

preserving inequalities limsup and liminf

### 1.2 Approximate sequences

### 1.3 Bounded sequences

monotone convergence Bolzano-Weierstrass

### 1.4 Recursive sequences

?

## Exercises

1.1. Every real sequence  $(a_n)_{n=1}^{\infty}$  has a monotonic subsequence  $(a_{n_k})_{k=1}^{\infty}$  such that  $\lim_{k \rightarrow \infty} a_{n_k} = \limsup_{n \rightarrow \infty} a_n$ .

# Chapter 2

## Series

### 2.1 Absolute convergence

2.1 (Unconditional convergence).

### 2.2 Convergence tests

comparison limit comparison cauchy condensation integral....

ratio root

2.2 (Abel transform).

$$A_k(B_k - B_{k-1}) + (A_k - A_{k-1})B_{k-1} = A_k B_k - A_{k-1} B_{k-1}$$

$$\sum_{m < k \leq n} A_k b_k = A_n B_n - A_m B_m - \sum_{m < k \leq n} a_k B_{k-1}.$$

abel test

2.3 (Dirichlet test).

2.4 (Mertens' theorem). If  $\sum_{k=0}^{\infty} a_k$  converges to  $A$  absolutely and  $\sum_{k=0}^{\infty} b_k$  converges to  $B$ , then their Cauchy product  $\sum_{k=0}^{\infty} c_k$  with  $c_k := \sum_{l=0}^k a_l b_{k-l}$  converges to  $AB$ .

*Proof.* Let

$$A_n := \sum_{k=0}^n a_k, \quad B_n := \sum_{k=0}^n b_k, \quad \text{and} \quad C_n := \sum_{k=0}^n c_k.$$

Consider the regions

$$T_n := \{(k, l) \in \mathbb{Z}_{\geq 0}^2 : k + l \leq n\}, \quad R_m := \{(k, l) \in \mathbb{Z}_{\geq 0}^2 : k \leq m\}.$$

Write

$$\begin{aligned} AB - C_n &= \sum_{k \leq m} \sum_{l > n-k} a_k b_l + \sum_{k > m} \sum_{l \geq 0} a_k b_l - \sum_{m < k \leq n} \sum_{l \leq n-k} a_k b_l \\ &= \sum_{k \leq m} a_k (B - B_{n-k}) + \sum_{k > m} a_k B - \sum_{m < k \leq n} a_k B_{n-k}. \end{aligned}$$

The first term

$$\left| \sum_{k \leq m} a_k (B - B_{n-k}) \right| \leq (\max_k |a_k|) \left( \sum_{l \geq n-m} |B - B_l| \right)$$

converges to zero as  $n \rightarrow \infty$  for fixed  $m$ , the second term

$$|\sum_{k>m} a_k B| \leq |A - A_m| |B|$$

converges to zero as  $m \rightarrow \infty$  for any  $n$ , and finally the third term

$$|\sum_{m<k\leq n} a_k B_{n-k}| \leq (\sum_{k>m} |a_k|)(\max_l |B_l|)$$

converges to zero as  $m \rightarrow \infty$  for any  $n$ .

Fix  $m$  such that the second and third terms are bounded by arbitrary  $\frac{\varepsilon}{2} > 0$  so that

$$|C_n - AB| \leq |\sum_{k\leq m} a_k (B - B_{n-k})| + \frac{\varepsilon}{2} + \frac{\varepsilon}{2}.$$

Then, by taking  $n \rightarrow \infty$ , we obtain

$$\limsup_{n \rightarrow \infty} |C_n - AB| \leq \varepsilon.$$

Since  $\varepsilon$  is arbitrary, we have

$$\lim_{n \rightarrow \infty} C_n = AB.$$

□

## Exercises

2.5. If  $a_n \rightarrow 0$ , then  $\frac{1}{n} \sum_{k=1}^n a_k \rightarrow 0$ .

2.6. If  $a_n \geq 0$  and  $\sum a_n$  diverges, then  $\sum \frac{a_n}{1+a_n}$  also diverges.

2.7. If  $a_n \downarrow 0$  and  $S_n \leq 1 + na_n$ , then  $S_n \leq 1$ .

## Chapter 3

# Metrics and norms

### 3.1 Metric spaces

**3.1** (Definition of metric spaces). Let  $X$  be a set. A *metric* is a function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  such that

- (i)  $d(x, y) = 0$  if and only if  $x = y$ , (nondegeneracy)
- (ii)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ , (symmetry)
- (iii)  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$ . (triangle inequality)

A pair  $(X, d)$  of a set  $X$  and a metric on  $X$  is called a *metric space*. We often write it simply  $X$ .

- (a) A normed space  $X$  is a metric space with a metric defined by  $d(x, y) := \|x - y\|$ .
- (b) A subset of a metric space is a metric space with a metric given by restriction.

**3.2** (System of open balls). A metric is often misunderstood as something that measures a distance between two points and belongs to the study of geometry. The main function of a metric is to make a system of small balls, sets of points whose distance from specified center points is less than fixed numbers. The balls centered at each point provide a concrete images of “system of neighborhoods at a point” in a more intuitive sense. In this viewpoint, a metric can be considered as a structure that lets someone accept the notion of neighborhoods more friendly.

Note that taking either  $\varepsilon$  or  $\delta$  in analysis really means taking a ball of the very radius. Investigation of the distribution of open balls centered at a point is now an important problem.

Let  $X$  be a metric space. A set of the form

$$\{y \in X : d(x, y) < \varepsilon\}$$

for  $x \in X$  and  $\varepsilon > 0$  is called an *open ball centered at  $x$  with radius  $\varepsilon$*  and denoted by  $B(x, \varepsilon)$  or  $B_\varepsilon(x)$ .

**3.3** (Convergence and continuity in metric spaces). Let  $\{x_n\}_n$  be a sequence of points on a metric space  $(X, d)$ . We say that a point  $x$  is a *limit* of the sequence or the sequence *converges to  $x$*  if for arbitrarily small ball  $B(x, \varepsilon)$ , we can find  $n_0$  such that  $x_n \in B(x, \varepsilon)$  for all  $n > n_0$ . If it is satisfied, then we write

$$\lim_{n \rightarrow \infty} x_n = x,$$

or simply  $x_n \rightarrow x$  as  $n \rightarrow \infty$ . We say a sequence is *convergent* if it converges to a point. If it does not converge to any points, then we say the sequence *diverges*.

A function  $f : X \rightarrow Y$  between metric spaces is called *continuous at  $x \in X$*  if for any ball  $B(f(x), \varepsilon) \subset Y$ , there is a ball  $B(x, \delta) \subset X$  such that  $f(B(x, \delta)) \subset B(f(x), \varepsilon)$ . The function  $f$  is called *continuous* if it is continuous at every point on  $X$ .



- (a) A sequence  $x_n$  in a metric space  $X$  converges to  $x \in X$  if and only if  $d(x_n, x)$  converges to zero.
- (b) Let  $f : X \rightarrow Y$  be a function between two metric spaces. If there is a constant  $C$  such that  $d(x, y) \leq C d(f(x), f(y))$  for all  $x$  and  $y$  in  $X$ , then  $f$  is continuous. In this case,  $f$  is particularly called *Lipschitz continuous* with the *Lipschitz constant*  $C$ .

## 3.2 Normed spaces

banach space

## 3.3 Open sets and closed sets

convergence, limit point

## 3.4 Compact sets

## 3.5 Connected sets

## Exercises

## **Part II**

# **Real functions**

## Chapter 4

# Continuous functions

### 4.1 Intermediate and extreme value theorems

### 4.2 Uniform continuity

### 4.3 Uniform convergence

#### Exercises

4.1. The set of local minima of a convex real function is connected.

4.2. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be continuous. The equation  $f(x) = c$  cannot have exactly two solutions for every constant  $c \in \mathbb{R}$ .

4.3. A continuous function that takes on no value more than twice takes on some value exactly once.

4.4. Let  $f$  be a function that has the intermediate value property. If the preimage of every singleton is closed, then  $f$  is continuous.

4.5. \* If a sequence of real functions  $f_n : [0, 1] \rightarrow [0, 1]$  satisfies  $|f(x) - f(y)| \leq |x - y|$  whenever  $|x - y| \geq \frac{1}{n}$ , then the sequence has a uniformly convergent subsequence.

## Chapter 5

# Differentiable functions

### 5.1 Monotonicity and convexity

### 5.2 Mean value theorem

Darboux

### 5.3 Taylor's theorem

### 5.4 Differentiable class

completeness

## Exercises

5.1. If  $\lim_{x \rightarrow \infty} f(x) = a$  and  $\lim_{x \rightarrow \infty} f'(x) = b$ , then  $a = 0$ .

5.2. Let  $f$  be a real  $C^2$  function with  $f(0) = 0$  and  $f''(0) \neq 0$ . Define a function  $\xi$  such that  $f(x) = xf'(\xi(x))$  with  $|\xi| \leq |x|$ , we have  $\xi'(0) = 1/2$ .

5.3. Let  $f$  be a  $C^2$  function such that  $f(0) = f(1) = 0$ . We have  $\|f\| \leq \frac{1}{8}\|f''\|$ .

5.4. A smooth function such that for each  $x$  there is  $n$  having the  $n$ th derivative vanish is a polynomial.

5.5. If a real  $C^1$  function  $f$  satisfies  $f(x) \neq 0$  for  $x$  such that  $f'(x) = 0$ , then in a bounded set there are only finite points at which  $f$  vanishes.

5.6. Let a real function  $f$  be differentiable. For  $a < a' < b < b'$  there exist  $a < c < b$  and  $a' < c' < b'$  such that  $f(b) - f(a) = f'(c)(b - a)$  and  $f(b') - f(a') = f'(c')(b' - a')$ .

5.7. Let  $f$  be a differentiable function on the unit closed interval. If  $f(0) = 0$  there is  $c$  such that  $cf'(c) = f(c)$ . (Flett)

5.8. Let  $f$  be a differentiable function on the unit closed interval. If  $f(0) = 0$  there is  $c$  such that  $cf(c) = (1 - c)f'(c)$ .

## Chapter 6

# Analytic functions

### 6.1 Power series

uniform convergence and absolute convergence, abel theorem? differentiation convergence of radius  
sum, product, composition, reciprocal? closed under uniform convergence

### 6.2 Complex analytic functions

complex domain (real analytic iff its domain contains real line) convergence of radius, revisited identity  
theorem

### 6.3 Special functions

hypergeometric, bessel, gamma, zeta

### Exercises

# **Part III**

## **Integration**

## Chapter 7

# Riemann integral

### 7.1 Riemann integral

tagged partition

### 7.2 Henstock-Kurzweil intergral

bounded compact support  $\leftrightarrow$  lebesgue

### 7.3 Improper integral

### 7.4 Fundamental theorem of calculus for continuous functions

### Exercises

7.1. Find the value of  $\lim_{n \rightarrow \infty} \frac{1}{n} \left( \sum_{k=1}^n \frac{1}{n} f\left(\frac{k}{n}\right) - \int_0^1 f(x) dx \right)$ .

7.2. If  $xf'(x)$  is bounded and  $x^{-1} \int_0^x f \rightarrow L$  then  $f(x) \rightarrow L$  as  $x \rightarrow \infty$ .

## Chapter 8

# Integrable functions

### 8.1



## Chapter 9

**Part IV**

**Multivariable Calculus**

## Chapter 10

# Fréchet derivatives

### 10.1 Tangent space

## **Chapter 11**

# **Inverse function theorem**

### **11.1 Banach fixed point theorem**

### **11.2 Variations of the inverse function theorem**

## **Chapter 12**

# **Differential forms**