

Haagerup's problems on normal weights

Ikhan Choi

Operator Algebra Seminars at the University of Tokyo
13 May 2025

Abstract

We will introduce three open problems related to normal weights posed by Haagerup fifty years ago in his master's thesis. Among these, we especially discuss a solution to the last problem and its meaning. This will provide a complete set of the positive Hahn-Banach separation theorems on operator algebras and shed light on why the weak* topologies on the duals of C^* -algebras are relatively difficult to work with. (arXiv:2501.16832.)

Contents

1	Weights	1
1.1	Normality	1
1.2	Semi-cyclic representations	2
2	Haagerup's three problems	2
2.1	Problem 1.10	2
2.2	Problem 1.11	2
2.3	Problem 2.7	3
3	Positive Hahn-Banach separation	4
3.1	Proof of (a)	4
3.2	Proof of (b)	5
3.3	Proof of (d)	6

1 Weights

1.1 Normality

Definition. Let A be a C^* -algebra. A *weight* on A is a homogeneous additive functional $\varphi : A^+ \rightarrow [0, \infty]$. A *subadditive weight* on A is a homogeneous subadditive functional $\varphi : A^+ \rightarrow [0, \infty]$.

Definition. For a weight φ on a C^* -algebra A , we say it is

- (i) *faithful* if $\varphi(a) = 0$ implies $a = 0$ for $a \in A^+$,
- (ii) *densely defined* if $\varphi^{-1}([0, \infty))$ is norm dense in A^+ ,
- (iii) *lower semi-continuous* if $\varphi^{-1}([0, 1])$ is norm closed in A^+ .

For a weight φ on a von Neumann algebra M , we say it is

- (ii') *semi-finite* if $\varphi^{-1}([0, \infty))$ is σ -weakly dense in M^+ ,
- (iii') *normal* if ...?

Theorem (Haagerup, 1975). *For a weight φ on a von Neumann algebra, the followings are all equivalent.*

- (1) φ is completely additive for positive elements.
- (2) φ preserves directed suprema.
- (3) φ is σ -weakly lower semi-continuous.
- (4) φ is given by the pointwise supremum of normal positive linear functionals.

Remark. Every von Neumann algebra admits a faithful semi-finite normal weight. There is a C^* -algebra that does not admit a faithful densely defined lower semi-continuous weight.

Let φ be a non-zero densely defined subadditive weight on a unital C^* -algebra A . Since φ is densely defined, there is $a \in A^+$ such that $\|1 - a\| < 1$ and $\varphi(a) < \infty$. Then, a is invertible so that $1 = a^{-1}a \leq \|a^{-1}\|a$, which implies $\varphi(1) < \infty$ and φ is bounded. Therefore, it follows that the condition that A admits a faithful densely defined weight means that A admits a faithful state. There is a compact Hausdorff space that does not admit a regular Borel measure of full support.

1.2 Semi-cyclic representations

Definition. A *semi-cyclic representation* of a C^* -algebra A is a representation $\pi : A \rightarrow B(H)$ together with a partially defined left A -linear operator $\Lambda : A \rightarrow H$ with dense range.

2 Haagerup's three problems

2.1 Problem 1.10

$$\left\{ \begin{array}{l} \text{For a subadditive weight on a von Neumann algebra,} \\ \text{is it } \sigma\text{-weakly lower semi-continuous if it preserves directed suprema?} \end{array} \right\}$$

Haagerup provided a proof of (1) $\Rightarrow$ (3) directly without an intermediate step (2). The proof is divided into two parts. First, it is proved for σ -finite von Neumann algebras, and extended to general von Neumann algebras. In a σ -finite von Neumann algebra, the σ -strong topology on the unit ball is metrizable. The additivity of a weight is used in the proof for the case of σ -finite von Neumann algebras, so it is enough to solve in the σ -finite case for this problem.

2.2 Problem 1.11

$$\left\{ \begin{array}{l} \text{For a weight on a von Neumann algebra,} \\ \text{is it normal if it is normal on every commutative von Neumann subalgebra?} \end{array} \right\}$$

Recall that the following theorem, which discuss normality of positive linear functionals.

Theorem (Dixmier, 1953). *For a positive linear functional ω on a von Neumann algebra, the followings are all equivalent.*

- (0) ω is completely additive for orthogonal projections.

- (1) ω is completely additive for positive elements.
- (2) ω preserves directed suprema.
- (3) ω is σ -weakly continuous.

Corollary. A (positive) linear functional on a von Neumann algebra is normal if it is normal on every commutative von Neumann subalgebra.

This corollary is used to prove some equivalent characterizations for weak compactness in M_* . See Theorem III.5.4.

Remark. (0) $\Rightarrow$ (1) is false for weights. Let φ be a weight on $\ell^\infty(\mathbb{N})$ defined by

$$\varphi(x) := \begin{cases} \sum_n x_n & \text{if } x = (x_n) \in c_c(\mathbb{N}), \\ \infty & \text{otherwise,} \end{cases} \quad x \in \ell^\infty(\mathbb{N})^+.$$

Then, it is not σ -weakly lower semi-continuous but is completely additive for orthogonal projections.

2.3 Problem 2.7

Definition. Let (E, E^*) be a dual pair of (directed partially) ordered real vector spaces such that E^+ and E^{*+} are mutually dual cones, i.e. $E^{*+} = \{x^* \in E^* : x^*(x) \geq 0 \text{ for } x \in E\}$. For a subset F of E^+ , we say it is *hereditary* if $0 \leq x \leq y \in F$ implies $x \in F$, and its *positive polar* is defined by the positive part of the real polar

$$F^{r+} := \{x^* \in E^{*+} : \sup_{x \in F} x^*(x) \leq 1\}.$$

Theorem (Haagerup 1975, C. 2025). For a von Neumann algebra M and a C^* -algebra A , consider the ordered real dual pairs (M^{sa}, M_*^{sa}) and (A^{sa}, A^{sa*}) .

- (a) If F is a σ -weakly closed convex hereditary subset of M^+ , then $F = F^{r+r+}$.
- (b) If F_* is a norm closed convex hereditary subset of M_*^+ , then $F_* = F_*^{r+r+}$.
- (c) If F is a norm closed convex hereditary subset of A^+ , then $F = F^{r+r+}$.
- (d) If F^* is weakly* closed convex hereditary subset of A^{*+} , then $F^* = (F^*)^{r+r+}$.

Corollary (Haagerup, 1975). For a weight φ on a von Neumann algebra, the followings are all equivalent.

- (3) φ is σ -lower semi-continuous.
- (4) φ is given by the pointwise supremum of normal positive linear functionals.

Proof. Let φ be a σ -weakly lower semi-continuous subadditive weight on a von Neumann algebra M . Define

$$F := \{x \in M^+ : \varphi(x) \leq 1\}, \quad F_* := \{\omega \in M_*^+ : \omega \leq \varphi\}.$$

Note that

$$F_* = F^{r+} := \{\omega \in M_*^+ : \sup_{x \in F} \omega(x) \leq 1\}$$

Then, the statement

$$\varphi(x) = \sup_{\omega \in F_*} \omega(x), \quad x \in M^+,$$

is equivalent to

$$F = F_*^{r+} := \{x \in M^+ : \sup_{\omega \in F_*} \omega(x) \leq 1\}.$$

Since F is σ -weakly closed by the σ -weak lower semi-continuity of φ , and since F is clearly convex and hereditary by definition of subadditive weights, we have $F = F^{r+r+}$ by the part (a) of the previous theorem. $\square$

Corollary (Combes, 1968). *For a weight φ on a C^* -algebra, the followings are all equivalent.*

(3) φ is lower semi-continuous.

(4) φ is given by the pointwise supremum of positive linear functionals.

Proof. Same as above but by (c) instead of (a). $\square$

Remark. (c) follows from the direct application of (a), but it can be also shown by purely C^* -algebraic methods with unitization, and this is what Combes essentially did in 1968. (probably?)

Corollary. *For a von Neumann algebra M and a C^* -algebra A , there are one-to-one correspondences*

$$\left\{ \begin{array}{c} \text{normal} \\ \text{subadditive} \\ \text{weights on } M \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \sigma\text{-weakly closed} \\ \text{convex hereditary} \\ \text{subsets of } M^+ \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{norm closed} \\ \text{convex hereditary} \\ \text{subsets of } M_*^+ \end{array} \right\}$$

and

$$\left\{ \begin{array}{c} \text{lower semi-continuous} \\ \text{subadditive} \\ \text{weights on } A \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{norm closed} \\ \text{convex hereditary} \\ \text{subsets of } A^+ \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{weakly}^* \text{ closed} \\ \text{convex hereditary} \\ \text{subsets of } A^{*+} \end{array} \right\}$$

Remark. For a subadditive weight φ on A , it is a weight if and only if

$$G^* := \{\omega \in A^{*+} : \exists \varepsilon > 0, (1 + \varepsilon)\omega \leq \varphi\}$$

is directed. Similarly for von Neumann algebras.

3 Positive Hahn-Banach separation

3.1 Proof of (a)

Definition (Suppression by the one-parameter family of functional calculi). For $\delta > 0$, we define a function $f_\delta : (-\delta^{-1}, \infty) \rightarrow \mathbb{R}$ such that

$$f_\delta(t) := t(1 + \delta t)^{-1}, \quad t > -\delta^{-1}.$$

Then, f_δ is operator monotone, σ -strongly continuous, and enjoys a semi-group property, etc.

Original proof of (a). Since the positive polar is represented by the real polar

$$F^{r+} = F^r \cap M_*^+ = F^r \cap (-M^+)^r = (F \cup -M^+)^r = (F - M^+)^r,$$

the positive bipolar can be written as

$$F^{r+r^+} = (F - M^+)^{r+r^+} = (\overline{F - M^+})^+$$

by the usual real bipolar theorem. Since $F = (F - M^+)^+ \subset (\overline{F - M^+})^+$, it suffices to show $(\overline{F - M^+})^+ \subset F$.

Let $x \in (\overline{F - M^+})^+$. Then, we can take a net $x_i \in F - M^+$ such that $x_i \rightarrow x$ σ -weakly in M , and a net $y_i \in F$ such that $x_i \leq y_i$ for all i . Observe that if x_i were bounded by $r > 0$, then assuming $x_i \rightarrow x$ σ -strongly and $f_\delta(y_i) \rightarrow y_\delta$ σ -weakly for each $0 < \delta < r^{-1}$, we get

$$x_i \leq y_i \in F, \quad f_\delta(x_i) \leq f_\delta(y_i) \in F, \quad 0 \leq f_\delta(x) \leq y_\delta \in F, \quad f_\delta(x) \in F, \quad x \in F.$$

However, x_i may not be bounded, so we find a proof that uses the Krein-Šmulian theorem.

Define

$$G := \{x \in M^{sa} : \text{for any sufficiently small } \delta > 0, f_\delta(x) \in F - M^+\}.$$

Since $F - M^+ \subset G$ and $G^+ \subset F$, we are done if we prove G is σ -weakly closed in M , and we can apply the Krein-Šmulian theorem to this intermediate subset. Fix $r > 0$ arbitrarily and let $M_r := \{x \in M : \|x\| \leq r\}$.

We first show that $G \cap M_r$ is σ -strongly closed. Let $x \in \overline{G \cap M_r}$ and fix arbitrary $0 < \delta < r^{-1}$. Take a net $x_i \in G \cap M_r$ such that $x_i \rightarrow x$ σ -strongly, and take a net $y_{i,\delta/2} \in F$ such that $f_{\delta/2}(x_i) \leq y_{i,\delta/2}$. Assuming $f_{\delta/2}(y_{i,\delta/2}) \rightarrow y_\delta$ σ -weakly, we get

$$f_{\delta/2}(x_i) \leq y_{i,\delta/2} \in F, \quad f_\delta(x_i) \leq f_{\delta/2}(y_{i,\delta/2}) \in F, \quad f_\delta(x) \leq y_\delta \in F, \quad f_\delta(x) \leq F - M^+, \quad x \in G.$$

We next show that $G \cap M_r$ is convex. It suffices to check

$$G \cap M_r = \overline{(F - M^+) \cap M_{2r}} \cap M_r.$$

If $x \in G \cap M_r$, then since $f_\delta(x) \in (F - M^+) \cap M_{2r}$ for small $\delta > 0$, we have $x \in \overline{(F - M^+) \cap M_{2r}} \cap M_r$. Conversely if $x \in \overline{(F - M^+) \cap M_{2r}} \cap M_r$, then there is a bounded net $x_i \in (F - M^+) \cap M_{2r} \subset G$ such that $x_i \rightarrow x$ σ -strongly, then by the σ -strong closedness of $G \cap M_{2r}$, we get $x \in G \cap M_r$.

Now we can deduce that, because G is convex and $G \cap M_r$ is σ -weakly closed, G is σ -weakly closed by the Krein-Šmulian theorem. $\square$

3.2 Proof of (b)

Definition (Bounded commutant Radon-Nikodym derivatives). Let M be a von Neumann algebra, and let $\psi \in M_*^+$. Consider the Gelfand-Naimark-Segal representation $\pi : M \rightarrow B(H)$ associated to ψ , together with the canonical cyclic vector $\Omega \in H$. Then, we have a positive bounded linear map $\theta : \pi(M)' \rightarrow M_*$ defined such that

$$\theta(h)(x) := \langle h\pi(x)\Omega, \Omega \rangle, \quad h \in \pi(M)', \quad x \in M.$$

Proof of (b). It is enough to prove $(\overline{F_* - M_*^+})^+ \subset F_*$.

Let $\omega \in (\overline{F_* - M_*^+})^+$. Then, we can take a sequence $\omega_n \in F_* - M_*^+$ such that $\omega_n \rightarrow \omega$ in norm of M_* , and a sequence $\varphi_n \in F_*$ such that $\omega_n \leq \varphi_n$ for all n . We may assume $\|\omega_n - \omega\| \leq 2^{-n}$ for all n by

passing to a subsequence. Define

$$\psi := \omega + \sum_n [\omega_n - \omega] + \sum_n 2^{-n} \frac{\varphi_n}{1 + \|\varphi_n\|} \in M_*^+,$$

and let $\theta : \pi(M)' \rightarrow M_*$ be the commutant Radon-Nikodym map associated to ψ . Since $-\psi \leq \omega_n \leq \psi$ implies the boundedness $\|\theta^{-1}(\omega_n)\| \leq 1$ for all n , the weak convergence $\omega_n \rightarrow \omega$ in M_* implies the convergence $\theta^{-1}(\omega_n) \rightarrow \theta^{-1}(\omega)$ in the weak operator topology of $\pi(M)'$. By the Mazur lemma, we can take a net ω_i in the convex hull of ω_n such that $\theta^{-1}(\omega_i) \rightarrow \theta^{-1}(\omega)$ strongly in $\pi(M)'$, and the corresponding $\varphi_i \in F^*$ can be defined such that $\omega_i \leq \varphi_i$ for all i . (In fact, the net ω_i can be taken to be a sequence because the commutant is σ -finite by the existence of the separating vector, but it is not necessary in here.) For each i and $0 < \delta < 1$, define

$$\omega_\delta := \theta(f_\delta(\theta^{-1}(\omega))), \quad \omega_{i,\delta} := \theta(f_\delta(\theta^{-1}(\omega_i))), \quad \varphi_{i,\delta} := \theta(f_\delta(\theta^{-1}(\varphi_i))).$$

Then,

$f_\delta(\theta^{-1}(\omega_i)) \rightarrow f_\delta(\theta^{-1}(\omega))$ strongly in $\pi(M)'$ implies $\omega_{i,\delta} \rightarrow \omega_\delta$ weakly in M_* ,

$f_\delta(\theta^{-1}(\omega)) \rightarrow \theta^{-1}(\omega)$ strongly in $\pi(M)'$ implies $\omega_\delta \rightarrow \omega$ weakly in M_* ,

definition $\varphi_\delta := \theta(\lim_{\mathcal{U}} f_\delta(\theta^{-1}(\varphi_i)))$ implies $\varphi_{i,\delta} \rightarrow \varphi_\delta$ weakly in M_* .

Since $\omega_i \leq \varphi_i$ and $0 \leq \varphi_{i,\delta} \leq \varphi_i \in F_*$, we get

$$\omega_{i,\delta} \leq \varphi_{i,\delta} \in F_*, \quad 0 \leq \omega_\delta \leq \varphi_\delta \in F_*, \quad \omega_\delta \in F_*, \quad \omega \in F_*.$$

□

3.3 Proof of (d)

The proof will start with two nets ω_i and φ_i such that $\omega_i \rightarrow \omega$ weakly* and $\omega_i \leq \varphi_i \in F^*$.

Question 1. How can we choose ψ with ω_i and φ_i ?

To construct ψ such that ω_i and φ_i are all absolutely continuous, it seems that we need to assume A is separable so that the weak* topology on the unit ball of A^* is metrizable and we may assume ω_i and φ_i are sequences. But even if this is the case, we can have $[\omega_n - \omega] \not\rightarrow 0$ in the weak* topology, contrary to the norm. So it seems to be difficult to construct ψ such that $\theta^{-1}(\omega_n)$ is bounded. However, we can still have $-2^n \leq \theta^{-1}(\omega_n) \leq \theta^{-1}(\varphi_n) \leq 2^n$, etc.

OK, now let us see why we need to give up using the weak operator topology of $\pi(A)'$.

First, anyway, $\theta^{-1}(\omega_n)$ is not, and should not be bounded by the second reason. Second, more importantly, the weak operator topology is too strong. The topologies on $\pi(A)'$ generated by vectors $\pi(A)\Omega$ and $\pi^{**}(A^{**})\Omega$ are totally different when they are induced in A^* via θ , i.e. weak* and weak topologies. The weak (operator) convergence of $\theta^{-1}(\omega_n)$ implies the weak convergence of ω_n . The Mazur lemma tells us that we may assume $\omega_n \rightarrow \omega$ in norm. Something is strange.

Then, use the compactness of the unit ball of A^* directly! Since $\theta^{-1}(\omega_n)$ is unbounded, we can try to take δ dynamically. We want to have the following properties:

$$-\delta_n < \theta^{-1}(\omega_n), \quad \theta(f_{\delta_n}(\theta^{-1}(\varphi_n))) \lesssim 1.$$

However, it is desparate to control the norm of $\theta(k)$ in terms of k , except by the norm of k . Also, by the “stiffness” or “rigidness” of the operator monotone functional calculi, two conditions are hardly

satisfied simultaneously.

Solution 1. Take ψ dynamically instead of δ !

Question 2. How can the limit of ω_i commute with f_δ without strong topology?

Solution 2. We can approximate f_δ with affine functions by

$$t - \delta^{\frac{1}{2}} \leq f_\delta(t) \leq t, \quad |t| \leq 2^{-1} \delta^{-\frac{1}{4}}$$

and

$$(1 + \delta^{-1})t \leq f_\delta(t) \leq t, \quad 0 \leq t \leq 1.$$

Proof of (d). As above, we will prove $(\overline{F^* - A^{*+}})^+ \subset F^*$. Define

$$G^* := \left\{ \omega \in A^{*sa} : \begin{array}{l} \text{there is } \psi \in A^{*+}, \text{ and there is } \varphi_\delta \in F^* \\ \text{for any sufficiently small } \delta > 0, \text{ such that} \\ \|\psi\| \leq 1, \|\varphi_\delta\| \leq \delta^{-1}, \text{ and } \omega \leq \varphi_\delta + \delta^{\frac{1}{2}}\psi \end{array} \right\}.$$

It suffices to show $F^* - A^{*+} \subset G^*$, $G^{*+} \subset F^*$, and $\overline{G^*} \subset G^*$.

Let $\omega \in F^* - A^{*+}$. Take $\varphi \in F^*$ such that $\omega \leq \varphi$. Define, for $\delta > 0$,

$$\psi := \frac{[\omega]}{1 + \|\omega\|} + \frac{\varphi}{(1 + \|\omega\|)(1 + \|\varphi\|)}, \quad \varphi_\delta := \theta(f_\delta(\theta^{-1}(\varphi))),$$

where θ is the commutant Radon-Nikodym map associated to ψ . The norm conditions $\|\psi\| \leq 1$ and $\|\varphi_\delta\| \leq \delta^{-1}$ are easily checked. For sufficiently small $\delta > 0$ such that $\|\theta^{-1}(\omega)\| \leq 1 + \|\omega\| \leq 2^{-1} \delta^{-\frac{1}{4}}$ and $\delta \leq 1$, we have

$$\omega = \theta(\theta^{-1}(\omega)) \leq \theta(f_\delta(\theta^{-1}(\omega)) + \delta^{\frac{1}{2}}) \leq \theta(f_\delta(\theta^{-1}(\varphi))) + \theta(\delta^{\frac{1}{2}}) = \varphi_\delta + \delta^{\frac{1}{2}}\psi,$$

so $\omega \in G^*$.

Let $\omega \in G^{*+}$. Take $\psi \in A^{*+}$ and $\varphi_\delta \in F^*$ such that $\|\psi\| \leq 1$, $\|\varphi_\delta\| \leq \delta^{-1}$, and $\omega \leq \varphi_\delta + \delta^{\frac{1}{2}}\psi$, for any sufficiently small $\delta > 0$. Let $\psi_\delta := \omega + \delta\varphi + \psi$, and let θ_δ be the associated commutant Radon-Nikodym map. For any fixed $\delta' > 0$, since $0 \leq \theta_\delta^{-1}(\omega) \leq 1$, we have

$$\begin{aligned} 0 &\leq (1 + \delta')^{-1} \theta_\delta(\omega) \leq f_{\delta'}(\theta_\delta^{-1}(\omega)) \leq f_{\delta'}(\theta_\delta^{-1}(\varphi_\delta + \delta^{\frac{1}{2}}\psi)) \\ &\leq f_{\delta'}(\theta_\delta^{-1}(\varphi_\delta) + \delta^{\frac{1}{2}}) \leq f_{\delta'}(\theta_\delta^{-1}(\varphi_\delta)) + \delta^{\frac{1}{2}}, \end{aligned}$$

and it implies $0 \leq (1 + \delta')^{-1} \omega \leq \theta_\delta(f_{\delta'}(\theta_\delta^{-1}(\varphi_\delta))) + \delta^{\frac{1}{2}}\psi_\delta$. Since $\|\psi_\delta\| \leq \|\omega\| + 2$ is bounded and $\theta_\delta(f_{\delta'}(\theta_\delta^{-1}(\varphi_\delta))) \in F^*$ is also bounded for fixed δ' , by considering the limit along a cofinal ultrafilter on the set of δ , we have $(1 + \delta')^{-1} \omega \in F^*$, so $\delta' \rightarrow 0$ gives $\omega \in F^*$.

To show G^* is weakly* closed, we claim for any $r > 0$ that

$$\overline{(F^* - A^{*+}) \cap A_{2r}^*} \subset G^*, \quad G^* \cap A_r^* \subset \overline{(F^* - A^{*+}) \cap A_{2r}^*},$$

where $A_r^* := \{\omega \in A^* : \|\omega\| \leq r\}$. If these are true, then

$$G^* \cap A_r^* = \overline{(F^* - A^{*+}) \cap A_{2r}^*} \cap A_r^*$$

is weakly* closed and convex in A^* for all $r > 0$, so the Krein-Šmulian theorem shows that G^* is weakly* closed.

Let $\omega_i \in (F^* - A^{*+}) \cap A_{2r}^*$ be a net such that $\omega_i \rightarrow \omega$ weakly* in A^* with $\|\omega_i\| \leq 2r$ for all i . Take $\varphi_i \in F^*$ such that $\omega_i \leq \varphi_i$ for all i , and define

$$\psi_i := \frac{[\omega_i]}{1 + \|\omega_i\|} + \frac{\varphi_i}{(1 + \|\omega_i\|)(1 + \|\varphi_i\|)}, \quad \varphi_{i,\delta} := \theta_i(f_\delta(\theta_i^{-1}(\varphi_i))),$$

where θ_i denotes the commutant Radon-Nikodym map associated to ψ_i . It clearly follows that $\|\psi_i\| \leq 1$ and $\|\varphi_{i,\delta}\| \leq \delta^{-1}$ for all i , and for sufficiently small $\delta > 0$ such that $1 + 2r \leq 2^{-1}\delta^{-\frac{1}{4}}$ and $\delta \leq 1$, we can also check

$$\omega_i = \theta_i(\theta_i^{-1}(\omega_i)) \leq \theta_i(f_\delta(\theta_i^{-1}(\omega_i)) + \delta^{\frac{1}{2}}) \leq \theta_i(f_\delta(\theta_i^{-1}(\varphi_i))) + \theta_i(\delta^{\frac{1}{2}}) = \varphi_{i,\delta} + \delta^{\frac{1}{2}}\psi_i.$$

Taking the limit along a cofinal ultrafilter on the index set of i , we obtain ψ and φ_δ for each δ by the limit of bounded nets ψ_i and $\varphi_{i,\delta}$ respectively, which satisfy $\omega \leq \varphi_\delta + \delta^{\frac{1}{2}}\psi$ with norm conditions $\|\psi\| \leq 1$ and $\|\varphi_\delta\| \leq \delta^{-1}$, so $\omega \in G^*$.

Let $\omega \in G^* \cap A_r^*$. Take $\psi \in A^{*+}$ and $\varphi_\delta \in F^*$ such that $\|\psi\| \leq 1$, $\|\varphi_\delta\| \leq \delta^{-1}$, and $\omega \leq \varphi_\delta + \delta^{\frac{1}{2}}\psi$, for any sufficiently small $\delta > 0$. If $\delta^{\frac{1}{2}} < r$, then $\omega - \delta^{\frac{1}{2}}\psi \in (F^* - A^{*+}) \cap A_{2r}^*$ converges to ω weakly* in A^* as $\delta \rightarrow 0$, we have $\omega \in \overline{(F^* - A^{*+}) \cap A_{2r}^*}$. This completes the proof. $\square$

(When A is non-commutative, $\omega_n \leq \varphi_n$ and $0 \leq \varphi_n$ do not imply $\omega_{n+} \leq \varphi_n$ because the ramp function is way far from operator monotone functions.)