

# **Calculus III**

**Calculus Early Transcendental 6th edition**

**Dr. Muath Karaki**



# An-Najah National University

## Department of Mathematics

### Calculus III

4C's Rule:

- Communication
- Collaboration
- Creativity
- Critical Thinking

**Muath Karaki**

Email: muath.karaki@najah.edu  
Website: [staff.najah.edu/en/profiles/2704](http://staff.najah.edu/en/profiles/2704)  
Office: 14F1460  
Office hours: See zajel

# Contents

I

## Part One

<b>10</b>	<b>Parametric Equation and Polar Coordinate .....</b>	<b>7</b>
10.1	Parametric Equation	7
10.2	Calculus with Parametric Equations.	11
10.3	Polar Coordinates.	16
10.4	Area and length	26
<b>12</b>	<b>Vectors and Geometry of Space .....</b>	<b>31</b>
12.1	3 Dimensional coordinate system	31
12.2	Vectors	34
12.3	The Dot Product	35
12.4	The Cross Product	40
12.5	Equation of linear & planes	45
<b>13</b>	<b>Vector Function .....</b>	<b>59</b>
13.1	Vector Functions and space curves	59
13.2	13.2	61
13.3	Arc Length	62

<b>14 Partial Derivatives .....</b>	<b>65</b>
14.1 Function of several variables	65
14.2 Partial Derivatives	69
14.3 Tangent Plane and Linear Approximation	72
14.4 Directional derivatives and Gradient vector	74
14.5 The Chain Rule	78

# Part One

## 10 Parametric Equation and Polar Coordinate ..... 7

- 10.1 Parametric Equation
- 10.2 Calculus with Parametric Equations.
- 10.3 Polar Coordinates.
- 10.4 Area and length

## 12 Vectors and Geometry of Space 31

- 12.1 3 Dimensional coordinate system
- 12.2 Vectors
- 12.3 The Dot Product
- 12.4 The Cross Product
- 12.5 Equation of linear & planes

## 13 Vector Function ..... 59

- 13.1 Vector Functions and space curves
- 13.2 13.2
- 13.3 Arc Length

## 14 Partial Derivatives ..... 65

- 14.1 Function of several variables
- 14.2 Partial Derivatives
- 14.3 Tangent Plane and Linear Approximation
- 14.4 Directional derivatives and Gradient vector
- 14.5 The Chain Rule




# 10. Parametric Equation and Polar Coordinate

## 10.1 Parametric Equation

$$x = 1 \Rightarrow y = 1$$

$$x = ? \Rightarrow y = ?$$

$$y = f(x)$$



$$y = t + 1$$

$$x = 1 - t^2$$

$$t = 1 \Rightarrow$$

$$x = 0, y = 2, (0, 2)$$

■ **Example 10.1** Sketch the curve define by the **Parametric equation**

$$x = t^2 - 2t, y = t + 1.$$

Solution

t	x	y
2	0	3
1	-1	2
0	0	1
-1	3	0
-2	8	-1
⋮	⋮	⋮


$$t = y - 1$$

⇓

$$x = (y - 1)^2 - 2(y - 1)$$

$$= y^2 - 2y + 1 - 2y + 2$$

$$x = y^2 - 4y + 2$$



■ **Example 10.2** Sketch  $x = t^2 - 2t$ ,  $y = t + 1$ ,  $-1 \leq x \leq 2$ .

[Solution](#)


*Parametric equation :*

$$x = f(t), y = g(t)$$

$$a \leq t \leq b$$

*Initial point*  $(f(a), g(a))$

*Terminal point*  $(f(b), g(b))$



■ **Example 10.3** Sketch  $x = \cos t$ ,  $y = \sin t$


[Solution](#)

$$\cos^2 t + \sin^2 t = 1$$

$$x^2 + y^2 = 1$$

$$t = 0 \Rightarrow (1, 0)$$

$$t = \frac{\pi}{2} \Rightarrow (0, 1)$$



■ **Example 10.4** Sketch  $x = \cos 2t$ ,  $y = \sin 2t$


[Solution](#)

$$\cos^2 2t + \sin^2 2t = 1$$

$$x^2 + y^2 = 1$$

$$t = 0 \Rightarrow (0, 0)$$


$$t = \frac{\pi}{2} \Rightarrow (-1, 0)$$



**Exercise 10.1** Sketch  $x = \cos \frac{1}{2}t$ ,  $y = \sin \frac{1}{2}t$


■ **Example 10.5** Sketch  $x = 2 \cos t$ ,  $y = 2 \sin t$   
Solution

$$\begin{aligned}x^2 + y^2 &= 4 \cos^2 t + 4 \sin^2 t \\x^2 + y^2 &= 4\end{aligned}$$




■ **Example 10.6** Sketch  $x = 2 \cos t + 1$ ,  $y = 2 \sin t + 2$   
Solution

$$\begin{aligned}x &= 2 \cos t + 1 \Rightarrow x - 1 = 2 \cos t \\y &= 2 \sin t + 2 \Rightarrow y - 2 = 2 \sin t \\(x - 1)^2 + (y - 2)^2 &= 4(\cos^2 t + \sin^2 t) \\(x - 1)^2 + (y - 2)^2 &= 4\end{aligned}$$



■ **Example 10.7** Sketch  $x = 2 \cos t$ ,  $y = 3 \sin t$   
Solution

$$\begin{aligned}x &= 2 \cos t \Rightarrow \cos t = \frac{x}{2} \\y &= 3 \sin t \Rightarrow \sin t = \frac{y}{3} \\\Rightarrow \left(\frac{x}{2}\right)^2 + \left(\frac{y}{3}\right)^2 &= 1\end{aligned}$$



■ **Example 10.8** Sketch  $x = \sin t$ ,  $y = \sin^2 t$

Solution

$$y = x^2$$


$$t = 0 \Rightarrow (0,0)$$

$$t = \frac{\pi}{2} \Rightarrow (1,1)$$

$$t = \pi \Rightarrow (0,0)$$

$$t = \frac{3\pi}{2} \Rightarrow (-1,1)$$

$$t = 2\pi \Rightarrow (0,0)$$

**Example 10.9**

$$\boxed{\frac{37}{628}}$$

$$(a) x = t^3, y = t^2, \Rightarrow t = x^{\frac{1}{3}} \Rightarrow y = x^{\frac{2}{3}}$$

$$(b) x = t^6, y = t^4, t = x^{\frac{1}{6}} \Rightarrow y = x^{\frac{4}{6}} \Rightarrow y = x^{\frac{2}{3}}$$

$$(c) x = e^{-3t}, y = e^{-2t}, \Rightarrow x = (e^{-t})^3 \Rightarrow e^{-t} = x^{\frac{1}{3}} \Rightarrow y = (e^{-t})^2 = (x^{\frac{1}{3}})^2 \Rightarrow y = x^{\frac{2}{3}}$$

Solution

$$(a) x = t^6$$

$$y = t^4$$

$$y = x^{\frac{2}{3}}$$

$$(b) x = t^6$$


$$y = t^4$$

$$y = x^{\frac{2}{3}}$$

$$(c) x = e^{-3t} = \frac{1}{e^{3t}}$$

$$y = e^{-2t} = \frac{1}{e^{2t}}$$


$$y = x^{\frac{2}{3}}$$



$$t = -1 \Rightarrow (-1,1)$$

$$t = 0 \Rightarrow (0,0)$$


$$t = 1 \Rightarrow (1,1)$$



$$t = -1 \Rightarrow (1,1)$$

$$t = 0 \Rightarrow (0,0)$$


$$t = 1 \Rightarrow (1,1)$$



$$t = -1 \Rightarrow (e^3, e^2)$$

$$t = 0 \Rightarrow (1,1)$$

$$t = 1 \Rightarrow \left(\frac{1}{e^3}, \frac{1}{e^2}\right)$$



**Problem 10.1** 1, 3, 4, 7, 8, 10, 12, 15, 16, 25, 26, 37.

## 10.2 Calculus with Parametric Equations.

let  $x = f(t), y = g(t)$  Then

$$\frac{dy}{dx} = \frac{\frac{d(y)}{d(t)}}{\frac{d(x)}{d(t)}} = \frac{g'(t)}{f'(t)}, f'(t) \neq 0$$

■ **Example 10.10** if  $x = \cos t, y = \sin t$

Solution

$$\frac{dy}{dx} = -\frac{\cos t}{\sin t} = -\cot t$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} \frac{dy}{dx} = \frac{\frac{d}{dt} \frac{dy}{dx}}{\frac{dx}{dt}} \\ &= \frac{\csc^2 t}{-\sin t} \\ &= -\csc^3 t \end{aligned} \tag{10.1}$$

■ **Example 10.11** if  $x = t^3, y = 3t$ , Find  $\frac{d^2y}{dx^2}$

Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{3}{3t^2} = t^{-2} & y &= 3x^{\frac{1}{3}} \\ \frac{d^2y}{dx^2} &= \frac{d}{dt} \frac{dy}{dx} & \frac{dy}{dx} &= x^{-\frac{2}{3}} \\ &= \frac{-2t^{-3}}{3t^2} = \frac{-2}{3}t^{-5} & \frac{d^2y}{dx^2} &= \frac{-2}{3}x^{-\frac{5}{3}} \\ \frac{d^2y}{dx^2} &= \frac{-2}{3}x^{-\frac{5}{3}} & \downarrow & \end{aligned}$$

**Exercise 10.2** if  $y = \cos t + t$ ,

$$x = 1 - t + t^2 \quad \text{Find } \frac{d^2y}{dx^2}$$

■ **Example 10.12** Consider the parametric equations :  $x = t^2, y = t^3 - 3t$

- (a) Show that the curve has two tangents at  $(3,0)$ , Find the tangent line
- (b) Find where the tangent is horizontal ? Vertical ?
- (c) Find where the curve is concave up? down ?
- (d) Sketch the curve .

#### Solution

$$(a) (3,0) \Rightarrow x = 3, y = 0$$

$$t^2 = 3 \quad t^3 - 3t = 0$$

↓

$$t = \sqrt{3}, -\sqrt{3}$$

↓

$$t = 0, \sqrt{3}, -\sqrt{3}$$

Refuse  $t=0$

The curve passes the point  $(3,0)$  two times at  $t = \sqrt{3}, -\sqrt{3}$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{3t^2 - 3}{2t}$$

$$\text{at } t = \sqrt{3} \Rightarrow \frac{dy}{dx} = \frac{3\sqrt{3}^2 - 3}{2\sqrt{3}} = \frac{3}{\sqrt{3}} = \sqrt{3}$$

$$\text{at } t = -\sqrt{3} \Rightarrow \frac{dy}{dx} = \frac{3 * 3 - 3}{-2\sqrt{3}} = -\sqrt{3}$$

So equation (1)

$$y - 0 = \sqrt{3}(x - 3)$$

And equation (2)

$$y - 0 = -\sqrt{3}(x - 3)$$

---


**Area:** If  $y = g(t), x = f(t)$

- Then the area between the curve  $C$  and  $x - \text{axis}$  is

$$A = \int_a^b g(t)f'(t)dt.$$

- The area with the  $y - \text{axis}$  is

$$A = \int_a^b f(t)g'(t)dt.$$




■ **Example 10.13** Find the area under one arc of cycloid

$$x = r(\theta - \sin \theta), y = r(1 - \cos \theta)$$

Solution


$$\begin{aligned} A &= \int_0^{2\pi} g(\theta)f'(\theta)d\theta = \int_0^{2\pi} r^2(1 - \cos \theta)^2 d\theta \\ &= r^2 \int_0^{2\pi} 1 - 2\cos \theta + \cos^2 \theta d\theta \\ &= r^2 \int_0^{2\pi} 1 - 2\cos \theta + \frac{1}{2} + \frac{1}{2}\cos 2\theta d\theta \\ &= r^2 [\theta - 2\sin \theta + \frac{\theta}{2} + \frac{1}{4}\sin 2\theta] \Big|_0^{2\pi} = [3\pi r^2] \end{aligned}$$



**Exercise 10.3** Q32 page 637 Find the area enclosed by the curve  $x = t^2 - 2t$ ,  $y = \sqrt{t}$  and the  $y - \text{axis}$

Solution

$$\begin{aligned} A &= \int_0^2 f(t)g'(t)dt \\ &= \int_0^2 (t^2 - 2t) \frac{1}{2}t^{-\frac{1}{2}} dt \\ &\vdots \end{aligned}$$



$$\text{Let } x = 0 \Rightarrow$$

$$t^2 - 2t = 0$$

$$t = 0, t = 2$$

**Arc Length:** Let  $x = f(t)$ ,  $y = g(t)$

$$L = \int_a^b \sqrt{f'^2(t) + g'^2(t)} dt$$




the curve should be traversed once in  $a \leq t \leq b$

■ **Example 10.14** Find the length of  $x = \cos \theta$ ,  $y = \sin \theta$ ,  $0 \leq \theta \leq 2\pi$

[Solution](#)

$$L = \int_0^{2\pi} \sqrt{\cos^2 \theta + \sin^2 \theta} d\theta$$

$$\int_0^{2\pi} 1 = 2\pi$$



■ **Example 10.15** Find the length of one arc of the cycloid

$$x = r(\theta - \sin \theta), y = r(1 - \cos \theta).$$

[Solution](#)

$$L = \int_0^{2\pi} \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta$$

$$= \int_0^{2\pi} \sqrt{r^2(1 - \cos \theta)^2 + r^2(\sin \theta)^2} d\theta$$

$$= r \int_0^{2\pi} \sqrt{1 - 2\cos \theta + \cos^2 \theta + \sin^2 \theta} d\theta$$

$$= r \int_0^{2\pi} \sqrt{2 - 2\cos \theta} d\theta$$

$$= r \int_0^{2\pi} \sqrt{4\sin^2 \frac{\theta}{2}} d\theta$$

$$= 2r \int_0^{2\pi} \left| \sin \frac{\theta}{2} \right| d\theta$$

$$= 2r \int_0^{2\pi} \sin \frac{\theta}{2}$$

$$2 - 2\cos \theta$$


$$= 2(1 - \cos \theta)$$

$$= 4\left(\frac{1}{2} - \frac{1}{2}\cos \theta\right)$$

$$= 4\left(\sin^2 \frac{1}{2}\theta\right)$$

$$2r(2) \cos \frac{\theta}{2} \Big|_0^{2\pi}$$

$$4r(1 + 1) = 8r$$



### **Surfaces Area:**

$$S = 2\pi \int_a^b g(t) \sqrt{f'^2(t) + g'^2(t)} dt \text{ about the } x-\text{axis}$$

$$\text{about the } x-\text{axis} = 2\pi \int_a^b y dL$$


$$S = 2\pi \int_a^b f(t) \sqrt{f'^2(t) + g'^2(t)} dt \text{ about the } y-\text{axis}$$

■ **Example 10.16** Find the area of the surface obtained by revolving one arc of the cycloid  $x = r(\theta - \sin \theta)$ ,  $y = r(1 - \cos \theta)$  about  $x-axis$

Solution:

$$S = 2\pi \int_0^{2\pi} r(1 - \cos \theta) \sqrt{2(1 - \cos \theta)} d\theta$$

⋮




■

■ **Example 10.17** Find the area of a sphere of radius  $r$   $S = 4\pi r^2$

Solution

$$\begin{aligned} S &= 2\pi \int_0^\pi r \sin \theta \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta \\ &= 2\pi r \int_0^\pi \sin \theta \sqrt{r^2} d\theta \\ &= 2\pi r^2 \int_0^\pi \sin \theta d\theta \\ &= 2\pi r^2 \cos \theta \Big|_0^\pi \\ &= 2\pi r^2(1 + 1) = 4\pi r^2 \end{aligned}$$



$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ 0 &\leq \theta \leq \pi \end{aligned}$$

■

---

**Problem 10.2** 1, 3, 5, 7, 11, 12, 13, 15, 17, 18, 19, 28, 29, 30, 33, 34, 37, 39, 40, 41, 43, 57, 59, 60, 65, 69.

**10.3 Polar Coordinates.**

.  $r = \sin(a\theta)$   $r = \cos(a\theta)$

.  $r = a + b \cos \theta$   $r = a + b \sin \theta$

---

■ **Example 10.18** Locate the following polar points .


1.  $(3, \frac{\pi}{4})$

2.  $(-3, \frac{\pi}{4})$

3.  $(1, \pi)$

4.  $(-1, 0)$

5.  $(1, -\pi)$

Solution

$$x = r\cos\theta \qquad r^2 = x^2 + y^2$$

$$y = r\sin\theta \qquad \theta = \tan^{-1} \frac{y}{x}$$

■ **Example 10.19** Find the cartesian coordinate for the following polar points. 1)  $(1, \frac{2\pi}{3})$

$$x = 1 * \cos \frac{2\pi}{3} = -\frac{1}{2}$$

$$y = 1 * \sin \frac{2\pi}{3} = \frac{\sqrt{3}}{2}$$

■ **Example 10.20** Convert from cartesian to polar


$$\text{Solution } (1, -\sqrt{3}) \quad r^2 = 1 + 3 = 4 \Rightarrow r = 2,$$

$$\theta = \tan^{-1} \sqrt{3}/1 \Rightarrow \theta = 300^\circ$$

**Polar Curves**

■ **Example 10.21** Sketch the following polar curves .

- $r = 2 \Rightarrow r^2 = 4 \Rightarrow x^2 + y^2 = 4$



- $\theta = \frac{\pi}{4}$

$$\tan \theta = \frac{y}{x}$$



$$1 = \frac{y}{x} \Rightarrow y = x$$

- $r = 2 * \cos \theta$


$$r^2 = 2 * r * \cos \theta \Rightarrow x^2 + y^2 = 2x$$

$$x^2 - 2x + ... + 1 + y^2 = 0 ... + 1$$


$$(x - 1)^2 + y^2 = 1$$



- $r = -4 * \cos \theta$



- $r = 2 * \sin \theta$




- $r = -4 * \sin \theta$  (H.W)
- $r = 2 * \sin \theta - 4 * \cos \theta$  (H.W)

- $r = 3 * \sec \theta$

$$r = \frac{3}{\cos \theta} \Rightarrow r * \cos \theta = 3$$

$$\Rightarrow x = 3$$





- $r = 2 * \sin \theta$

- $r = -3 * \csc \theta$  (H.W)

- $r = 1 + \sin \theta$

First sketch the equation in the cartesian system

$\theta$	$r$
$0 \rightarrow 2$	$1 \rightarrow 2$
$\frac{\pi}{2} \rightarrow \pi$	$2 \rightarrow 1$
$\pi \rightarrow \frac{2\pi}{3}$	$1 \rightarrow 0$
$\frac{2\pi}{3} \rightarrow \pi$	$0 \rightarrow 1$




- $r = 1 - \sin \theta$  (H.W)

- $r = 1 + \cos \theta$  (H.W)

- $r = 1 - \cos \theta$  (H.W)


- $r = 2 + 2\cos \theta$  (H.W)



- $r = 2 + \sin \theta$  (H.W)


$\theta$	$r$
$0 \rightarrow 2$	$2 \rightarrow 3$
$\frac{\pi}{2} \rightarrow \pi$	$3 \rightarrow 2$
$\pi \rightarrow \frac{2\pi}{3}$	$2 \rightarrow 1$
$\frac{2\pi}{3} \rightarrow \pi$	$1 \rightarrow 2$

- $r = 3 + 2\cos\theta$  (H.W)
- $r = 3 - 2\cos\theta$  (H.W)



■ **Example 10.22** Sketch  $r = 1 + 2\cos\theta$

Solution




$$r = 0 \Rightarrow$$

$$1 + 2\cos\theta = 0$$

$$\cos\theta = -\frac{1}{2}$$


$$\theta = \frac{2}{3}\pi, \frac{4}{3}\pi$$

$\theta$	$r$
$0 \rightarrow \frac{\pi}{2}$	$3 \rightarrow 1$
$\frac{\pi}{2} \rightarrow \frac{\pi}{3}$	$1 \rightarrow 0$
$\frac{2}{3}\pi \rightarrow \pi$	$0 \rightarrow -1$
$\pi \rightarrow \frac{4}{3}\pi$	$0 \rightarrow -1$
$\frac{4}{3}\pi \rightarrow \frac{3}{2}\pi$	$0 \rightarrow 1$
$\frac{3}{2}\pi \rightarrow 2\pi$	$1 \rightarrow 3$



$$r = a + b\sin\theta$$


■ **Example 10.23** Sketch  $r = -3 - 6\sin\theta$


Solution

■ **Example 10.24** Sketch  $r = \cos 2\theta$

Solution

$\theta$	$r$
$0 \rightarrow \frac{\pi}{4}$	$1 \rightarrow 0$
$\frac{\pi}{4} \rightarrow \frac{\pi}{2}$	$0 \rightarrow -1$
$\frac{\pi}{2} \rightarrow \frac{3\pi}{4}$	$-1 \rightarrow 0$
$\frac{3\pi}{4} \rightarrow \pi$	$0 \rightarrow 1$
$\pi \rightarrow \frac{5\pi}{4}$	$1 \rightarrow 0$
$\frac{5\pi}{4} \rightarrow \frac{3\pi}{2}$	$0 \rightarrow -1$
$\frac{3\pi}{2} \rightarrow \frac{7\pi}{4}$	$-1 \rightarrow 0$
$\frac{7\pi}{4} \rightarrow 2\pi$	$0 \rightarrow 1$





**Exercise 10.4** Sketch:

1.  $r = \sin 2\theta$
2.  $r = \cos 3\theta$
3.  $r = \cos 4\theta$


■ **Example 10.25** Sketch  $r^2 = \cos 2\theta$

Solution

$$r = \sqrt{\cos 2\theta}$$

$$r = -\sqrt{\cos 2\theta}$$


$\theta$	$r = \cos 2\theta$	$r^2 = \cos 2\theta$
$0 \rightarrow \frac{\pi}{4}$	$1 \rightarrow 0$	$1 \rightarrow 0$
$\frac{\pi}{4} \rightarrow \frac{\pi}{2}$	$0 \rightarrow -1$	$\times$
$\frac{\pi}{2} \rightarrow \frac{3\pi}{4}$	$-1 \rightarrow 0$	$\times$
$\frac{3\pi}{4} \rightarrow \pi$	$0 \rightarrow 1$	$0 \rightarrow 1$
$\pi \rightarrow \frac{5\pi}{4}$	$1 \rightarrow 0$	$1 \rightarrow 0$
$\frac{5\pi}{4} \rightarrow \frac{3\pi}{2}$	$0 \rightarrow -1$	$\times$
$\frac{3\pi}{2} \rightarrow \frac{7\pi}{4}$	$-1 \rightarrow 0$	$\times$
$\frac{7\pi}{4} \rightarrow 2\pi$	$0 \rightarrow 1$	$0 \rightarrow 1$



### Symmetry:

When we sketch polar curves, it is sometimes helpful to take advantage of symmetry .The following three rules are explained below :

1. If a polar equation is unchanged when  $\theta$  is replaced by  $-\theta$  , the curve is symmetric about the polar axis .
2. If the equation is unchanged when  $r$  is replaced by  $-r$  , or when  $\theta$  is replaced by  $\theta + \pi$  the curve is symmetric about the pole.(This means that the curve remains unchanged if we rotate it through  $180^\circ$  about the origin)
3. If the equation is unchanged when  $\theta$  is replaced by  $\pi - \theta$ , the curve is symmetric about the vertical line  $\theta = \frac{\pi}{2}$



■ **Example 10.26**  $r = \cos\theta$  is symmetric

1. about the polar axis .

2. about the origin .

3. about  $\theta = \frac{\pi}{2}$

■

Tangents in polar system  $\frac{dy}{dx}$

$$x = r\cos\theta \quad y = r\sin\theta$$

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{\frac{dr}{d\theta} * \sin\theta + r\cos\theta}{\frac{dr}{d\theta} * \cos\theta - r * \sin\theta}$$

■ **Example 10.27** Let  $r = 1 + \sin\theta$

1. Find the slope at  $\theta = \frac{\pi}{3}$

2. Find where the tangent is horizontal? Vertical ?

Solution:  $x = r\cos\theta \quad , y = r\sin\theta$

$$1. \frac{dy}{dx} = \frac{\frac{dr}{d\theta} * \sin\theta + r\cos\theta}{\frac{dr}{d\theta} * \cos\theta - r * \sin\theta}$$

$$\frac{dy}{d\theta} = \cos\theta$$

$$\cos\frac{\pi}{3} = \frac{1}{2}$$

$$\sin\frac{\pi}{3} = \frac{\sqrt{3}}{2}$$

$$\begin{aligned} \Rightarrow \frac{dy}{dx} &= \frac{\cos\frac{\pi}{3} * \sin\frac{\pi}{3} + \cos\frac{\pi}{3}(1 + \sin\frac{\pi}{3})}{\cos\frac{\pi}{3} * \cos\frac{\pi}{3} - (1 + \sin\frac{\pi}{3}) * \sin\frac{\pi}{3}} \\ &= \frac{\frac{\sqrt{3}}{2} * \frac{1}{2} + \frac{1}{2}(1 + \frac{\sqrt{3}}{2})}{\frac{1}{2} - \frac{\sqrt{3}}{2}} = -1 \end{aligned}$$

Equation of the tangent:

Slope=-1

$$x_0 = \left(1 + \sin\frac{\pi}{3} * \cos\frac{\pi}{3}\right) = \left(\frac{1}{2} + \frac{\sqrt{3}}{4}\right)$$

$$y_0 = \left(1 + \sin\frac{\pi}{3} * \sin\frac{\pi}{3}\right) = \left(1 + \frac{\sqrt{3}}{4}\right) * \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{2} + \frac{3}{4}$$

$$\text{Equation : } y - \frac{\sqrt{3}}{2} + \frac{3}{4} = -1(x - \left(\frac{1}{2} + \frac{\sqrt{3}}{4}\right))$$

$$2. \frac{dy}{dx} = \frac{\cos\theta * \sin\theta + (1 + \sin\theta) * \cos\theta}{\cos\theta * \cos\theta - (1 + \sin\theta) * \sin\theta}$$

$$\frac{2 * \cos\theta * \sin\theta + \cos\theta}{\cos^2\theta - \sin^2\theta - \sin\theta} = \frac{\sin 2\theta + \cos\theta}{\cos 2\theta - \sin\theta}$$

$$\frac{dy}{dx} = 0 \Rightarrow 2 * \cos\theta * \sin\theta + \cos\theta = 0 \Rightarrow \cos\theta(2\sin\theta + 1) = 0$$

$$\cos\theta = 0 \quad \text{or} \quad \sin\theta = -\frac{1}{2}$$

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2} \quad \theta = \frac{7\pi}{6}, \frac{11\pi}{6}$$

$$\Rightarrow \frac{dx}{d\theta} = 0 \Rightarrow \cos^2\theta - \sin^2\theta - \sin\theta = 0$$

$$1 - \sin^2\theta - \sin\theta - \sin\theta = 0$$

$$(2\sin\theta - 1)(\sin\theta + 1) = 0$$

$$\sin\theta = \frac{1}{2}, \quad \sin\theta = -1$$

$$\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$$

$$\text{H.T } \theta = \frac{7\pi}{6}, \frac{11\pi}{6}, \frac{\pi}{2}$$

$$\text{V.T } \theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$$

$$\text{at } \theta = \frac{3\pi}{2}$$


$$\lim_{\theta \rightarrow \frac{3\pi}{2}^+} \frac{dy}{dx} = \lim_{\theta \rightarrow \frac{3\pi}{2}^+} \frac{\sin 2\theta + \cos\theta}{\cos 2\theta - \sin\theta} = \left(\frac{0}{0}\right)$$

$$\text{L'Hopital} = \lim_{\theta \rightarrow \frac{3\pi}{2}^+} \frac{3\pi}{2} = \frac{2\cos 2\theta - \sin\theta}{2\sin 2\theta - \cos\theta} = -\infty$$

**Problem 10.3** 1, 3, 7, 8, 9, 11, 13, 15, 16, 17, 19, 21, 24, 25, 29, 30, 31, 34, 37, 39, 40, 42, 43, 47, 57, 58, 61, 63, 65, 67, 70.

## 10.4 Area and length

$$A = \frac{1}{2} \int_a^b r^2 d\theta$$




■ **Example 10.28** Find the area of one leaf of the rose  $r = \cos 2\theta$

$$A = 2 * \frac{1}{2} * \int_0^{\frac{\pi}{4}} (\cos 2\theta)^2 d\theta$$

Solution:  $A = \int_0^{\frac{\pi}{2}} \left( \frac{1}{2} + \frac{1}{2} * \cos 4\theta \right) d\theta$

$$A = \frac{\theta}{2} + \frac{1}{8} \sin 4\theta \Big|_0^{\frac{\pi}{2}}$$

$$A = \frac{\pi}{4}$$





■ **Example 10.29** Find the area of the region that lies inside  $r = 3 \sin \theta$  and outside  $r = 1 + \sin \theta$ .

$$r = f(\theta)$$

Solution:

$$A = \frac{1}{2} \int_a^b r^2 d\theta$$






$$A = \frac{1}{2} \int_{\pi}^{\frac{5\pi}{6}} (3 \sin \theta)^2 - (1 + \sin \theta)^2 d\theta$$

$$= 2 \cdot \frac{1}{2} \int_{\pi}^{\frac{\pi}{2}} (3 \sin \theta)^2 - (1 + \sin \theta)^2 d\theta$$

■ **Example 10.30** Find the area inside both  $r = 1 + \cos \theta$ ,  $r = 1 - \cos \theta$

Solution:



$$A = 4 \cdot \frac{1}{2} \int_0^{\frac{\pi}{2}} (1 - \cos \theta)^2 d\theta$$

$$\text{or } A = 4 \cdot \frac{1}{2} \int_{\frac{\pi}{2}}^{\pi} (1 - \cos \theta)^2 d\theta$$

■ **Example 10.31** Find the area inside the inner loop of  $r = 1 + 2 \cos \theta$

Solution:

$$r = 0$$

$$\Rightarrow 1 + 2\cos \theta = 0$$


$$\Rightarrow \cos \theta = -\frac{1}{2}$$

$$\Rightarrow \theta = \frac{2}{3}\pi, \frac{4}{3}\pi$$

$$A = 2 \cdot \frac{1}{2} \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} (1 + 2\cos \theta)^2 d\theta$$

$$= 2 \cdot \frac{1}{2} \int_{\pi}^{\frac{4}{3}\pi} (1 + 2\cos \theta)^2 d\theta$$

$$= \frac{1}{2} \int_{\frac{4}{3}\pi}^{\frac{2}{3}\pi} (1 + 2\cos \theta)^2 d\theta$$




■

- **Example 10.32** Find the area the lies between the inner and the outer loop of  $r = 1 + 2\cos \theta$ .

Solution:

$$A = 2 \cdot \frac{1}{2} \left( \int_0^{\frac{2}{3}\pi} (1 + 2\cos \theta)^2 d\theta - \int_{\frac{4}{3}\pi}^{\frac{2}{3}\pi} (1 + \cos \theta)^2 d\theta \right)$$




■

- **Example 10.33** Find the area... inside both  $r = \cos 2\theta$ ,  $r = \sin 2\theta$ .

Solution:

$$A = 8 \cdot 2 \cdot \frac{1}{2} \int_0^{\frac{\pi}{8}} (\sin 2\theta)^2 d\theta$$

$$\text{or } = 16 \cdot \frac{1}{2} \int_{\frac{\pi}{8}}^{\frac{\pi}{4}} (\cos 2\theta)^2 d\theta$$




---


**Problem 10.4** 2, 3, 5, 6, 7, 8, 9, 11, 13, 17, 21, 23, 24, 25, 26, 28, 29, 31, 33, 35, 37, 41, 45, 47.

■ **Example 10.34** Find the area of the loop  $r^2 = 9 \cos 2\theta$  Solution:

$$A = 2 * \frac{1}{2} \int_0^{\frac{\pi}{4}} 9 \cos 2\theta d\theta$$

$$A = \frac{9}{2} \sin 2\theta \Big|_0^{\frac{\pi}{4}}$$

$$A = \frac{9}{2}$$




---

### Parametric Equation

- $x = f(\theta) * \cos(\theta)$        $r = f(\theta)$
- $y = f(\theta) * \sin(\theta)$

---

### Arc Length :

$$L = \int_a^b \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta$$

$$= \int_a^b \sqrt{(r' * \cos(\theta) - r * \sin(\theta))^2 + (r' * \sin(\theta) + r * \cos(\theta))^2} d\theta$$

$$= \int_a^b \sqrt{(r')^2 * \cos(\theta)^2 - 2 * r' * r * \cos(\theta) * \sin(\theta) + (r')^2 * \sin(\theta)^2 + 2 * r' * r * \cos(\theta) * \sin(\theta)} d\theta$$

$$L = \int_a^b \sqrt{(r')^2 + r^2} d\theta$$

■ **Example 10.35** Find the length of the cardioid  $r = 1 + \sin \theta$  Solution:

$$\begin{aligned}
L &= \int_0^{2\pi} \sqrt{(r')^2 + r^2} d\theta \\
&= \int_0^{2\pi} \sqrt{(1 + \sin \theta)^2 + \cos^2(\theta)} d\theta \\
&= \int_0^{2\pi} \sqrt{1 + 2 \sin \theta + \sin^2 \theta + \cos^2(\theta)} d\theta \\
&= \int_0^{2\pi} \sqrt{2 + 2 \sin \theta} d\theta * \sqrt{\frac{2 - 2 \sin \theta}{2 - 2 \sin \theta}} \\
&= \int_0^{2\pi} \sqrt{\frac{4(1 - \sin^2 \theta)}{2(1 - \sin \theta)}} \\
&= \int_0^{2\pi} \frac{2\sqrt{\cos^2 \theta}}{\sqrt{2(1 - \sin \theta)}} \\
&= \int_0^{2\pi} \frac{2|\cos \theta|}{\sqrt{2(1 - \sin \theta)}}
\end{aligned}$$


- 
- **Example 10.36** 1.  $\delta = \int_a^b 2\pi r * \sin \theta * \sqrt{(r')^2 + r^2} d\theta$  about the polar axis.  
 2. Find the surface area generated by rotating the lemniscate  $r^2 = \cos 2\theta$  about the polar axis .

Solution:


$$\begin{aligned}
\delta &= 2 \int_0^{\frac{\pi}{4}} 2\pi r * \sin \theta * \sqrt{(r')^2 + r^2} d\theta \\
&= 4\pi \int_0^{\frac{\pi}{4}} r * \sin \theta * \sqrt{(\cos 2\theta) + \frac{\sin^2(2\theta)}{\cos(2\theta)}} d\theta \\
&= 4\pi \int_0^{\frac{\pi}{4}} r * \sin \theta * \sqrt{\frac{1}{\cos(2\theta)}} d\theta \\
&= 4\pi \int_0^{\frac{\pi}{4}} r * \sin \theta * \frac{1}{r} d\theta \\
&= 4\pi * \cos \theta * (0 - \frac{\pi}{4}) \\
&= 4\pi * (1 - \frac{1}{\sqrt{2}})
\end{aligned}$$

## 12. Vectors and Geometry of Space

### 12.1 3 Dimensional coordinate system




- $XZ - plane$
- $XY - plane$
- $YZ - plane$




■ **Example 12.1** Describe the following equations :

1. (a)  $y = 3$  in  $2D(R^2)$   
 (b)  $y = 3$  in  $3D(R^3)$  (Plane parallel  $xz$ -plane)
2.  $z^2 = 1$  in  $R^3$ 
  - $z=1$  plane above parallel  $xy$ -plane
  - $z=-1$  plane below parallel  $xy$ -plane
3.  $x^2 + y^2 = 1 \ \& z = 2$
4.  $y = x$  in  $R^3$ 
  - plane  $\frac{\pi}{4}$  with  $XZ$ -plane ,  $YZ$ -plane





Solution:




1. (a)



(b)



### 3D Coordinate System



- $(x_1, y_1, z_1)$ ,  $(x_2, y_2, z_2)$ , Distance:

$$D = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

- $(h, k, l)$ ,  $r$

$$(x - h)^2 + (y - k)^2 + (z - l)^2 = r^2$$

*Sphere center  $(h, k, l)$*

*radius r*

---

- **Example 12.2** Describe the region represented by:  $1 \leq x^2 + y^2 + z^2 \leq 4$

Solution

The equation represented the region between:

The sphere of center (0,0,0), radios 2

and the sphere of center (0,0,0), radios 1


■

- **Example 12.3** Find an equation of the largest sphere of center (5,4,9) in the first octant.

Solution

$$r = 4$$

$$(x - 5)^2 + (y - 4)^2 + (z - 9)^2 = 16$$




- **Example 12.4** Find the distance between the point (2,3,-4) and :

1. the  $x-axis$

$$d = \sqrt{3^2 + 4^2} = 5$$

2. the  $xz-plane$


$$d = 3$$




---

**Problem 12.1** 2,3,7,9,10,11,13,14,15,17,19,20,21,23,25,29,32,33,35,38

## 12.2 Vectors



$$\vec{v} = \overrightarrow{PQ} = \langle x_2 - x_1, y_2 - y_1 \rangle$$

in 2D  $\vec{d} = \langle x, y \rangle$

$$|\vec{d}| = \sqrt{x^2 + y^2}$$

in 3D  $\vec{d} = \langle x, y, z \rangle$

$$|\vec{d}| = \sqrt{x^2 + y^2 + z^2}$$

if  $\vec{d} = \langle a_1, a_2, a_3 \rangle, \vec{b} = \langle b_1, b_2, b_3 \rangle \Rightarrow$

$$\begin{aligned}\vec{d} + \vec{b} &= \langle a_1 + b_1, a_2 + b_2, a_3 + b_3 \rangle \\ c\vec{d} &= \langle ca_1, ca_2, ca_3 \rangle\end{aligned}$$

$$\begin{aligned}\vec{d} &= \langle a_1, a_2, a_3 \rangle = a_1 \underbrace{\langle 1, 0, 0 \rangle}_i + a_2 \underbrace{\langle 0, 1, 0 \rangle}_j + a_3 \underbrace{\langle 0, 0, 1 \rangle}_k \\ &= a_1 i + a_2 j + a_3 k\end{aligned}$$

**Definition 12.2.1 — Unit vector:** a vector  $\vec{u}$  is called a unit if  $|\vec{u}| = 1$

■ **Example 12.5** Find a unit vector in the opposite direction of  $\vec{v} = \langle 2, -2, 1 \rangle$

•

Solution:  $|\vec{v}| = \frac{\vec{v}}{|\vec{v}|} = \frac{-1}{\sqrt{4+4+1}} \langle 2, -2, 1 \rangle = \langle \frac{-2}{3}, \frac{2}{3}, \frac{-1}{3} \rangle$

▪

**Problem 12.2** 7,11,13,15,17,19,21,23,24,25,37,41,42

### 12.3 The Dot Product

**Definition 12.3.1** if  $\vec{d} = \langle a_1, a_2, a_3 \rangle, \vec{b} = \langle b_1, b_2, b_3 \rangle \Rightarrow$

$$\vec{d} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

■ **Example 12.6** if  $\vec{d} = \langle 2, 1, -2 \rangle, \vec{b} = \langle 1, 1, 3 \rangle$

Solution

$$\vec{d} \cdot \vec{b} = 2 + 1 - 6 = -3$$

$$\vec{d} \cdot \vec{d} = a_1^2 + a_2^2 + a_3^2 = (\sqrt{a_1^2 + a_2^2 + a_3^2})^2 = |\vec{d}|^2$$

**Properties:**

1.  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$
2.  $\vec{a} \cdot \vec{a} = |\vec{a}|^2$
3.  $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$
4.  $(c\vec{a}) \cdot \vec{b} = \vec{a} \cdot (c\vec{b}) = c(\vec{a} \cdot \vec{b})$
5.  $\vec{0} \cdot \vec{a} = 0$

**Theorem 12.3.1** if  $\theta$  is the angle between  $\vec{a}$  &  $\vec{b}$  then  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$   
 $\theta \in [0, \pi]$

**Corollary 1:**  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$   $|\vec{a}| \neq 0, |\vec{b}| \neq 0$

■ **Example 12.7** Find the angle between  $\vec{a} = \langle 2, 2, -1 \rangle$ ,  $\vec{b} = \langle 5, -3, -2 \rangle$

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{10 - 6 - 2}{3\sqrt{25+9+4}} = \frac{2}{3\sqrt{38}}$$

$$g = \cos^{-1}\left(\frac{2}{3\sqrt{38}}\right) \approx 1.46(84)^\circ$$

**Corollary 2:** Two non-zero vectors are orthogonal iff  $\vec{a} \cdot \vec{b} = 0$

■ **Example 12.8** if  $\vec{a} = \langle 2, 1, -2 \rangle$ ,  $\vec{b} = \langle c, 2, 1 \rangle$

Find  $c$  such that  $\vec{a} \perp \vec{b} = 0$

$$\vec{a} \cdot \vec{b} = 0 \Leftrightarrow 2c + 2 - 2 = 0 \Leftrightarrow c = 0$$

$$\vec{a} = \langle a_1, a_2, a_3 \rangle \Rightarrow$$

$$\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

$$\vec{b} = \langle b_1, b_2, b_3 \rangle \quad \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta \quad \vec{a} \cdot \vec{b} =$$

$$0 \Leftrightarrow \vec{a} \perp \vec{b}$$

- $-|a||b| \leq \vec{a} \cdot \vec{b} \leq |a||b|$

$$|\vec{a} \cdot \vec{b}| \leq |a||b|$$

■ **Example 12.9** if  $|\vec{a}| = 2$ ,  $|\vec{b}| = 3$ ,  $\theta = \frac{2}{3}\pi$ . Find  $|\vec{a} - 2\vec{b}|$ .  $|\vec{u}|^2 = \vec{u} \cdot \vec{u}$

$$|\vec{a} - 2\vec{b}|^2 = (\vec{a} - 2\vec{b}) \cdot (\vec{a} - 2\vec{b})$$

$$= \vec{a} \cdot \vec{a} - 2\vec{a} \cdot \vec{b} - 2\vec{b} \cdot \vec{a} + 4\vec{b} \cdot \vec{b}$$

$$= |\vec{a}|^2 - 4|\vec{a}||\vec{b}| \cos \theta + 4|\vec{b}|^2$$

$$= 4 - 4 \cdot 2 \cdot 3 \left(\frac{-1}{2}\right) + 4 \cdot 9$$

$$= 16 + 36 = 52 \quad |\vec{a} - 2\vec{b}| = \sqrt{52}$$

$$|\vec{a} + \vec{b}| = \sqrt{|a|^2 + |b|^2 + 2|a||b| \cos \theta}$$

■ **Example 12.10** Prove that  $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$ .

Pf:

$$|\vec{a} + \vec{b}|^2 = (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b})$$

$$\begin{aligned}
 &= |\vec{a}|^2 + |\vec{b}|^2 + 2 \vec{a} \cdot \vec{b} \leq |\vec{a}|^2 + |\vec{b}|^2 + 2 |a| |b| \\
 &\quad = (\underbrace{|a| + |b|}_{\gamma})^2 \\
 \Rightarrow |\vec{a} + \vec{b}| &\leq |\vec{a}| + |\vec{b}|
 \end{aligned}$$

Direction Angles.  
is the angle b/w  $\vec{v}$  & the  $x-axis$   $\beta$   
 $\gamma$

Direction Cosines.  $\cos\alpha = \frac{a_1}{|\vec{v}|}$   
 $\cos\beta = \frac{a_2}{|\vec{v}|}$   $\cos\gamma = \frac{a_3}{|\vec{v}|}$

$$\begin{aligned}
 a_1 &= \cos\alpha |\vec{v}|, a_2 = \cos\beta |\vec{v}|, a_3 = \cos\gamma |\vec{v}| \\
 \vec{v} &= |\vec{v}| \langle \cos\alpha, \cos\beta, \cos\gamma \rangle \\
 \frac{\vec{v}}{|\vec{v}|} &= \langle \cos\alpha, \cos\beta, \cos\gamma \rangle \\
 \Rightarrow \cos^2\alpha + \cos^2\beta + \cos^2\gamma &= 1
 \end{aligned}$$

■ **Example 12.11** can  $\alpha = \frac{\pi}{4}$ ,  $\beta = \frac{\pi}{4}$ ,  $\gamma = \frac{\pi}{4}$  be direction cosines. Answer.  
No, because

$$\cos^2\frac{\pi}{4} + \cos^2\frac{\pi}{4} + \cos^2\frac{\pi}{4} = \frac{3}{2} \neq 1$$

■ **Exercise 12.1** if  $\vec{a} = \langle 1, 2, 3 \rangle$  Find direction angles.

### Projections

- Scalar Projection:

$$\begin{aligned}
 \text{comp}_{\vec{a}} \vec{b} &= \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} & \frac{|\vec{a}| |\vec{b}| \cos\theta}{|\vec{a}|} &= \mathcal{L} & \vec{v} &= \mathcal{L} \frac{\vec{a}}{|\vec{a}|} \\
 && \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} &= & &= \frac{\vec{a} \cdot \vec{a}}{|\vec{a}|^2} \vec{a}
 \end{aligned}$$

- Vector Projection:

$$\text{Proj}_{\vec{a}} \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \vec{a}$$

■ **Example 12.12** if  $\vec{a} = \langle 2, 1, -1 \rangle$

Find

$$\vec{b} = \langle 2, -1, 2 \rangle$$

$$1. \ comp_{\vec{b}}^{\vec{a}} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{4 - 1 - 2}{3} = \frac{1}{3}$$

$$2. \ Proj_{\vec{b}}^{\vec{a}} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = \frac{1}{9} \langle 2, -1, 2 \rangle$$

■

■ **Example 12.13** if  $Proj_{\vec{b}}^{\vec{a}} = \langle 2, 1, -2 \rangle$ ,  $\underbrace{\theta}_{\text{angle b/w } \vec{a} \text{ & } \vec{b}} = \frac{8}{15}\pi$

Find:

$$1. \ Proj_{\vec{b}}^{2\vec{a}} = 2 \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = 2 \langle 2, 1, -2 \rangle = \langle 4, 2, -4 \rangle$$

$$2. \ Proj_{2\vec{b}}^{\vec{a}} = \frac{\vec{a} \cdot 2\vec{b}}{4|\vec{b}|} |2\vec{b}| = \langle 2, 1, -2 \rangle$$

$$3. \ Proj_{-2\vec{b}}^{\vec{a}} = \frac{\vec{a} \cdot (-2\vec{b})}{4|\vec{b}|} (-2\vec{b}) = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} = \langle 2, 1, -2 \rangle$$

$$4. \ comp_{\vec{b}}^{\vec{a}} = -3$$

■

■ **Example 12.14** Find  $x$  such that the angle between  $\langle 2, 1, -1 \rangle$ ,  $\langle 1, x, 0 \rangle$  is  $45^\circ$

$$\cos 45^\circ = \frac{2+x-0}{\sqrt{6}\sqrt{1+x^2}} \quad |\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}|$$

$$\Leftrightarrow \frac{\sqrt{6}}{\sqrt{2}} \sqrt{1+x^2} = 2+x$$

$$\Rightarrow 3(1+x^2) = (2+x)^2$$

$$3+3x^2-4x-1=0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

■

■ **Example 12.15** Find the two unit vectors that make an angle  $60^\circ$  with  $\vec{v} = \langle 3, 4 \rangle$ .


$$\vec{u} = \langle a, b \rangle$$

$$a^2 + b^2 = 1 \cdots (1)$$

$$\cos 60^\circ = \frac{\vec{u} \cdot \vec{v}}{|\vec{u}| \cdot |\vec{v}|}$$

$$\frac{1}{2} = \frac{3a+4b}{1.5}$$


$$3a+4b = \frac{5}{2} \cdots (2)$$



$$\theta^\circ = \tan \frac{4}{3} = x_0$$

■ 53: the distance between the fine  $ax + by + c = 0$  and the point  $(x_1, y_1)$  is

$$D = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$




**Question:** Find the distance between  $(-1, 3)$  & the line  $3x - 4y + 5 = 0$

$$D = \frac{|3(-1) - 4(3) + 5|}{\sqrt{3^2 + 4^2}} = \frac{13}{5}$$

■ **Example 12.16** if  $\vec{c} = |\vec{d}| \vec{b} + |\vec{b}| \vec{d}$ ,  $\vec{d}, \vec{b}, \vec{c}$  not zero vectors  
Show that  $\vec{0}$  bisects  $\vec{d}$  &  $\vec{b}$

Solution:

$$\begin{aligned}\cos \alpha &= \frac{\vec{d} \cdot \vec{c}}{|\vec{d}| |\vec{c}|} = \frac{\vec{d} \cdot [|\vec{d}| \vec{b} + |\vec{b}| \vec{d}]}{|\vec{d}| |\vec{c}|} \\ &= \frac{\vec{d} \cdot \vec{b} + |\vec{d}| |\vec{b}|}{|\vec{c}|}\end{aligned}$$



$$\begin{aligned}\cos \beta &= \frac{\vec{b} \cdot \vec{c}}{|\vec{b}| |\vec{c}|} = \frac{\vec{b} \cdot [|\vec{d}| \vec{b} + |\vec{b}| \vec{d}]}{|\vec{b}| |\vec{c}|} \\ &= \frac{\vec{d} \cdot \vec{b} + |\vec{d}| |\vec{b}|}{|\vec{c}|} \\ \Rightarrow \alpha &= \beta\end{aligned}$$

**Exercise 12.2** Show that  $\text{proj}_{\vec{b}} \vec{d} \cdot \text{proj}_{\vec{d}} \vec{b} = (\vec{d} \cdot \vec{b}) \cos^2 \theta$

**Problem 12.3** 1,3,7,9,10,11,15,19,20,21,23,25,26,27,31,35,39,43,45,49,54,59

## 12.4 The Cross Product

**Definition 12.4.1** if  $\vec{d} = \langle a_1, a_2, a_3 \rangle$ ,  $\vec{b} = \langle b_1, b_2, b_3 \rangle$  then the cross product of  $\vec{d}$  &  $\vec{b}$  is:

$$\begin{aligned}\vec{d} \times \vec{b} &= \begin{vmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \\ &= \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} i - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} j + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} k \\ &= \langle a_2 b_3 - a_3 b_2, -(a_1 b_3 - a_3 b_1), a_1 b_2 - a_2 b_1 \rangle\end{aligned}$$

**Example 12.17** If  $\vec{d} = \langle 1, 2, -1 \rangle$ ,  $\vec{b} = \langle 2, 2, -3 \rangle$  Find  $\vec{d} \times \vec{b}$   
Solution:

$$\vec{d} \times \vec{b} = \begin{vmatrix} i & j & k \\ 1 & 2 & -1 \\ 2 & 2 & -3 \end{vmatrix} = (-4)i + 1(j) + (-2)k = \langle -4, 1, -2 \rangle.$$


**Theorem 12.4.1**  $\vec{a} \times \vec{b} \perp \vec{a}$  &  $\vec{a} \times \vec{b} \perp \vec{b}$

**proof:**  $(\vec{a} \times \vec{b}) \cdot \vec{a} = \langle a_2b_3 - a_3b_2, -a_1b_3 + a_3b_1, a_1b_2 - a_2b_1 \rangle \cdot \langle a_1, a_2, a_3 \rangle$   
 $= a_1a_2b_3 - a_1a_3b_2 - a_2a_3b_1 + a_1a_3b_2 - a_2a_3b_2 = \text{Zero}$   
 $\Rightarrow (\vec{a} \times \vec{b}) \perp \vec{a}$ .

**Theorem 12.4.2**  $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$ .  $0 \leq \theta \leq 180^\circ$

**proof:**  $|\vec{a} \times \vec{b}|^2 = (a_2b_3 - a_3b_2)^2 + (a_1b_3 - a_3b_1)^2 + (a_1b_2 - a_2b_1)^2$   
 $= a_2^2b_3^2 + a_3^2b_2^2 - 2a_2a_3b_2b_3 + a_1^2b_3^2 + a_3^2b_1^2 - 2a_1a_3b_1b_3 + a_1^2b_2^2 + a_2^2b_1^2 - 2a_1a_2b_1b_2$   
 $= (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2) - (a_1b_1 + a_2b_2 + a_3b_3)^2$   
 $= |\vec{a}|^2 |\vec{b}|^2 - (\vec{a} \cdot \vec{b})^2$   
 $= |\vec{a}|^2 |\vec{b}|^2 - |\vec{a}|^2 |\vec{b}|^2 \cos \theta^2$

**Corollary:** if  $\vec{a}, \vec{b}$  non-zero vectors, then  
 $\vec{a} \parallel \vec{b} \iff \vec{a} \times \vec{b} = \vec{0} \iff |\vec{a} \times \vec{b}| = 0 \iff \vec{a} = c \vec{b}$  for some  $c$



- $\vec{a} \times \vec{b} = \begin{vmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$
- $a \times b \perp a \& a \times b \perp b$
- $|a \times b| = |a| |b| \sin \theta$   $\theta \in [0, \pi]$
- $|a \times b| = \text{area of the parallelogram that determined by } a \times b$

■ **Example 12.18** find the area of the triangle with vertices

Solution :

- $P(2, 1, 3)$
  - $Q(1, -1, 1)$
  - $R(3, 2, -2)$
  - $\overrightarrow{PQ} = \langle -1, -2, -2 \rangle$
  - $\overrightarrow{PR} = \langle 1, 1, -5 \rangle$
  - $A = |\overrightarrow{PQ} \times \overrightarrow{PR}|$
- $$\overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} i & j & k \\ -1 & -2 & -2 \\ 1 & 1 & -5 \end{vmatrix} = \langle 12, -7, 1 \rangle$$
- $$\Rightarrow A = |\langle 12, -7, 1 \rangle|$$
- $$= \sqrt{144 + 49 + 1}$$
- $$= \sqrt{194}$$
- 
- 

■ **Example 12.19** Find

1.  $i \times i$
2.  $i \times j$
3.  $k \times j$
4.  $\langle -1, -2, -2 \rangle \times \langle 1, 1, -5 \rangle$

Solution :

1.  $\overrightarrow{0} = \langle 0, 0, 0 \rangle$
2.  $k$
3.  $-i$
4.  $= (-i - 2j - 2k) \times (i + j - 5k)$   
 $= -k - 5j + 2k + 10i - 2j + 2i$   
 $= \langle 12, -7, 1 \rangle$

■

---

■ **Example 12.20** True or False :

1.  $\overrightarrow{d} \times \overrightarrow{b} = \overrightarrow{b} \times \overrightarrow{d}$
2.  $(\overrightarrow{d} \times \overrightarrow{b}) \times \overrightarrow{c} = \overrightarrow{d} \times (\overrightarrow{b} \times \overrightarrow{c})$
3.  $i \times (i \times j) = i \times k = -j$   
 $(i \times i) \times j = \overrightarrow{0} \times i = \overrightarrow{0}$

Solution :

1. False :  $i \times j = k \neq j \times i = -k$
2. False

- $i \times (i \times j) = i \times k = -j$
- $(i \times i) \times j = \vec{0} \times i = \vec{0}$

■

Properties :

1.  $\vec{d} \times \vec{b} = -\vec{b} \times \vec{d}$
2.  $(c\vec{d}) \times \vec{b} = \vec{d} \times (c\vec{b}) = c(\vec{d} \times \vec{b})$
3.  $\vec{d} \times (\vec{b} + \vec{c}) = \vec{d} \times \vec{b} + \vec{d} \times \vec{c}$
4.  $(\vec{d} + \vec{b}) \times \vec{c} = \vec{d} \times \vec{c} + \vec{b} \times \vec{c}$
5.  $\vec{d} \cdot (\vec{b} \times \vec{c}) = (\vec{d} \times \vec{b}) \cdot \vec{c}$
6.  $\vec{d} \times (\vec{b} \times \vec{c}) = (\vec{d} \cdot \vec{c})\vec{b} - (\vec{d} \cdot \vec{b})\vec{c}$

Pf(5) if :

- $\vec{d} = \langle a_1, a_2, a_3 \rangle$
- $\vec{b} = \langle b_1, b_2, b_3 \rangle$
- $\vec{c} = \langle c_1, c_2, c_3 \rangle$

L.H.S

$$\begin{aligned}
 &= \vec{d} \cdot (\vec{b} \times \vec{c}) \\
 &= \langle a_1, a_2, a_3 \rangle \cdot \langle b_2c_3 - b_3c_2, b_3c_1 - b_1c_3, b_1c_2 - b_2c_1 \rangle \\
 &= a_1b_2c_3 - a_1b_3c_2 + a_2b_3c_1 - a_2b_1c_3 + a_3b_1c_2 - a_3b_2c_1
 \end{aligned}$$

R.H.S

$$\begin{aligned}
 &= (\vec{d} \times \vec{b}) \cdot \vec{c} \\
 &= \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1 \rangle \cdot \langle c_1, c_2, c_3 \rangle \\
 &= a_2b_3c_1 - a_3b_2c_1 + a_3b_1c_2 - a_1b_3c_2 + a_1b_2c_3 + a_2b_1c_3 \\
 &\Rightarrow \text{L.H.S} = \text{R.H.S}
 \end{aligned}$$

■ **Example 12.21** If  $\vec{d} \cdot (\vec{b} \times \vec{c}) = 2$ ,

Find  $2\vec{b} \cdot (\vec{d} \times 2\vec{c})$

Solution :

$$\begin{aligned}
 &= 4\vec{b} \cdot (\vec{d} \times \vec{c}) \\
 &= 4(\vec{d} \times \vec{c}) \cdot \vec{b} \\
 &= 4\vec{d} \cdot (\vec{c} \times \vec{b}) \\
 &= -4\vec{d} \cdot (\vec{b} \times \vec{c}) \\
 &= -8
 \end{aligned}$$

■

$$\vec{d} \cdot (\vec{b} \times \vec{c}) = (\vec{d} \times \vec{b}) \cdot \vec{c} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

triple product  $\vec{d} \cdot (\vec{b} \times \vec{c})$

volume of the parallelepiped :

$$\vec{v} = |\vec{d} \cdot (\vec{b} \times \vec{c})|$$

■ **Example 12.22** find the volume of the parallelepiped that determine by

$$\vec{d} = \langle 1, 2, -1 \rangle$$

$$\vec{b} = \langle 2, 1, 1 \rangle$$

$$\vec{c} = \langle 3, 2, -2 \rangle$$

Solution :

$$\begin{aligned} \vec{v} &= |\vec{d} \cdot (\vec{b} \times \vec{c})| = \left\| \begin{vmatrix} 1 & 2 & -1 \\ 2 & 1 & 1 \\ 3 & 2 & -2 \end{vmatrix} \right\| \\ &= |1(-4) - 2(-7) + -1(1)| \\ &= 9 \end{aligned}$$

if  $\vec{d} \cdot (\vec{b} \times \vec{c}) = 0$  then we say that  $\vec{d}$ ,  $\vec{b}$  and  $\vec{c}$  are called coplaner ■

■ **Example 12.23** show that the following vectors are coplaner

$$\vec{d} = \langle 2, 1, -1 \rangle$$

$$\vec{b} = \langle -1, 3, 2 \rangle$$

$$\vec{c} = \langle 0, 7, 3 \rangle$$

Solution :

$$\begin{aligned} \vec{d} \cdot \vec{b} \times \vec{c} &= \begin{vmatrix} 2 & 1 & -1 \\ -1 & 3 & 2 \\ 0 & 7 & 3 \end{vmatrix} \\ &= 2(-5) - 1(-3) + -1(-7) \\ &= -10 + 3 + 7 = 0 \end{aligned}$$

Thus **coplaner** ■

**Problem 12.4** 1,5,7,9,13,18,19,27,29,31,34,35,38,43

## 12.5 Equation of linear & planes

in 3D :

To determine a line , we need :

1. point  $(x_., y_., z_.)$
2. parallel vector  $\vec{v} = \langle a, b, c \rangle$

find it is equations !

note that  $\vec{v} \parallel \vec{r}$

$$\vec{v} = t \vec{d}, t \in R$$

**vector equation of the line .**

$$\langle x - x_0, y - y_0, z - z_0 \rangle = \langle ta, tb, tc \rangle$$

parametric equations of the line :

- $x = x_0 + at$
- $y = y_0 + bt$
- $z = z_0 + ct \quad -\infty < t < \infty$

- **Example 12.24** 1. find the parametric equations of the line that passes through the point  $(2, 1, 3)$  & parallel vector  $\vec{v} = \langle 2, 1, -1 \rangle$ .
2. Find the point on the line
3. does the point  $(0,0,5)$  lie on the line ?

Solution :

$$\begin{aligned} 1. \quad &x = 2 + 2t \\ &y = 1 + t \\ &z = 3 - 2t \quad -\infty < t < \infty \end{aligned}$$

$$\begin{aligned} 2. \quad &t = 2 \Rightarrow (6, 3, -1) \\ &t = \frac{7}{2} \Rightarrow \left(9, \frac{9}{2}, -4\right) \\ 3. \quad &0 = 2 + 2t \Rightarrow t = -1 \\ &0 = 1 + t \Rightarrow t = -1 \\ &5 = 3 - 2t \Rightarrow t = -1 \quad \text{yes.} \end{aligned}$$

■

- $x = x_0 + at \Rightarrow t = \frac{x - x_0}{a} \quad a \neq 0$
- $y = y_0 + bt \Rightarrow t = \frac{y - y_0}{b} \quad b \neq 0$
- $z = z_0 + ct \Rightarrow t = \frac{z - z_0}{c} \quad c \neq 0$

So, if

- $a \neq 0$
- $b \neq 0$
- $c \neq 0$

$$\Rightarrow \frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c} \quad a \neq 0, b \neq 0, c \neq 0$$

**symmetric equation**

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}$$

if  $a = 0$

$$x = x_0, \frac{y - y_0}{b} = \frac{z - z_0}{c}$$

- **Example 12.25** 1. find the equation of the line that passes thorough the point  $P(1, 2, -1)$  &  $Q(3, 1, 2)$   
 2. find where the line intersected the xy-plane !

Solution :

1. We need

- (a) point  $P(1, 2, -1)$
- (b)  $\vec{v} = \langle 2, -1, 3 \rangle$

**Parametric eq.s**

$$x = 1 + 2t$$

$$y = 2 - t$$

$$z = -1 + 3t$$

**Symmetric eq.**

$$\frac{x - 1}{2} = \frac{y - 2}{-1} = \frac{z + 1}{3}$$

2.  $z = 0$

$$\bullet \Rightarrow \frac{x - 1}{2} = \frac{1}{3}$$

$$\Rightarrow x - 1 = \frac{2}{3}$$

$$\Rightarrow x = \frac{5}{3}$$

$$\bullet \frac{y - 2}{-1} = \frac{1}{3} \rightarrow y - 2 = \frac{-1}{3} \rightarrow y = \frac{5}{3}$$

$$\bullet \left( \frac{5}{3}, \frac{5}{3}, 0 \right)$$

■ **Example 12.26** Find the parametric equations of the line that passes through the point  $(-2, 1, 1)$  & parallel to the line :

$$L_1 = \frac{x - 2}{1} = \frac{2 - y}{1} = \frac{2z + 1}{1}$$

Solution: We need

1. point  $(-2, 1, 1)$
2.  $\vec{v} = \langle 1, -1, \frac{1}{2} \rangle$

$$x = 2 + t$$

$$y = 1 - t$$

$$z = 1 + \frac{1}{2}t \quad t \in R$$

### line

1. point  $(x_0, y_0, z_0)$
2. parallel vector  $\vec{v} = \langle a, b, c \rangle$

### Parametric equation

- $x = x_0 + at$
- $y = y_0 + bt$
- $z = z_0 + ct \quad -\infty < t < \infty$

### Symmetric equations

$$\bullet \frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}$$



two lines are parallel iff their vector are parallel

■ **Definition 12.5.1** two lines are called skew if they are not parallel & they do not intersect .

■ **Example 12.27** show that the following lines are skew

$$L_1 : x = 1 + t, y = -2 + 3t, z = 4 - t / \vec{v}_1 = \langle 1, 3, -1 \rangle$$

$$L_2 : x = 2s, y = 3 + s, z = -3 + 4s / \vec{v}_2 = < 2, 1, 4 >$$

### Solution

$v_1 \neq v_2 \Rightarrow L_1 \neq L_2$  ( $L_1 \& L_2$  are not parallel )

$$1 + t = 2s \quad t - 2s = -1 \rightarrow (1)$$

$$-2 + 2t = 3 + s \quad 3t - s = 5 \rightarrow (2)$$

$$4 - t = -3 + 4s \quad -t - 4s = -7 \rightarrow (3)$$

Solve (1)&(3)

$$0 - 6s = -8 \Rightarrow s = \frac{8}{6} = \frac{4}{3}$$

$$t = -1 + 2s \Rightarrow t = -1 + \frac{8}{3} = \frac{5}{3}$$

$$s = \frac{4}{3}, t = \frac{5}{3}$$

in Equation 2

$$3\frac{5}{3} - \frac{4}{3} \neq 5 \Rightarrow 5 - \frac{4}{3} \neq 5$$

$L_1 \& L_2$  do not intersect  $\Rightarrow L_1 \& L_2$  are skew

■

Planes : to determine a plane we need

1. point  $(x_0, y_0, z_0)$
2. normal vector  $\vec{n} = < a, b, c >$  Note that  $\vec{v} \perp \vec{n}$   
 $\Rightarrow \vec{v} \cdot \vec{n} = 0$

$$\Rightarrow < x - x_0, y - y_0, z - z_0 > \cdot < a, b, c > = 0$$

$$\Rightarrow a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$$

$$\Rightarrow ax + by + cz + -(ax_0 + by_0 + cz_0) = 0$$

$$\Rightarrow ax + by + cz + d = 0$$

$$\Rightarrow d = -(ax_0 + by_0 + cz_0)$$

- **Example 12.28** 1. find the equation of the plane that passes through the point  $(1, -1, 3)$  & normal vector  $\vec{v} = < 2, 1, -1 >$ .

2. find two points on the plane .

### Solution

1. point p(1,-1,3)

2. normal vector  $\vec{v} = < 2, 1, -1 >$ .

$$2(x - 1) + 1(y + 1) + -1(z - 3) = 0$$

$$2x + y - z + 6 = 0$$

$$2) (0,0,6) (-3,0,0) (0,-6,0)$$

- **Example 12.29** find the equation of the plane that passes through the points  
 $P(2, 1, -2)$     $Q(1, 1, -1)$     $R(3, -2, 1)$

Solution

1. Point  $(2, 1, -2)$
2.  $\vec{n} = \overrightarrow{PQ} \times \overrightarrow{PR}$   
 $= <3, 4, 3>$   
 $= 3x + 4y + 3z + -4$   
 $= 0$   
 $\overrightarrow{PQ} = <-1, 0, 1>$   
 $\overrightarrow{PR} = <1, -3, 3>$

$$\vec{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} i & j & k \\ -1 & 0 & 1 \\ 1 & -3 & 3 \end{vmatrix}$$

$$= <3, 4, 3>.$$

---

Plane

- $p(x_0, y_0, z_0)$
- $\vec{n} = <a, b, c>$   
 $ax + by + cz + d = 0$   
 $d = -(ax_0 + by_0 + cz_0)$

- **Example 12.30** Find the equation of the plane that passes through the point  
 $p(1, 2, 1)$ ,  $Q(2, 3, 2)$ ,  $R(-1, -1, 3)$

Solution

$$ax + by + cz + d = 0$$

- if  $d \neq 0$   
 $Ax + By + Cz + 1 = 0$   
 $A + 2B + C + 1 = 0$   
 $2A + 3B + 2C + 1 = 0$   
 $-A - B - 3C + 1 = 0 \quad (\text{Rejected})$
- $d = 0$   
 $ax + by + cz + 1 = 0$

■ **Example 12.31** Find the equation of the plane that passes through the point  $(1, -1, 2)$  & contains the line

$$L_1 : \frac{x-1}{2} = \frac{y+1}{2} = \frac{z-1}{3}, \vec{v} = \langle 2, 3, 1 \rangle$$

Solution

1. Point  $(1, -1, 2)$
2. normal vector

$$\begin{aligned}\vec{n} &= \vec{r} \times \vec{v} \\ &= \vec{RQ} \times \vec{v} \\ &= \langle 0, 0, 1 \rangle \times \langle 2, 3, 1 \rangle \\ &= -2x + 2y + 0z + 4 = 0 \\ &\Rightarrow x - y - 2 = 0\end{aligned}$$

■ **Example 12.32** find the equations of the line of intersection of the following planes .

$$P_1 : 2x - y + z = 0, \vec{n}_1 = \langle 2, -1, 1 \rangle$$

$$P_2 : x - 3y - z - 1 = 0, \vec{n}_2 = \langle 1, -3, -1 \rangle$$

Solution

- Point :  $P(0, -1, 2)$
- Parallel vector

$$\begin{aligned}\vec{v} &= \vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} i & j & k \\ 2 & -1 & 1 \\ 1 & -3 & -1 \end{vmatrix} \\ &= \langle 4, 3, -5 \rangle \text{ let } x=0\end{aligned}$$

$$-y + z = 3$$

+

$$-3y - z = 1$$

$$\Rightarrow -4y = 4 \Rightarrow y = -1$$

$$\Rightarrow z = 2$$

$$x = 0 + 4t$$

$$y = -1 + 3t$$

$$z = 2 - 5t$$

Solution 2 : Pick two points on the line :

- $P(0, -1, 2)$

- $Q\left(\frac{4}{3}, 0, \frac{1}{3}\right)$

Let  $y = 0$

$$2x + z = 3$$

+

$$x - z = 1$$

$$\Rightarrow 3x = 4$$

$$\Rightarrow x = \frac{4}{3}$$

$$\Rightarrow z = \frac{4}{3} - 1 = \frac{1}{3}$$

$$\vec{v} = \overrightarrow{PQ} = \left\langle \frac{4}{3}, 1, \frac{-5}{3} \right\rangle$$

- $x = 0 + \frac{4}{3}t$

- $y = -1 + t$

- $z = 2 - \frac{5}{3}t$

Solution 3 :

$$2x - y + z - 3 = 0$$

$$x - 3y - z - 1 = 0$$

let  $x = t$

$$\Rightarrow 2t - y + z - 3 = 0$$

$$\Rightarrow t - 3y - z - 1 = 0$$

↓

$$-y + z = 3 - 2t$$

+

$$-3y - z = 1 - t$$

$$\Rightarrow -4y = 4 - 3t$$

$$\Rightarrow y = -1 + \frac{3}{4}t$$

$$-3\left(-1 + \frac{3}{4}t\right) - z = 1 - t$$

$$z = 3 - \frac{9}{4}t - 1 + t$$

$$z = 2 - \frac{5}{4}t$$

R

- two plane are parallel if their normal vectors are parallel .
- the angle btw two plane is defined to be the acute angle btw  $\vec{n}_1$  &  $\vec{n}_2$ .

■ **Example 12.33** find the angle btw the following plane :  $2x - 2y = z - 1 = 0$

$x = 3y - z = 7 = 0$ , . [Solution](#):

- $\vec{n}_1 = \langle 2, -2, 1 \rangle$
- $\vec{n}_2 = \langle 1, 3, -1 \rangle$

$$\theta : \cos\theta = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|} = \frac{2 - 6 - 1}{3\sqrt{11}} = -\frac{5}{3\sqrt{11}} \approx 120.1$$

$$\Rightarrow \alpha = 59.9$$

■ **Example 12.34** find the intersection btw the following line :

$$P : 2x - 2y + z_1 = 0$$

$$L : x = 1 + t, y = 1 - t, z = t.$$

[Solution](#):

$$2(1+t) - 2(1-t) + t - 1 = 0$$

$$2 + 2t - 2 + 2t + t = 1$$

$$5t = 1 \Rightarrow t = \frac{1}{5}$$

$$x = 1 + \frac{1}{5} \Rightarrow \frac{6}{5}$$

$$y = 1 - \frac{1}{5} \Rightarrow \frac{4}{5}$$

$$z = \frac{1}{5}$$

[Distances](#) :

$$\frac{1}{2} * |\overrightarrow{QR}| * D = \frac{1}{2} * |\overrightarrow{QR} * \overrightarrow{PQ}|$$

$$D = \frac{|\overrightarrow{QR}| * |\overrightarrow{PQ}|}{|QR|}$$

OR

$$D = \frac{|\overrightarrow{v} * \overrightarrow{PQ}|}{|\overrightarrow{v}|}$$

■ **Example 12.35** Find the distance below  $(1, 2 - 1)$  & the line

$$x = 1 + t$$

$$y = 1 - t$$

$$z = t$$

Solution : Pick two points on the line

$$t = 0 \Rightarrow Q(1, 1, 0)$$

$$t = 1 \Rightarrow R(2, 0, 1)$$

$$D = \frac{|\overrightarrow{QR}| * |\overrightarrow{PQ}|}{|QR|}$$

$$\overrightarrow{PQ} = \langle 0, -1, 1 \rangle$$

$$\overrightarrow{QR} = \langle 1, -1, 1 \rangle$$

$$\overrightarrow{PQ} * \overrightarrow{QR} = \begin{vmatrix} i & j & k \\ 0 & -1 & 1 \\ 1 & -1 & 1 \end{vmatrix}$$

$$= \frac{\sqrt{0+1+1}}{\sqrt{1+1+1}} = \sqrt{\frac{2}{3}}$$

■ **Example 12.36** Find the distance below the point  $(x, y, z)$  & the plane

$$ax + by + cz = 0$$

Solution :

$$D = \left| \text{Comp}_{\vec{n}} \overrightarrow{r} \right|$$

$$= \frac{|\overrightarrow{r} \cdot \vec{n}|}{|\vec{n}|} \quad \overrightarrow{r} = \langle x_0 - x_1, y_0 - y_1, z_0 - z_1 \rangle$$

$$= \frac{|a(x_1 - x_0) + b(y_1 - y_0) + c(z_1 - z_0)|}{\sqrt{a^2 + b^2 + c^2}} \quad \vec{n} = \langle a, b, c \rangle$$

$$= \frac{|ax_1 + by_1 + cz_1 - (ax_0 + by_0 + cz_0)|}{\sqrt{a^2 + b^2 + c^2}}$$

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

- **Example 12.37** Find the distance between the point  $(1, 2, -1)$  and the plane  $2x + 2y - z - 3 = 0$

$$\begin{aligned} D &= \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}} \\ \text{Solution : } &= \frac{|2(1) + 2(2) - 1(-1) - 3|}{\sqrt{9}} \\ &= \frac{4}{3} \end{aligned}$$

- **Example 12.38** Find the distance between the following planes

$$P_1 : x + y - z + 1 = 0$$

$$P_2 : 2x - 2y + 3z + 7 = 0$$

$$\text{Solution : } \vec{n}_1 = \langle 1, 1, -1 \rangle$$

$$\vec{n}_2 = \langle 2, -2, 3 \rangle$$

$$\Rightarrow \vec{n}_1 \nparallel \vec{n}_2$$

$$\Rightarrow P_1 \nparallel P_2$$

$\Rightarrow P_1$  and  $P_2$  are intersected

$$\Rightarrow D = 0$$

■

- **Example 12.39** Find the distance between the following plane

$$P_1 : x - 2y + 2z + 3 = 0$$

$$P_2 : -2x + 4y - 4z - 5 = 0$$

Solution :

$$\vec{n}_1 = \langle 1, -2, 2 \rangle$$

$$\vec{n}_2 = \langle -2, 4, -4 \rangle$$

Thus  $\vec{n}_1 \parallel \vec{n}_2$

Pick any point on  $P_1 \Rightarrow p(-3, 0, 0)$

$$D = \frac{|-2(-3) + 4(0) - 4(0) - 5|}{\sqrt{4 + 16 + 16}} = \frac{1}{6}$$

■

- **Example 12.40** Find the distance between the line and plane

$$L : x = 1 + t, \quad y = 1 - t, \quad z = t$$

$$P : 2x - y + z + 3 = 0$$

Solution (1):

$$\vec{v} = \langle 1, -1, 1 \rangle$$

$$\vec{n} = \langle 2, -1, 1 \rangle$$

$$P \parallel L \iff \vec{v} \perp \vec{n} \iff \vec{n} \cdot \vec{v} = 0$$

Note that

$$\begin{aligned}\vec{n} \cdot \vec{v} &= 2 + 1 + 1 \\ &= 4 \\ &\neq 0 \\ \Rightarrow \vec{n} &\not\perp \vec{v} \\ \Rightarrow P &\nparallel L \\ \Rightarrow D &= 0\end{aligned}$$

### Solution (2):

Try to find an intersection point

$$2(1+t) - (1-t) + t + 3 = 0$$

$$2 + 2t - 1 + t + t + 3 = 0$$

$$4t = -4 \Rightarrow t = -1$$

P & L are intersecting

$$\Rightarrow D = 0$$

### ■ Example 12.41 Find the distance between the following plane & line

$$L : x = 1 - t, \quad y = t, \quad z = 2 - t$$

$$P : 2x + y - z + 3 = 0$$

### Solution :

$$\vec{v} = \langle -1, 1, -1 \rangle$$

$$\vec{n} = \langle 2, 1, -1 \rangle$$

$$\vec{n} \cdot \vec{v} = 0 \Rightarrow P \parallel L$$

pick a point on the line  $t = 0 \Rightarrow (1, 0, 2)$

$$\begin{aligned}D &= \frac{|2+0-2+3|}{\sqrt{4+1+4}} \\ &= \frac{3}{\sqrt{6}} \\ &= \sqrt{\frac{3}{2}}\end{aligned}$$

### ■ Example 12.42 Find the distance between the following lines

$$L_1 : x = 1 - t, \quad y = 1 + t, \quad z = t$$

$$L_2 : x = -1 + 2t, \quad y = -2t, \quad z = 1 - 2t$$

### Solution :

$$\vec{v_1} = \langle -1, 1, 1 \rangle$$

$$\vec{v_2} = \langle 2, 2, -2 \rangle$$

$$\vec{v_1} \parallel \vec{v_2} \Rightarrow L_1 \parallel L_2$$

**Exercise 12.3** Pick a point on  $L_1$   $t = 0 \Rightarrow p(1, 1, 0)$

■ **Example 12.43** If

$$L_1 : x = 1 + t, \quad y = 2 + 3t, \quad z = 4 - t$$

$$L_2 : x = 2s, \quad y = 3 + s, \quad z = -3 + 4s$$

are skew , Find the distance between them .

Solution : Construct two parallel planes that contain  $L_1 \& L_2$  respectively

$P_1$

$$\text{point } t = 0 \Rightarrow (1, -2, 4)$$

$$\vec{n} = \vec{v_1} * \vec{v_2}$$

$$\vec{n} = 13i - 6j - 5k$$

$$13x - 6y - 5z + 0 = 0$$

$$d = -(13 + 12 - 20)$$

$$D = \frac{|13(0) - 6(3) - 5(-3) + 0|}{\sqrt{13^2 + 6^2 + 25}} = \frac{|-8|}{\sqrt{230}} = \frac{8}{\sqrt{230}}$$

$P_2$

$$\text{point } s = 0 \Rightarrow (0, 3, -3)$$

$$\vec{n} = \langle 13, -6, -5 \rangle$$

$$13x - 6y - 5z + 3 = 0$$

■ **Example 12.44** Determine whether each sentence is true or false .

1. Two lines parallel to a third line are parallel .
2. Two lines perpendicular to a third line are parallel .
3. Two planes parallel to a third plane are parallel .
4. Two planes perpendicular to a third plane are parallel .
5. Two lines parallel to a plane are parallel.
6. Two lines perpendicular to a plane are parallel .
7. Two planes parallel to a line are parallel .
8. Two planes perpendicular to a line are parallel .
9. Two planes either intersect or are parallel .
10. Two lines either intersect or are parallel .
11. A plane and a line either intersect or are parallel .

Solution

1. T

5. F

9. T

2. F

6. T

10. F

3. T

7. F

11. T

4. F              8. T

---

■ **Example 12.45** Show that the distance between the following planes

$$P_1 : ax + by + cz + d_1 = 0$$

$$P_2 : ax + by + cz + d_2 = 0, \text{ is}$$

$$D = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$

Solution :

Pick a point on  $P_1(x_1, y_1, z_1)$

$$\text{So } D = \frac{|ax_1 + by_1 + cz_1 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$


---

■ **Example 12.46** Find equations of the parallel planes to

$$P_1 : x - 2y - 2z + 1 = 0 \text{ and units away from it.}$$

Solution :

$$P_2 : x - 2y - 2z + d = 0$$

$$D = 2$$

$$\frac{|d - 1|}{\sqrt{1+4+4}} = 2$$

$$\Rightarrow |d - 1| = 6$$

$$\Rightarrow d - 1 = -6 \text{ OR } d - 1 = 6$$

$$\Rightarrow d = -5 \text{ OR } d = 7$$


---

■ **Example 12.47** Find the projection of the point  $(1, 2, -1)$  on the plane

$$2x - 2y + z - 1 = 0$$

Solution:

Let's construct a line that passes through  $(1, 2, -1)$  & perpendicular to the plane .

point  $(1, 2, -1)$

parallel vector  $\vec{n} = \langle 2, -2, 1 \rangle$

$$x = 1 + 2t$$

$$y = 2 - 2t$$

$$z = -1 + t$$

we will find the intersection between the line and the plane

$$2(1 + 2t) - 2(2 - 2t) + (-1 + t) - 1 = 0$$

$$9t = 4 \Rightarrow t = \frac{4}{9}$$

$$\text{point } \left(1 + \frac{8}{9}, 2 - \frac{8}{9}, -1 + \frac{4}{9}\right) = \left(-\frac{17}{9}, \frac{10}{9}, -\frac{5}{9}\right)$$

■

■ **Example 12.48** 1.  $x + y + z = c$

2.  $x + y + z = 1$

3.  $(\cos c)y + (\sin c)z = 1$

■

---

**Problem 12.5** 1,3,5,9,11,12,13,14,17,19,21,25,26,29,30,31,35,37,38,39,45,46,48,51-57(odd),61-71(odd),74,76

# 13. Vector Function

$f : \mathbb{R} \mapsto \text{vector}$  (real valued function)

$$f(x) = \sin x$$

$$f(\pi/2) = 1$$

## 13.1 Vector Functions and space curves

$$\begin{aligned}\vec{r}(t) &:= \langle f(t), g(t), h(t) \rangle \} \text{ vector function} \\ &= f(t)i + g(t)j + h(t)k\end{aligned}$$

---

■ **Example 13.1**  $\vec{r}(t) = \langle t^2, 1-t, t^2+1 \rangle$   
 $\vec{r}(1) = \langle 1, 0, 2 \rangle$  ■

---

■ **Example 13.2** Find the domain of  $\vec{r}(t) = \langle \sqrt{t}, \ln(1-t), t^2 \rangle$   
 $D_{\vec{r}(t)} = D_f \cap D_g \cap D_h = [0, 1)$  ■

---

Limit and continuity :

if  $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$   
then  $\lim_{t \rightarrow t_0} \vec{r}(t) = \langle \lim_{t \rightarrow t_0} f(t), \lim_{t \rightarrow t_0} g(t), \lim_{t \rightarrow t_0} h(t) \rangle$

---

■ **Example 13.3** Find  $\lim_{t \rightarrow t_0} \langle 2t, \frac{\sin t}{1 - e^t}, \frac{t}{t} \rangle = \langle 0, 1, -1 \rangle$

$\vec{r}(t)$  is cont at  $t_0 \Leftrightarrow \lim_{t \rightarrow t_0} \vec{r}(t) = \vec{r}(t_0)$

■ **Definition 13.1.1** Space Curve: suppose that  $f, g, h$  are cont real-value function on I(interval) then

$$x = f(t), y = g(t), z = h(t)$$

is called space curve it can be represented using  $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$

■ **Example 13.4** Describe the curve defined by

$$1. \vec{r}(t) = \langle t, 1+t, 2-t \rangle$$

$$x = t$$

$$y = 1+t, \quad -\infty < t < \infty$$

$$z = 2-t$$

$$2. \vec{r}(t) = \langle \cos t, \sin t, t \rangle$$

$$x = \cos t, y = \sin t, z = t$$

■ **Definition 13.1.2** Line Segment :the line segment from  $\vec{r}_0$  to  $\vec{r}_1$ ,  $0 \leq t \leq 1$

■ **Example 13.5** Find the vector function that represented the line segment  $P(1, 2, -1)$  &  $Q(2, 3, 2)$

$$\vec{r}(t) = (1-t) \langle 1, 2, -1 \rangle + t \langle 2, 3, 2 \rangle, \quad 0 \leq t \leq 1$$

■

■ **Example 13.6** Find a vector function that represented the intersection of  $x^2 + y^2 = 1$  and  $y + z = 2$  [Solution](#)

$$\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$$

$$\vec{r}(t) = \langle \cos t, \sin t, 2 - \sin t \rangle, \quad 0 \leq t \leq 1$$

■

**Problem 13.1** 1,3,5,15,17,25,27,35,37,42

**13.2 13.2**

$$\begin{aligned}\vec{r}'(t) &= \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h} \\ &= \langle f'(t), g'(t), h'(t) \rangle \\ &= f'(t)\mathbf{i} + g'(t)\mathbf{j} + h'(t)\mathbf{k}\end{aligned}$$


---

■ **Example 13.7** if  $\vec{r}(t) = (1+t)\mathbf{i} + te^{-t}\mathbf{j} + \sin 2t\mathbf{k}$

a Find  $\vec{r}'$

b unit tangent vector at  $t = 0$

Solution

$$\vec{r}'(t) = 2t\mathbf{i} + (e^{-t} - te^{-t})\mathbf{j} + 2\cos 2t\mathbf{k}$$

$\vec{r}'(0) = \langle 0, 1, 2 \rangle$  tangent vector

$$\vec{T}(t) = \frac{\vec{r}'(0)}{|\vec{r}'(0)|} = \frac{1}{\sqrt{0+1+4}} \langle 0, 1, 2 \rangle = \langle 0, \frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}} \rangle$$


---

■ **Example 13.8** Find parametric equation for tangent line to the helix:

$$\vec{r}(t) = \langle \cos t, \sin t, t \rangle \text{ at } t = \pi$$

Solution

point  $(-\mathbf{1}, 0, \pi)$

parallel vector  $\vec{r}'(t) = \langle -\sin t, \cos t, 1 \rangle$

$$\vec{r}'(\pi) = \langle 0, -1, 1 \rangle$$

$$x = -1 + 0t$$

$$y = 0 - t$$

$$z = \pi + t$$


---

Differential Rule:

1.  $\frac{d}{dt}(\vec{u}(t) + \vec{v}(t)) = \frac{d}{dt}\vec{u}(t) + \frac{d}{dt}\vec{v}(t)$
  2.  $(c\vec{u}(t))' = c\vec{u}'(t)$
  3.  $\frac{d}{dt}(f(t)\vec{u}(t)) = f'(t)\vec{u}(t) + f(t)\vec{u}'(t)$
  4.  $\frac{d}{dt}(\vec{u}(t) \cdot \vec{v}(t)) = \vec{u}'(t) \cdot \vec{v}(t) + \vec{u}(t) \cdot \vec{v}'(t)$
  5.  $\frac{d}{dt}(\vec{u}(t) \times \vec{v}(t)) = \vec{u}'(t) \times \vec{v}(t) + \vec{u}(t) \times \vec{v}'(t)$
  6.  $\frac{d}{dt}\vec{u}(f(t)) = f'(t)\vec{u}'(f(t))$
-

■ **Example 13.9** if  $\vec{u}'(1) \times \vec{v}(1) = \langle 2, -1, 3 \rangle$

$$\vec{v}'(1) \times \vec{u}'(1) = \langle 2, -1, 3 \rangle$$

$$\text{Find } (\vec{u} \times \vec{v})'(1) = \vec{u}' \times \vec{v} + \vec{u} \times \vec{v}'$$

$$= \langle 2, -1, 3 \rangle + \langle -2, 1, -3 \rangle = \langle 0, 0, 0 \rangle$$

■

■ **Example 13.10** if  $|\vec{r}(t)| = c$  (*constant*)

show that  $\vec{r}'(t) \perp \vec{r}(t)$

Proof

$$|\vec{r}(t)| = \text{const}$$

$$|\vec{r}(t)|^2 = \text{const}$$

$$\vec{r}(t) \cdot \vec{r}(t) = \text{const}$$

■

Differentiate:

$$\vec{r}'(t) \cdot \vec{r}'(t) + \vec{r}(t) \cdot \vec{r}'(t) = 0$$

$$2\vec{r}'(t) \cdot \vec{r}'(t) = 0$$

$$\vec{r}'(t) \cdot \vec{r}'(t) = 0$$

$$\vec{r}'(t) \perp \vec{r}(t)$$

■ **Definition 13.2.1** Integrals:  $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$

$$\text{then } \int \vec{r}(t) dt = \langle \int f(t) dt, \int g(t) dt, \int h(t) dt \rangle$$

■ **Example 13.11** if  $\vec{r}(t) = 2ti - e^t j + lntk$

$$\text{Find } \vec{r}(t) \text{ where } \vec{r}(1) = \langle 0, 0, 1 \rangle \quad \vec{r}(t) = \langle t^2 + c_1, -e^t + c_2, tlnt - t + c_3 \rangle$$

$$1 + c_1 = 0 \Rightarrow c_1 = -1$$

$$c_2 - e = 0 \Rightarrow c_2 = e$$

$$-1 + c_3 = 1 \Rightarrow c_3 = 2$$

■

**Problem 13.2** 3,4,5,6,9,11,12,17,19,21,23,25,32,34,37,39,49

### 13.3 Arc Length

$$L = \int_a^b \sqrt{f'^2(t) + g'^2(t) + h'^2(t)} dt$$

$$L = \int_a^b |\vec{r}'(t)|$$

■ **Example 13.12** Find the length of the helix  $\vec{r}(t) = \langle \cos t, \sin t, t \rangle \quad 0 \leq t \leq 2\pi$

$$t \leq \pi$$

Solution

$$L = \int_0^\pi \sqrt{\sin^2 t + \cos^2 t + 1} dt = \int_0^\pi \sqrt{2} dt = \pi\sqrt{2}$$

■ **Example 13.13** Two particles travel along the curves

$$\vec{r}_1(t) = < t, t^2, t^3 >$$

$$\vec{r}_2(t) = < 1+2s, 1+6s, 1+14s >$$

1. Do the particles collide?

2. Do their paths intersect?

$$t = 1+2s, \quad t^2 = 1+6s, \quad t^3 = 1+14s \Rightarrow$$

$$(1+2s)^2 = 1+6s \Rightarrow 1+4s+4s^2 = 1+6s \Rightarrow$$

$$s(4s-2) = 0 \Rightarrow s=0, s=1/2 \Rightarrow$$

$$s=0, t=1 \Rightarrow 1 \stackrel{?}{=} 1$$

$$s=1/2, t=2 \Rightarrow 8 \stackrel{?}{=} 8$$

the paths intersect two lines, at the point  $(1, 1, 1)$  &  $(2, 4, 8)$

But they do not collide, the paths intersect at different  $t, s$

■

---

**Problem 13.3** 1,3,5



# 14. Partial Derivatives

## 14.1 Function of several variables

**Definition 14.1.1** A function of two variables is a rule that assigns for each  $(x, y)$  in the domain one value  $z = f(x, y)$  in the range  
 $x, y$  are called independent variables  
 $z$  is called an independent variable

■ **Example 14.1** Find the domain of the following function

1.  $f(x, y) = \sqrt{y - x + 1}$

$D = \{(x, y) : y - x + 1 \geq 0\}$

Range:  $R = [0, \infty)$

2.  $f(x, y) = \sqrt{9 - x^2 - y^2} + \sqrt{x}$

$D = \{(x, y) : 9 - x^2 - y^2 \geq 0 \text{ & } x \geq 0\}$

3.  $f(x, y) = \ln(x^2 + y^2 - 9)$

$D = \{(x, y) : x^2 + y^2 - 9 > 0\}$

$R = (-\infty, \infty)$

4.  $f(x, y) = \frac{\sin^{-1}(x - y)}{\sqrt{x - y^2}} \mid \sin[-\pi/2, \pi/2] \rightarrow [-1, 1]$

$D = \{(x, y) : -1 \leq x - y \leq 1 \text{ & } x - y^2 > 0\}$

(a)  $-1 \leq x - y$

(b)  $x - y \leq 1$

(c)  $x - y^2 > 0$

■ **Example 14.2** Find the range of  $f(x,y) = \sqrt{9 - x^2 - y^2}$

$$D = \{(x,y) : x^2 + y^2 \leq 9\}$$

$$R = [0, 3]$$

$$0 \leq x^2 + y^2 \leq 9$$

$$0 \geq -x^2 - y^2 \geq -9$$

$$9 \geq 9 - x^2 - y^2 \geq 0$$

$$3 \geq \sqrt{9 - x^2 - y^2} \geq 0$$

Domain 2D.

Range 1D.

■ **Definition 14.1.2** let  $z = f(x,y)$  then the graph of the function is the set:

$$G = \{(x,y,z) : (x,y) \in D, z = f(x,y)\}$$

■ **Example 14.3** Sketch the following

$$1. f(x,y) = 6 - 2x - 3y$$

$$z = 6 - 2x - 3y$$

$$2x + 3y + z - 6 = 0$$

*x-intercept*  $y = 0, z = 0 \Rightarrow x = 3$  *y-intercept*  $x = 0, z = 0 \Rightarrow y = 2$

*z-intercept*  $x = 0, y = 0 \Rightarrow z = 0$

$$2. g(x,y) = \sqrt{9 - x^2 - y^2}$$

$$z = \sqrt{9 - x^2 - y^2}$$

$$z^2 = 9 - x^2 - y^2$$

$$x^2 + y^2 + z^2 = 9, z \geq 0$$

■ **Definition 14.1.3** let  $z = f(x,y)$  then the level curve of  $f$  at  $k \in \text{Range}$  is the set

$$L = \{(x,y) : k = f(x,y)\} \supseteq \mathbb{R}^2$$

■ **Example 14.4** Find level curve to  $f(x,y) = 6 - 2x - 3y$  at  $k = 0, 6, -6\dots$

Solution

$$k = 0 \Rightarrow 0 = 6 - 2x - 3y \Rightarrow y = 2 - \frac{2}{3}x$$

$$k = 6 \Rightarrow 6 = 6 - 2x - 3y \Rightarrow y = -\frac{2}{3}x$$

■ **Example 14.5** Find level curve for  $f(x, y) = \sqrt{9 - x^2 - y^2}$  at:

$$k = 1 \Rightarrow 1 = \sqrt{9 - x^2 - y^2} \Rightarrow x^2 + y^2 = 8$$

$$k = 2 \Rightarrow 2 = \sqrt{9 - x^2 - y^2} \Rightarrow x^2 + y^2 = 5$$

$$k = 0 \Rightarrow 0 = \sqrt{9 - x^2 - y^2} \Rightarrow x^2 + y^2 = 9$$

■

■ **Example 14.6** Sketch some level curve of the function  $f(x, y) = 4x^2 + y^2$

$$k = 1 \Rightarrow 4x^2 + y^2 = 1$$

$$\frac{x^2}{1} + \frac{y^2}{\frac{1}{4}} = 1$$

$$k = 4 \Rightarrow 4x^2 + y^2 = 4$$

$$\frac{x^2}{4} + \frac{y^2}{1} = 1$$

■

**Definition 14.1.4** Function of three variable

$$\begin{array}{ccc} w & = & f(x, y, z) \\ \text{dependent variable} & & \text{independent variable} \end{array}$$

Domain: 3D

Range: 1D

Graph: 4D

Level surface: 3D

■ **Example 14.7** Find the domain of  $f(x, y, z) = \sqrt{9 - x^2 - y^2 - z^2}$

$$D = \{(x, y, z) : x^2 + y^2 + z^2 \leq 9\}$$

■

■ **Example 14.8** Find the following limits if exists:

$$1. \lim_{(x,y) \rightarrow (1,2)} (2x + y) = 2(1) + 2 = 4$$

$$2. \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x - y} = \lim_{(x,y) \rightarrow (0,0)} \frac{(x - y)(x + y)}{x - y} = 0$$

$$3. \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2} \text{ we will take different paths.}$$

path 1: along  $y = 0$

$$\lim_{x \rightarrow 0} \frac{x^2}{x^2} = 1$$

path 2: along  $x = 0$

$$\lim_{y \rightarrow 0} \frac{0 - y^2}{0 + y^2} = -1$$

$$-1 \neq 1 \Rightarrow D.N.E$$

4.  $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$

path  $y = 0 \Rightarrow \lim_{x \rightarrow 0} \frac{0}{x^2} = 0$

path  $x = 0 \Rightarrow \lim_{y \rightarrow 0} \frac{0}{y^2} = 0$

path  $y = x \Rightarrow \lim_{y \rightarrow 0} \frac{y^2}{y^2 + y^2} = 1/2$

$0 \neq 1/2 \Rightarrow D.N.E$

path  $y = mx \Rightarrow \lim_{x \rightarrow 0} \frac{mx^2}{x^2 + m^2 x^2} = \frac{m}{1 + m^2}$   
depends on  $m$  so D.N.E

5.  $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$

path  $y = mx \rightarrow \lim_{x \rightarrow 0} \frac{xm^2 x^2}{x^2 + m^4 x^4}$

$= \lim_{x \rightarrow 0} \frac{x^3}{x^2} \frac{m^2}{1 + m^4 x^2}$

$= \lim_{x \rightarrow 0} \frac{x^3}{x^2} \frac{m^2}{1 + m^4 x^2}$

$\rightarrow 0 \rightarrow m$

$= 0$

path  $x = my^2$

$\lim \frac{xy^2}{x^2 + y^2} = \lim_{y \rightarrow 0} \frac{my^2 y^2}{m^2 y^4 + y^4} = \frac{m}{m^2 + 1}$  depends on  $m$  D.N.E

6.  $\lim_{(x,y) \rightarrow (0,0)} \frac{\sin(x^2 + y^2)}{x^2 + y^2}$

$r = \sqrt{x^2 + y^2}$

$g = \tan^{-1} y(x, y) \rightarrow 0$

$r \rightarrow 0^+ g \rightarrow ??$

$= \lim_{r \rightarrow 0^+} \frac{\sin r^2}{2} = 1$

7.  $\lim_{(x,y) \rightarrow (0,0)} \frac{1 - e^{\sqrt{x^2 + y^2}}}{\sqrt{x^2 + y^2}} = \lim_{r \rightarrow 0^+} \frac{1 - e^r}{r} = \lim_{r \rightarrow 0^+} \frac{-e^r}{1} = -1$

$$8. \lim_{(x,y) \rightarrow (0,0)} \frac{x^3 - y^3}{x^2 + y^2} = \lim_{r \rightarrow 0^+} \frac{r^3 \cos^3 \Theta - r^3 \sin^3 \Theta}{r^2} \\ = \lim_{r \rightarrow 0^+} r(\cos^3 \Theta - \sin^3 \Theta) = 0$$

$$9. \lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2} = \lim_{r \rightarrow 0^+} \frac{r \cos \theta r \sin \theta}{r^2} = \cos \theta \sin \theta \text{ depends on } \theta \text{ dose not exist}$$

$$10. \lim_{(x,y) \rightarrow (0,0)} \frac{xy}{\sqrt{x^2 + y^2}} = \lim_{r \rightarrow 0^+} \frac{r \cos \theta r \sin \theta}{r} = 0$$

$$11. \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{x^2 + y^2} = \lim \frac{r^3 \cos^3 \theta + r^3 \sin^3 \theta}{r^2 (\cos^2 \theta - \sin^2 \theta)}$$

$$= \lim_{r \rightarrow 0^+} r \left( \frac{\cos^3 \theta + \sin^3 \theta}{\cos^2 \theta - \sin^2 \theta} \right) \text{ D.N.E / if } \theta = \frac{\pi}{y}$$

continuity  $z = f(x, y)$  is cont at  $(x, y)$

if  $\lim_{(x,y) \rightarrow (x,y)} f(x, y) = f(x, y)$

■ **Example 14.9** find where the function  $f(x, y) : \begin{cases} \frac{x^2 - y^2}{x^2 + y^2} & : (x, y) \neq (0, 0), \\ 0 & : (x, y) = (0, 0) \end{cases}$ ,

is cont

### Solution

$f$  is cont for  $(x, y) \neq (0, 0)$  at  $(0, 0)$ ?

- $f(0, 0) = 0$

- $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$  D.N.E  $\rightarrow f$  is not cont at  $(0, 0)$

$f$  is cont  $R^2 (0, 0)$

■

**Problem 14.1** 6, 7, 9, 11, 13, 15, 19, 20, 21, 23, 25, 29, 41, 61, 63

## 14.2 Partial Derivatives

■ **Definition 14.2.1** if  $z = f(x, y)$

$$\frac{\partial z}{\partial x} = \frac{\partial f}{\partial x} = f_x = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

$$\frac{\partial z}{\partial y} = \frac{\partial f}{\partial y} = f_y = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}$$

Rules:

1. to find  $f_x$  treat y as a constant.
  2. to find  $f_y$  treat x as a constant.
- 

■ **Example 14.10** find  $f_x, f_y$ :

$$1. \quad f(x, y) = x + y + xy$$

$$f_x = 1 + 0 + y = 1 + y$$

$$f_y = 1 + x$$

$$2. \quad f(x, y) = \frac{x}{y}$$

$$f_x = \frac{1}{y}$$

$$f_y = x \left( -\frac{1}{y^2} \right)$$

$$3) \quad f(x, y) = xe^{xy}$$

$$f_x = e^{xy} + xye^{xy}$$

$$f_y = x^2 e^{xy}$$

$$f(x, y) = \frac{2}{x} - x \ln y$$

$$f_x = \frac{-2}{x^2} - \ln y$$

$$f_y = 0 - \frac{x}{y}$$

■

---

Function of three variables:

$w = f(x, y, z)$  to find

$f_x$  treat y,z as constant .

$f_y$  treat x,z as constant .

$f_z$  treat x,y as constant .

Ex : if  $f(x, y, z) = xy + \frac{1}{z} - e^{xz}$

find  $f_x = y + 0 + -ze^{xz}$

$f_y = x$

$f_z = 0 + \frac{-1}{z^2} - xe^{xz}$

Higher Derivatives:

---

■ **Example 14.11** if  $f(x,y) = x^2y + xy - 2ye^x$

find  $f_{xx}, f_{xy}, f_{yy}$

$$f_x = 2xy + y - 2ye^x$$

$$f_y = x^2 + x - 2e^x$$

$$f_{xx} = 2y - 2ye^x$$

$$f_{xy} = 2x + 1 - 2e^x$$

$$f_{yx} = 2x + 1 - 2e^x$$

$$f_{yy} = 0$$

clairaut's theorem : if  $f$  defined on a disc  $D$  that contains  $(a,b)$  if  $f_{xy}, f_{yx}$  cont on  $D$  then  $f_{xy}(a,b) = f_{yx}(a,b)$  ■

■ **Example 14.12** T/F : there exists a function s.t  $f_x = 21x + 3y$

$$f_y = x^2 - 2y$$

Solution: note that  $f_{xy} = 3f_{yx} = 2x$

$f_{xy} \neq f_{yx}$  false . ■

■ **Example 14.13** if  $f(x,y,z) = z^2 \cos(x+2y)$  find  $f_{zxyz}$

Solution  $f_z = 2z \cos(x+2y)$

$$f_{zx} = -2z \sin(x+2y)$$

$$f_{zxy} = -4z \cos(x+2y)$$

$$f_{zxyz} = -y \cos(x+2y)$$
 ■

### Partial differential Equation:

1. Laplace's Equation:  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$

Solution of this equation are called :harmonic equation

■ **Example 14.14** show that  $u(x,y) = x^2 - y^2$  is harmonic

$$u_x = 2x, u_y = -2y$$

$$u_{xx} = 2, u_{yy} = -2$$

$$\text{so } u_{xx} + u_{yy} = 0$$
 ■

■ **Example 14.15** show that  $u(x,y) = e^x \cos y$  satisfies Laplace's equation

$$u_x = e^x \sin y, u_y = e^x \cos y$$


$$u_{xx} = e^x \sin y, u_{yy} = -e^x \sin y$$

$$\Rightarrow u_{xx} + u_{yy} = 0$$

---

## 2. Wave equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial x^2}$$



■ **Example 14.16** show that  $u(x, t) = \sin x - at$  satisfies the wave equation

$$u_t = -a \cos(x - at), u_x = \cos(x - at)$$

$$\text{L.H.S} = u_{tt} = -a^2 \sin(x - at), \text{R.H.S} = u_{xx} = -\sin(x - at)$$

$$\text{L.H.S} = \text{R.H.S}$$

---

**Problem 14.2** 5,7,8,9,12,13,14,18,21,25,29,31,33,37,39,40

### 14.3 Tangent Plane and Linear Approximation

Let  $z = f(x, y)$  then the tangent plane at  $(x_0, y_0)$

$$z = z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

---

■ **Example 14.17** Find the tangent plane of  $f(x, y) = 2x^2 + y^2$

at  $(1, 1, 3)$

$$f_x = 4x \Rightarrow f_x(1, 1) = 4$$

$$f_y = 2y \Rightarrow f_y(1, 1) = 2$$

$$z - 3 = 4(x - 1) + 2(y - 1)$$

$$4x + 2y - z - 3 = 0$$

---

Linear Approximation: let  $z = f(x, y)$ ,  $f_x, f_y$  cont

The linear approximation of  $f$  at  $(a, b)$

$$\text{is } L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

---

■ **Example 14.18** let  $f(x, y) = xe^{xy}$

1. Find linear approximation at  $(1, 0)$

2. Approximation  $f(1.1, -0.1)$

#### Solution

$$f(1, 0) = 1$$

$$f_x = xy e^{xy}$$

$$f_x(1,0) = 1$$

$$f_y = x^2 e^{xy}$$

$$f_y(1,0) = 1$$

$$\Rightarrow L(x,y) = 1 + 1(x-1) + 1(y-0)$$

$$L(x,y) = x+y \approx xe^{xy} \text{ around } (1,0)$$

$$f(1.1, -0.1) \approx L(1.1, -0.1) = 1.1 - 0.1 = 1$$

$$f(1.1, -0.1) = 0.9854$$

■

■ **Example 14.19** if the tangent plane to  $z = f(x,y)$  at  $(2,3)$

$$\text{is } 2x - 3y + z = 1 \Rightarrow L(x,y) = 1 - 2x + 3y$$

$$\text{Approximation } f(2.1, 2.9)$$

$$\text{sol } f(2.1, 2.9) \approx 1 - 2(2.1) + 3(2.9) = 1 - 4.2 + 8.7 = 5.5$$

■

■ **Example 14.20** Approximation  $12\sqrt{8.9} - 12\sqrt[3]{8.1}$

### Solution

$$f(x,y) = 12\sqrt{x} - 12\sqrt[3]{y} \text{ at } (9,8)$$

$$L(x,y) = 12 + 2(x-9) - 1(y-8)$$

$$f(8.9, 8.1) \approx L(8.9, 8.1) = 12 + 2\left(\frac{-1}{10} + \frac{-1}{10}\right)$$

$$12 - \frac{3}{10} = 11.7$$

■

**Definition 14.3.1** Differentials: if  $z = f(x,y)$ , then we define the differential

$$dz = f_x \partial x + f_y \partial y$$

$$dz = \frac{\partial f}{\partial x} \partial x + \frac{\partial f}{\partial y} \partial y$$

let:

$$\partial x = \delta x = x - a$$

$$\partial y = \Delta y = y - a$$

$$\partial z = \delta z = z - z_0 = f(x,y) - f(a,b)$$

$$dz = \frac{\partial f}{\partial x} |_{(a,b)} (x-a) + \frac{\partial f}{\partial y} |_{(a,b)} (y-b)$$

■ **Example 14.21** let  $t = f(x,y) = x^2 + 3xy - y^2$

1. find the differential

$$\partial t = (2x + 3y)\partial x + (3x - 2y)\partial y$$

2. if  $x$  change from 2 to 2.05

$y$  change from 3 to 2.96

com pane  $\partial z, \delta z$

$$\partial z = (2(2) + 3(3))\frac{5}{100} + (3.2.2.3)\frac{-4}{100}$$

$$= \frac{65}{100} = 0.65$$

$$(2, 3) \rightarrow (2.05, 2.96)$$

$$\Delta z = f((2.05, 2.96)) - f(2, 3) = 0.6449$$

$$dz = \delta z = z - z.$$

■

functions of three variables

$$w = f(x, y, z)$$

$$\partial w = f_x \partial x + f_y \partial y + f_z \partial z$$

**Problem 14.3** 1-43(odd), 44, 48, 49, 50, 51, 53, 59, 61, 65, 71, 72(a,d), 73, 75, 77, 87\*, 89\*, 93, 94

## 14.4 Directional derivatives and Gradient vector

$$f_x = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

$$f_y = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}$$

**Definition 14.4.1** the direction derivative of  $f$  at  $(x_0, y_0)$  in the direction of the limit vector  $\vec{u} = \langle a, b \rangle$  is  $D_{\vec{u}} f(x_0, y_0) = \lim \frac{f(x_0 + ah, y_0 + bh) - f(x_0, y_0)}{h}$

Gradient vector:

**Definition 14.4.2** if  $z = f(x, y)$ , then the gradient of  $f$  at  $(x_0, y_0)$  is  $\nabla f = \langle f_x(x_0, y_0), f_y(x_0, y_0) \rangle$

■ **Example 14.22** if  $f(x, y) = x^2 - y^2$  find  $\nabla f |_{(1, 2)}$

$$f_x = 2x \rightarrow f_x(1, 2) = 2$$

$$f_y = 2y \rightarrow f_y(1, 2) = -4$$

Solution

$$\nabla f|_{(1,2)} = \langle 2, -4 \rangle$$

**Theorem 14.4.1**  $D_{\vec{u}} f(x, y) = \nabla f \cdot \vec{u}$

- **Example 14.23** if  $f(x, y) = \sin x + e^{xy}$  find the directional derivative of  $f$  at  $(0, 1)$  in the direction of  $\vec{u} = \langle 3, -4 \rangle$

so,  $D_{\vec{u}} f(0, 1) = \nabla f \cdot \vec{u}$

$$= \langle 2, 0 \rangle \cdot \left\langle \frac{3}{5}, \frac{4}{5} \right\rangle$$

$$= \frac{6}{10} + 0 \cdot \frac{4}{5} = (0, 6)$$

$$\nabla f = \langle f_x, f_y \rangle = \langle 2, 0 \rangle$$

$$f_x = \cos x + ye^{xy}$$

$$f_y = xe^{xy}$$

$$f_x(0, 1) = 2$$

$$f_y(0, 1) = 0$$

function of three variables  $w = f(x, y, z)$

$$\nabla f = \langle f_x, f_y, f_z \rangle$$

$$D_{\vec{u}} f(a, b, c) = \nabla f \cdot \vec{u}$$

■

- **Example 14.24** find the direction derivative of  $f(x, y, z) = \frac{x-y}{z} + x^2 + e^y$  at  $p(1, 0, 1)$  in the direction to the point  $Q(-1, 2, 0)$

Solution

$$: D_{\vec{u}} f(-1, 2, 0) = \nabla f \cdot \vec{u} = \langle 3, 0, -1 \rangle \cdot \left\langle \frac{2}{3}, \frac{2}{3}, \frac{-1}{3} \right\rangle = -2 + 0 + \frac{1}{3} =$$

$$-1 \frac{2}{3}$$

$$\vec{u} = \frac{\vec{PQ}}{|\vec{PQ}|} = \left\langle \frac{-2}{3}, \frac{2}{3}, \frac{-1}{0} \right\rangle$$

$$f_x = \frac{1}{z} + 2x, f_x(1, 0, 1) = 1 + 2 = 3$$

$$f_y = \frac{-1}{z} + e^y \rightarrow f_y(1, 0, 1) = -1 + 1 = 0$$

$$f_z = \frac{y-x}{z^2}, f_z(1, 0, 1) = \frac{0-1}{1^2} = -1$$

$$\nabla f = \langle 3, 0, -1 \rangle$$

■

■ **Example 14.25** if  $D_{\vec{u}} f = 3\nabla f \cdot \vec{u}$

$$\text{if } D_{\vec{u}} \nabla f = 6$$

$$D_{\vec{u}} \nabla f = -6$$

$$D_{\vec{u}}^2 f = 3$$

$$D_{-\vec{u}}^2 f = -3$$

■

Question  $z = f(x, y)$  ( $x, y$ )

$$\vec{u} = ??$$

find  $\vec{u}$  that maximize  $D_{\vec{u}} f(x, y) = \nabla f \cdot \vec{u} = |\nabla f| |\vec{u}| \cos$

**Theorem 14.4.2**

- the max directional derivative of  $f$  at  $(x, y)$  is  $|\nabla f|$  and it accrues if  $\vec{u}$  has the same direction of  $\nabla f$ .
- the min directional derivative of  $f$  is  $-|\nabla f|$  and it accrues if  $\vec{u}$  has the opposite direction of  $\nabla f$ .

■ **Example 14.26** Let  $z = f(x, y) = xe^y$  find the max directional derivative at  $(2, 0)$ .

### Solution

$$\nabla f = \langle f_x, f_y \rangle, f_x = e^y, f_y = xe^y$$

$$\nabla f = \langle 1, 2 \rangle$$

$$\max D_{\vec{u}} f(2, 0) = |\nabla f| = \sqrt{5}$$

it occurs if  $\vec{u}$  has the same direction of  $\langle 1, 2 \rangle$

max directional derivative  $\leftrightarrow$  max rate of change

$\leftrightarrow$  increasing most rapidly


min directional derivative  $\leftrightarrow$  min rate of change

$\leftrightarrow$  decreasing most rapidly

■

### Tangent plane for level surfaces

$k = f(x, y, z)$  level surfaces  
 to find the plane we need 1. point  
 $(x_0, y_0, z_0)$  2.  $\vec{n} = \nabla f$



- **Example 14.27** 1. find the equation of the tangent plane to  $\frac{x^2}{4} + y^2 + \frac{z^2}{9} = 3$  at  $(2, 1, 3)$   
 2. find the equation of the normal line

### Solution

1. plane ! point  $(2, 1, 3)$

$$\vec{n} = \nabla f = \left\langle \frac{2x}{4}, 2y, \frac{2}{9} \right\rangle = \left\langle 1, 2, \frac{2}{3} \right\rangle$$

$$|(x-2) + 2(y-1) + \frac{2}{3}(z-3)|$$

2. point  $(2, 1, 3)$

$$\vec{n} = \left\langle 1, 2, \frac{2}{3} \right\rangle, x = 2 + t$$

$$, y = 1 + 2t$$

$$, z = 3 + \frac{2}{3}t$$


$$\nabla f = \langle f_x, f_y, f_z \rangle$$

$$D_{\vec{u}} f(x_0, y_0, z_0) = \nabla f \cdot \vec{u}$$

max  $D_{\vec{u}} f = |\nabla f|$  it accrues if  $\nabla f, \vec{u}$  have the same direction


$F(x, y, z) = K$  the  $n$  the normal to the tangent  $D_{\vec{n}} f = |\nabla f|$

## 14.5 The Chain Rule



Case I:


$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$$



Case II:

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s}$$



Case III:

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial s}$$


■ **Example 14.28** If  $u = x^4y + y^2z^3$

$$x = rse^t$$

$$y = rs^2e^{-t}$$

$$z = r^2s \sin t$$

$$\text{Find } \frac{\partial u}{\partial s} \text{ when } r = 2, s = 1, t = 0$$



Solution


$$\begin{aligned} \frac{\partial u}{\partial s} &= u_x \frac{\partial x}{\partial s} + u_y \frac{\partial y}{\partial s} + u_z \frac{\partial z}{\partial s} \\ &= (4x^3y)re^t + (x^4 + 2yz^3)(2rse^{-t}) + (3y^2t^2)(r^2 \sin t) \\ \frac{\partial u}{\partial s} \Big|_{(r,s,t)=(2,1,0)} &= (64)2 + (16+0)4 + 0 = 128 + 64 = 192 \end{aligned}$$

■

■ **Example 14.29** .

$$\text{if } g(s, t) = f(s^2 - t^2, t^2 - s^2)$$

$$\text{show that } g \text{ satisfies } t \frac{\partial g}{\partial s} + s \frac{\partial g}{\partial t} = 0$$



Solution

$$g(s, t) = f(x, y)$$

$$\frac{\partial g}{\partial s} = \frac{\partial g}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial g}{\partial y} \frac{\partial y}{\partial s}$$

$$\frac{\partial g}{\partial s} = f_x(2s) + f_y(-2s) \cdots 1$$

$$\frac{\partial g}{\partial t} = \frac{\partial g}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial g}{\partial y} \frac{\partial y}{\partial t}$$

$$\frac{\partial g}{\partial t} = f_x(-2t) + f_y(2t) \cdots 2$$

$$\text{so, } t \frac{\partial g}{\partial s} + s \frac{\partial g}{\partial t} = f_x(2st) + f_y(-2st) + f_x(-2st) + f_y(2st) = 0$$

■ **Example 14.30 .**

let  $z = f(x, y)$  , ( $f$  has cont second partial derivative)

if  $x = r^2 + s^2$  ,  $y = 2rs$

find:




$$1. \frac{\partial z}{\partial r}$$

$$2. \frac{\partial^2 z}{\partial r^2}$$

Solution

$$\begin{aligned} \frac{\partial z}{\partial r} &= f_x \frac{\partial x}{\partial r} + f_y \frac{\partial y}{\partial r} = f_x(2r) + f_y(2s) \\ &= 2rf_x + 2sf_y \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 z}{\partial r^2} &= 2[r(f_{xx} \frac{\partial x}{\partial r} + f_{xy} \frac{\partial y}{\partial r}) + f_x] + \\ &2s[f_{yx} \frac{\partial x}{\partial r} + f_{yy} \frac{\partial y}{\partial r}] \\ &= (4r^2)f_{xx} + 8rsf_{xy} + 4s^2f_{yy} \end{aligned}$$



■ **Definition 14.5.1** Implicit differentiation

if  $F(x, y, z) = 0$

$$\begin{aligned} \text{then } \frac{\partial z}{\partial x} &= -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}} \\ \frac{\partial z}{\partial y} &= -\frac{\frac{\partial F}{\partial y}}{\frac{\partial F}{\partial z}} \end{aligned}$$

■ **Example 14.31** if  $x^3 + y^3 + z^3 = 1 - 6xyz$

Find  $\frac{\partial z}{\partial x}, \frac{\partial z}{\partial y}$

Solution

$$\begin{aligned} x^3 + y^3 + z^3 + 6xyz - 1 &= 0 \\ \frac{\partial z}{\partial x} &= -\frac{F_x}{F_z} = -\frac{3x^2 + 6yz}{3z^2 + 6xy} \\ \frac{\partial z}{\partial y} &= -\frac{F_y}{F_z} = -\frac{3y^2 + 6xz}{3z^2 + 6xy} \end{aligned}$$

■

**Problem 14.5** 3,4,5,7,9,11,12,13,14,15,16,17,19,21,23,26,27,28,31,33,35,39,43,46,48,50,51