
Index

- A_n -singularity, 411
 - and double points, 411
 - blow-up of a curve at a singularity, 423–424
 - Abel’s theorem, 572–573
 - Abel–Jacobi maps, 572
 - and projective bundles, 573–574, 587–589
 - birationality onto image, 573
 - class of image, 585
 - fibers, 587
 - used to describe cohomology of the Jacobian, 583
 - abstract secant variety, 371
 - of a rational normal curve, 374
 - adjunction formula, 40–41
 - gives genus formula, 69
 - statement, 41
 - used to calculate canonical classes of hypersurfaces, 41
 - adjunction map, 526–527
 - affine space, 24
 - affine stratification, 26–28
 - definition, 27
 - gives basis for Chow group, 28
 - of $\mathbb{G}(1, 3)$ via Schubert cells, 102, 126
 - of $G(k, n)$, 135–137, 161
 - of a blow-up of \mathbb{P}^n , 58, 80
 - of flag manifolds, 160
 - of projective space, 44
 - of the product of projective spaces, 51
 - affine tangent space, 10
 - algebraic cycles modulo algebraic equivalence, 551
 - algebraic cycles modulo homological equivalence, 380, 552
 - algebraic stacks, 502
 - alternative cycle theories, 550–552
 - a comparison, 553
 - advantages over Chow ring, 564–565
 - analytic topology, 344
 - apparent nodes, 114
 - asterisk, 65
 - Atiyah class, 251
 - Azumaya algebras, 346
 - Bézout’s theorem, 14–15, 17, 46–48
 - for dimensionally transverse intersections, 14
 - general statement, 46
 - base change, 524
 - associated natural map, 524–525
 - commuting with direct image, 525–526
 - for finite morphisms, 525–526
 - for flat base change, 525
 - of an affine morphism, 525
 - commuting with higher direct image, 531–533
 - Bertini’s theorem, 11, 230
 - application to lines on a cubic, 221
 - extension, 237–238
 - strong form, 169–170
 - Betti numbers
 - odd Betti numbers of a smooth projective variety are even, 549
 - of a hypersurface, 182
 - of a K3 surface, 192
 - of the quadric line complex, 192
 - binodal curves
 - in a net, 425
 - birational equivalence
 - between the Hilbert scheme and Kontsevich space, 315
 - blow-up, 36
 - applied to the five conic problem, 291–292
 - as projective bundle, 337–338
 - Chow ring, 471–473
 - generators, 473
 - relations, 476–477
 - of \mathbb{P}^2 at a point, 339
 - of \mathbb{P}^3 along a line, 81
 - class of proper transform of a smooth surface, 357
 - of \mathbb{P}^3 along a smooth curve, 473–475, 478–479
 - of \mathbb{P}^4 along a line, 357
 - of \mathbb{P}^5 , 301–302
 - of \mathbb{P}^n along a linear space, 337–339, 357, 358
 - Chow ring, 479
 - various classes, 357
 - of \mathbb{P}^n at a point, 56–60, 72, 80
 - of a curve at an A_n -singularity, 423–424
 - of a singular curve, 74
 - of a surface at a point, 72–74
 - of the Veronese surface in \mathbb{P}^5 , 480
 - applied to the five conic problem, 480
 - resolving indeterminacy of a rational map, 282
 - utility in calculating intersection multiplicities, 61–62
- boundary, 289
 - branch divisor, 278
 - Brauer group, 346
 - Brauer–Severi variety, 344–346
 - Brill–Noether locus, 573

- as degeneracy locus, 580–583, 591–592
- class, 577, 590–592
- description, 578–579
- dimension, 576, 579, 590
 - bound for a general curve, 579–580
- nonemptiness, 583, 590
- scheme structure, 583
- Brill–Noether theorem, 564
 - existence part, 578–579
 - used to count linear series, 577
 - version 1, 566
 - version 2, 576
 - comparison with Castelnuovo’s bound, 576
 - version 3, 590–592
 - via curves on a K3 surface, 580
 - via degeneration to singular curves, 579
- bundle of k -frames, 561
- bundle of principal parts, 244, 247–248
 - advantage, 254
 - Chern class, 254
 - via Grothendieck Riemann–Roch, 508
 - definition and properties, 248–250
 - does not behave well in families, 272
 - of a line bundle on \mathbb{P}^n , 251
- bundle of relative principal parts, 391–392
 - Chern class, 508
 - properties, 392
- canonical bundle, 39
- canonical class, 14, 39–40
 - of a blow-up, 73–74
 - of a subvariety, 40
 - of hypersurfaces and complete intersections, 41–42
 - of projective space, 39–40
- Cartesian flex, 271
- Cartier divisor, *see* divisor, Cartier
- Castelnuovo bound, 386
- Castelnuovo’s theorem, 570–571
 - open problem, 570
- Castelnuovo–Mumford regularity, 261
- Catalan number, 149, 577
- Čech complex, 528–529
- characteristic classes, 559
- Chern character, 177, 484–486
 - computes the Chern class of a tensor product, 486
 - definition, 485–486
 - is a ring homomorphism, 485–486
 - is an isomorphism up to torsion, 486–487
 - of the tangent bundle of $\mathbb{G}(1, 3)$, 507
- Chern class, 14, 134, *see also* topological Chern class
 - alternative definition via Grassmannian, 170
 - and the Chow ring of a projective bundle, 331
 - as degeneracy locus, 167–168, 177, 426
 - characterization, 167–169
 - coincides with topological Chern class for an algebraic vector bundle, 561
 - construction, 170–172
 - definition for smooth quasi-projective varieties, 169
 - first Chern class, 37, 167
 - is a homomorphism, 37
 - of a line bundle, 37–42
 - on a singular variety, 38
 - general definition, 171
 - Grothendieck’s definition via projective bundles, 332
 - in connection with lines on a cubic, 199–200
 - in connection with the degree of a discriminant hypersurface, 253
 - information about the Grothendieck ring, 486–487
 - introductory example, 166
 - is the reciprocal of the Segre class, 364
 - of $\mathcal{E} = \text{Sym}^3 \mathcal{S}^*$, 200
 - via Grothendieck Riemann–Roch, 490–493
 - of $\mathcal{O}_{\mathbb{P}^r}(1)$, 177
 - of $\Phi(3, 3, 1)$, 232
 - of $\text{Sym}^d \mathcal{S}^*$, 227
 - of a bundle of principal parts, 252–255
 - on \mathbb{P}^n , 254
 - of a coherent sheaf is well-defined, 507
 - of a determinant, 173–174
 - of a direct sum, 169, 174
 - of a Fano scheme, 198
 - of a smooth curve, 179
 - of a symmetric power tensored with a line bundle, 255
 - of a symmetric square, 174
 - of a tensor product with a line bundle, 174–176
 - of tautological quotient bundle, 347
 - of the dual bundle, 173
 - of the relative tangent bundle of a projective bundle, 394
 - of the structure sheaf of a point in \mathbb{P}^n , 507
 - of the structure sheaf of a smooth curve in \mathbb{P}^3 , 507
 - of the tangent bundle, 179–183
 - of $\mathbb{G}(1, 3)$, 507
 - of a Grassmannian, 183
 - of a hypersurface, 179–180
 - of a product of projective spaces, 192
 - of a quadric in \mathbb{P}^5 , 192
 - of products of projective spaces, 508
 - of projective space, 179
 - of the Grassmannian of $G(2, 4)$, 191
 - of the tensor product of a vector bundle and a dual bundle, 428
 - of the tensor product of bundles, 176, 191
 - of the universal bundles on the Grassmannian, 178–179
 - of the universal quotient bundle on projective space, 177–178
 - parallel with Segre class, 364
 - top Chern class of a tensor product via the resultant, 428
 - topological, *see* topological Chern class
 - vanishes above the rank, 173
 - via Grothendieck Riemann–Roch, 490
- Chern polynomial, 427
- Chinese remainder theorem, 11
- chords, 113
 - of two twisted cubics, 85, 115, 129
 - via specialization, 122, 130
- to a curve, 113–115
 - class, 163
 - computing the class via specialization, 121–122, 130

- Chow cohomology, 77
 - Chow group, 16–17
 - analogy with (co)homology, 15, 28
 - definition*, 16
 - grading, 17
 - of a Grassmannian via Young diagrams, 153
 - of a projective bundle, 331
 - of a subset of affine space, 26
 - of an affine space, 24
 - theory carries over to cycles modulo algebraic equivalence, 551
 - via affine stratification, 28
 - Chow ring, 14
 - computation, 22–36
 - definition*, 19
 - examples, 44–62
 - existence, 19
 - introduction, 15–19
 - of a 0-dimensional scheme, 22
 - of a blow-up, 471–473
 - generators, 473
 - of \mathbb{P}^3 along a smooth curve, 474–475
 - of \mathbb{P}^n along a linear space, 338–339
 - of \mathbb{P}^n at a point, 56–61
 - of a surface, 72
 - of the Veronese surface in \mathbb{P}^5 , 480
 - relations, 476–477
 - of a flag variety, 356, 359
 - of a Grassmannian bundle, 346–347
 - of a parameter space, 62, 86
 - of conics in \mathbb{P}^3 , 349
 - of a product of projective spaces, 51–52, 79
 - of a projective bundle, 331–335
 - of projective space, 44
 - of the Grassmannian, 137, 183–187
 - of lines in \mathbb{P}^3 , 105–109
 - of the space of complete conics, 306–309
 - of the universal hyperplane, 336
 - of the universal line, 337, 394
 - of the universal plane, 335–336
 - relation to transversality, 17–19
 - circle
 - circular points, 66
 - in complex projective space, 66–67
 - tangent to a given circle, 67–68
 - tangent to three general circles, *see* circles of Apollonius
 - circles of Apollonius, 66–68
 - classical topology
 - and algebraic geometry, 544
 - is finer than the Zariski topology, 543
 - Clemens conjecture, 239
 - Clifford's theorem, 569–570
 - codimension
 - definition*, 10
 - expected, *see* expected codimension
 - of a Schubert cycle, 133
 - of a subvariety, 17
 - cofactor map, 297
 - Cohen–Macaulay variety
 - and Bézout's theorem, 46–47
 - and intersection multiplicity, 72
 - and the theory of liaison, 71
 - pullback, 30
 - coherent sheaf
 - criterion to be a vector bundle, 536
 - has a locally free resolution, 486
 - cohomology
 - group vs. ring, 192
 - of \mathbb{P}^3 , 192
 - of a smooth quadric threefold, 192
 - collinearity, 80
 - compactification, 290, 292
 - of the total space of a vector bundle, 343
 - complete conic, *see* space of complete conics
 - complete flag, 102, 132
 - complete intersection, 109
 - and the excess intersection formula, 455
 - Chern class, 180
 - counting lines, 240
 - homology and cohomology, 555
 - hyperplane section with triple points, 286, 421
 - normal bundle, 212
 - singular curves on a complete intersection, 286
 - subvariety contained in no smooth hypersurface, 555
 - subvariety of codimension-1 is intersection with hypersurface, 556–557
- complete linear series
 - allowed by Castelnuovo, 576
 - corresponding to a line bundle on a curve, 567
 - completeness of the adjoint series, 386
 - complex of flat modules
 - approximation by finitely generated free modules, 537
 - for quasi-projective schemes, 541
 - complex projective variety
 - as holomorphic subvariety/submanifold, 543
 - complexification
 - and Hodge theory, 546
 - of the cotangent bundle of a complex manifold, 546
 - composition series, 11
 - cone, 83, 257, 453
 - cone construction, 512
 - conic curve
 - in \mathbb{P}^4 , 361
 - conormal variety, 380
 - contact problem, 389
 - cotangent bundle, 179
 - cubic surface
 - can have at most four isolated singular points, 236, 241
 - cannot have three collinear isolated singular points, 241
 - must contain lines, 201
 - curvature form, 562
 - curvature matrix, 562
 - curve of genus 4
 - expressed as a 3-sheeted cover of \mathbb{P}^1 , 564, 577–578
 - curve of genus 6
 - expressed as a degree-6 curve in \mathbb{P}^2 , 564
 - curve of genus 8

- embedded as a degree-9 curve in \mathbb{P}^3 , 564
 - image curve of embedding does not lie on a cubic, 593
- curves expressed as covers of \mathbb{P}^1 , 566
- cuspidal, 34, 410
 - in a net of plane curves, 413–419
 - ordinary, *see* ordinary cuspidal
- cycle, 15
 - definition*, 15
 - of plane conics in \mathbb{P}^3 meeting a given line, 348
 - class, 349, 359
 - tangent spaces, 352–353
- Debarre–de Jong conjecture, 227, 236–239, 243
- defective variety, 371
 - curves are not defective, 372
 - defective Veronese varieties, 373, 385–386
 - equivalent characterization via tangent spaces, 372
- deformations, *see* first-order deformations
- degeneracy locus, 167, 168, 426–427
 - class, 429, *see also* Porteous’ formula
 - is independent of sections chosen, 187
 - expected codimension, 427
 - from pencils, 251–252
 - geometry, 168
 - reducedness, 252
- degeneration to singular curves
 - in connection with Brill–Noether, 579
- degree map, 29
- degree of a covering, 28
- del Pezzo surface, 288
- derivations
 - identification with Zariski tangent space, 100
- derived category, 534
 - is formal, 535, 540–541
- determinant of a bundle, 173
- determinantal variety, 430
 - degree, 433, 442
 - scheme structure on the Brill–Noether locus, 583
- diagonal of $\mathbb{P}^r \times \mathbb{P}^r$
 - class, 53–54
 - generalization, 189
 - via specialization, 79
- dimensional transversality, 31–33, 46
 - definition*
 - for cycles, 32
 - for subschemes, 32
 - fails for cycles in a proper subvariety, 465
 - to a morphism, 518
 - sufficient condition, 518
 - weaker than generic transversality, 33
- dimensionally proper intersection, 14
- direct image, 520–521, 523–528
 - conditions to be a vector bundle, 525–526
 - definition*, 523
 - for a projective morphism, 523
 - for an affine morphism, 523
 - for finite morphisms, 525
 - higher, *see* higher direct image
 - of a line bundle, 541
- direct image complex, 533–534
 - explicit computation, 534, 541–542
 - with terms given by sums of line bundles, 541
- directrix, 357
- discriminant, 244, 245, 418
 - definition*, 258
 - degree in the space of forms of degree d , 253
 - is an irreducible hypersurface, 245–246
 - linearizing the description, 247
 - multiplicity, 280–281
 - at a double conic, 288
 - at a double line, 288
 - of a net
 - of plane curves, 274–276
 - of a quartic polynomial, 246, 247
 - of a very ample linear series, 258
 - smooth locus, 284
 - tangent cone, 284
 - tangent space, 282–284
 - of discriminant of degree- d polynomials, 285, 304
- divisor, 15
 - associated to a rational function, 23
 - Cartier, 37, 41, 165
 - cohomology of an ample divisor, 553
 - of a function, 23
 - of a nonaffine variety, 23
- divisor class group, 22, 306
- divisor classes in the space of complete conics, 307–309
- Dolbeault complex, 548
- double conic, 478
- double point, 34, 411
 - is an A_n -singularity, 411
 - of a curve, 411–412
- dual conic
 - degeneration, 293–295
- dual variety, 259, 297, 380
 - of a hypersurface, 49–50, 78
 - degree, 50, 382
 - of a quadric, 296–299
 - of a smooth complete intersection, 388
 - of a smooth conic, 293
 - divisor, 298
 - of a smooth hypersurface, 381
 - in char p , 387
 - of a smooth variety failing to be a hypersurface, 382, 388
 - of a smooth variety tends to be singular, 382, 388
 - of the Veronese embedding, 247
 - reflexivity of projective varieties, 294, 381
- dualizing sheaf, 503
- dynamic projection
 - in a family of projective spaces, 333–334
 - introduction, 117–119
- dynamic specialization, 150–152, 162–163
- Eckhart point, 420
- elementary symmetric functions, 169, 175, 485
- elliptic curve
 - elliptic quintic lies on no planes or quadrics, 452

- secant varieties, 388
- symmetric power, 388
- elliptic normal curve, 287
 - characterized by independence of divisors, 386
- embedded component, 9
- embedded point, 121, 129
- enumerative formula, 85, 87–88
 - for singular elements of linear series, 258
- enumerative geometry, 62
 - 19th century achievements, 2
 - applications of intersection theory, 289
 - aspects of problems, 88
 - used to analyze geometric questions, 564
- enumerative problem, 86
 - generality, 88
 - geometry of set of solutions, 87
 - negative expected dimension analog, 397
 - steps to solving, 86–88
- étale equivalence relations, 575
- étale topology, 344
- Euler characteristic, *see also* topological Euler characteristic
 - constancy for a sheaf on a family, 534–535
 - of a coherent sheaf, 482
 - of the structure sheaf of a smooth projective threefold, 508
 - via the Todd class, 487
- Euler sequence, 97–99, 179
- evaluation map, 582–583
- exceptional divisor, 36, 57, 60, 61, 72
 - normal bundle, 472
 - of a blow-up of \mathbb{P}^n along a linear space, 338, 357
- excess intersection
 - elementary examples, 447–452
 - in a subvariety, 465–466
 - of a pullback to a subvariety, 469–470
 - of hypersurfaces, 464–465, 477, 478
 - via blowing up, 479
 - of three surfaces intersecting in a curve and a 0-dimensional scheme, 445, 449–451
 - via blowing up, 475–476
- excess intersection formula, 5, 292, 446, 454–456
 - applied to the five conic problem, 462–464
 - does not extend to irreducible components, 458
 - for a pullback via an inclusion, 470
 - for cycles on a subvariety, 465
 - via specialization to the normal curve, 468–469
 - for hypersurfaces, 465
 - for three surfaces meeting in a curve and a 0-dimensional scheme, 451
 - for Veronese surface in \mathbb{P}^5 , 478
 - heuristic explanation, 456–458
 - statement, 454–455
 - three-surface case restated in general form, 452–453
 - utility, 455–456
- excision, 25
- expected codimension, 19, 187
 - of a degeneracy locus, 427
- expected dimension, 446
 - negative, 396
 - of the secant variety, 371
- family of bundles, 489
 - on \mathbb{P}^1 , 494–497
- family of divisors, 571–576
- family of lines, 230–233
- Fano scheme, 193, 194
 - as special case of Hilbert schemes, 201
 - bounds on k, n necessary to obtain expected dimension, 237
- Chern class, 198
- definition, 196–197
- dimension, 194
 - bound via the normal bundle, 209
 - expected dimension, 195
- expression via the Grassmannian, 198–199
- has expected dimension when $d \ll n$, 237
- of 2-planes on a quadric, 240
- of a cubic surface
 - with one ordinary double point, 234–236
 - with two ordinary double points, 241
 - must have ≤ 4 isolated singular points, 241
- of a smooth cubic surface, 226
- of a smooth degree-4 hypersurface, 238
- of a smooth quintic threefold, 228
- of a smooth surface, 212
- of lines on a quadric surface, 197
- of lines on a smooth hypersurface, 226, 243
- of planes of maximal dimension, 222
- of the Fermat quartic in \mathbb{P}^4 , 226
- potential generalization, 239
- reducedness, 197, 208, 228, 238
- scheme structure, 196–197
- smoothness and the normal bundle, 208–212
- the nonsmooth case, 234–236, 241, 242
- universal property, 203
- upper bound on dimension, 197
- Zariski tangent space, 208
- Fermat surface
 - Fermat quartic, 238, 243
 - hyperflexes, 422
 - lines contained in, 421
- fiber of a vector bundle, 10
- fine moduli space, 575
 - of degree- d line bundles, 573, 575–576
- first-order deformations, 212–219
 - geometric view of lines on a cubic, 218
 - identification with global sections of the normal sheaf, 214–215
 - identification with morphisms from a double point, 213–214
 - utility in identifying tangent spaces, 213
 - vector space structure, 215–216
- five conic problem, 3, 289
 - answer, 308
 - generalization, 321
 - naive approach, 290–291

- transversality in the variety of complete conics, 303–306
- via blowing up the excess locus, 291, 480
- via the excess intersection formula, 292, 462–464
- via the space of complete conics, 302–308
- flag bundle, 347
- flag variety, 125, 159–160
 - Chow ring, 356, 359
 - class, 356
- flecnodal locus, 398–401
 - degree, 399
 - general point of a smooth surface of degree ≥ 3 is not flecnodal, 399
 - geometry, 421
- flecnodal, 398
- flex, 266
 - flex tangent, 266
 - flexes approaching a cusp, 423
 - hyperflex, 244, 266, 271–272
 - in a general pencil of curves, 405
 - in a pencil of plane curves, 287
 - in families, 401
 - of a family of curves, 405–408
 - of a general curve, 286
 - of a plane curve, 270–271
 - alternate notion, 271
 - via defining equations, 401–409
- flex line, 401–402
 - geometry of the curve of flex lines, 408–409
 - hyperflex lines, 403–405
 - on a curve defined by a homogeneous form, 402–403
- flip, 315
- footprint of a subvariety, 333
- form of type (p, q) , 547
- frame manifolds, 561
- fundamental class, 544
 - of a codimension- k subvariety, 549
 - of a scheme, 22
- fundamental class map, 545
 - applied to the Chern class, 559
 - image, 545, 552
 - is a ring homomorphism, 545
- fundamental cycle, 544
- GAGA theorems, 543–544
- Gauss map, 49–50, 218, 219, 263, 439
 - definition, 263–264
 - from a surface to its dual, 260
 - of a hypersurface is either birational or has positive-dimensional fibers, 381
 - of a smooth hypersurface is finite and birational, 381
- general polynomial of degree $2m - 1$
 - unique expression as a sum of m d -th powers, 362, 376–377
- general position lemma, 370
- general quadratic polynomial has no rational solution, 346, 356
- generality, 9, 88
 - of a curve, 565–566
 - of a rational curve, 441
 - very general, 9
- generalized principal ideal theorem, 11, 363
- generic finiteness, 28
- generic transversality, 46
 - definition
 - for a subvariety and a function, 30
 - for subvarieties, 18
 - is stronger than dimensional transversality, 33
 - of Schubert cycles, 139
 - reasons for nontransversality, 516–517
 - to a cycle, 512–517
 - to a morphism, 518–519
 - necessity of char 0, 518
 - sufficient condition, 518
- generically finite morphism
 - degree, 470–471
- genus formula, 69–70
 - applications, 70–71
 - for singular curves, 74
- geometric genus, 74
- Giambelli's formula, 157–159
 - in conjunction with Pieri's formula, 158–159
 - inductively via Pieri's formula, 164
 - statement, 158
- graph of a map, 54–55
- Grassmannian
 - as Hilbert scheme, 96, 201
 - Chow ring, 137
 - generators and relations via Chern classes, 183–187
 - is generated by special Schubert classes, 158
 - class of the pullback to the product of Grassmannians, 189
 - covering by affine spaces, 92–94
 - cut out by quadrics, 91, 94, 125
 - definition, 89
 - degree, 150, 164
 - differential of a morphism into the Grassmannian, 99–100
 - generalizations, 159–160
 - Lagrangian Grassmannian, 160
 - natural identification with Grassmannian of dual space, 89, 134
 - notation, 89
 - of lines in \mathbb{P}^3
 - Chow ring, 105–109
 - of lines in \mathbb{P}^n , 126, 147–150
 - degree, 131, 149–150
 - of 2-planes, 91–92
 - orthogonal Grassmannian, 160
 - smoothness, 91
 - tangent bundle, 96–99
 - expression via the universal bundles, 96
 - tangent space, 129
 - via the universal property, 100–102
 - universal bundles, 95–96
 - universal property, 201–203
 - universal quotient bundle, 95
 - Chern class, 178

- is globally generated, 178
 - universal subbundle, 95
 - Chern class, 178–179
 - existence, 95
 - lacks nonzero global section, 178
 - used to define Fano schemes, 198–199
- Grassmannian bundle, 346
 - Chow ring, 346–347
- Grothendieck ring of vector bundles, 484
- group of cycles, 15
- “hairy coconut” theorem, 179
- hard Lefschetz theorem, 558
- Hessian, 271
- higher direct image, 489, 498, 528–532
 - computation via Serre’s coherence theorem, 531
 - definition*, 528–529
 - is coherent for a projective morphism and coherent sheaf, 530
 - natural map to Čech cohomology group, 529
 - properties, 529–530
 - relation to direct image, 528
- higher direct image functors, 528
- Hilbert polynomial, 35, 45, 204–207
 - of a hypersurface, 206
 - subschemes with polynomial $2m + 1$, 350–352
- Hilbert scheme, 3, 83, 201, 203–207, 310–312
 - advantages, 311–312
 - as better compactification than symmetric power, 370
 - construction, 205–207
 - definition* and uniqueness, 204–205
 - disadvantages, 312
 - extraneous components, 312
 - mysterious closure of locus of smooth curves, 312
 - of a hypersurface, 206
 - of conics and cubics in \mathbb{P}^2 , 311
 - of divisors on a smooth scheme identified with symmetric power, 575–576
 - of hypersurfaces in \mathbb{P}^n , 207
 - of plane conics in \mathbb{P}^3 , 311
 - of subschemes of \mathbb{P}^3 with Hilbert polynomial $2m + 1$, 350–352
 - as projective bundle, 350
 - of twisted cubics, 311
 - singularities, 312
- Hilbert series
 - of a graded complete intersection, 185
- Hodge bundle, 503
 - of a pencil of quartics, 505, 510
- Hodge conjecture, 545, 549–550
 - integral codimension-1 case, 550
- Hodge decomposition, 546–548
 - algebraic computation of $H^{p,q}(X)$, 547–548
- Hodge diamond, 548–549
 - of a smooth quartic surface, 548
 - symmetries, 549
- Hodge number, 549
 - as invariant, 549
- Hodge structure, 548
 - polarization, 548
- Hodge–Riemann bilinear relations, 558–559
- holomorphic
 - map is algebraic, 544
 - subvariety is algebraic, 543
- homological equivalence, 545
- hook formula, 164
- Hopf index theorem, 506
- hyperelliptic curves, 570
- hyperplane
 - contact with a curve, 244, 265
- hypersurface
 - criterion to contain a line, 240
 - in \mathbb{P}^4 containing a complete intersection, 192
 - lines having point of contact of order 7, 420
 - of sufficiently low degree contains lines, 226
- ideal sheaf
 - of three points in \mathbb{P}^2 , 521–522
 - direct image, 526
 - direct image complex, 534–535
 - higher direct images, 532–533
 - of two points in \mathbb{P}^2 , 521
- inflection point, 265–273
 - weight on a general curve, 268
- integrals of algebraic functions, 571–572
- interpolation problem, 373
- intersection multiplicity, 14, 31–33
 - coinciding with the order of contact, 389
 - connection with multiplicity of a scheme at a point, 36
 - definition* and existence, 32
 - in the Cohen–Macaulay case, 32
 - necessity, 19
 - of a curve and a Cartier divisor, 265
 - of a divisor and a subvariety, 447–448
 - Serre’s formula, 48
 - via blow-ups, 61–62
- intersection number, 69
- intersection product, 15, 20, 41, 76
 - correction terms, 47–48
 - existence, 7, 19
 - for curves on surfaces, 68–74
 - in Chow ring corresponds to cup product in cohomology, 545
 - necessity of smoothness, 20
 - of a Cartier divisor and a subvariety, 447
 - geometric view, 447–448
 - of cycles on a proper subvariety, 465–466
 - on singular varieties, 38, 75–77, 455
 - semi-refined version via the strong moving lemma, 511
 - via the basic moving lemma, 511
- intersection theory
 - applications to enumerative geometry, 289
 - dependence on the moving lemma, 511, 512
 - goals, 14–15
 - influence on algebraic geometry, 1–2, 5
 - motivation, 1–2
 - on algebraic stacks, 502
- invariants of families of curves, 502–504

- degree of the Hodge bundle, 503
- for a pencil of plane quartics, 504–505
- inequalities, 504, 510
- number of nodes, 503
- of a pencil of curves of bidegree (a, b) , 510
- of a pencil of plane curves, 510
- of a pencil of plane sections of a smooth surface, 510
- relations, 504
- self-intersection of the relative canonical divisor, 504
- irreducible component, 9
- isotropic, 155
- Jacobi inversion theorem, 573
 - classical form, 592
- Jacobian, 571–572
 - Chow vs. cohomology rings, 583
 - cohomology ring, 583–584
 - classes of interest, 584
 - cotangent space, 574
 - definition, 572
 - identified with linear equivalence classes of effective divisors, 573
 - isomorphism with $\text{Pic}^d(C)$, 575
- join, 9
- Jordan–Hölder theorem, 11
- jumping lines, 225, 493–494
 - examples, 501–502
 - on a bundle of rank 2 on \mathbb{P}^2 , 497–502
 - on a bundle of rank 2 on \mathbb{P}^3 , 508–509
- K3 surface, 192
 - curves on a very general K3 surface, 580
- Kleiman’s transversality theorem, 20–21
- Kontsevich space, 292
 - and tangency conditions, 317
 - application, 322
 - birationality to the Hilbert scheme, 313
 - cross-ratio, 318
 - disadvantages, 316–317
 - extraneous components, 316
 - introduction, 312–313
 - is proper, 313
 - mysterious closure of locus of smooth curves, 317
 - number of PGL_4 -orbits, 322
 - of plane conics, 314
 - in \mathbb{P}^3 , 315
 - of plane cubics, 315–316
 - is not irreducible, 315
 - of rational plane curves, 317–321
 - of twisted cubics, 316
 - singularities, 317
- Krull’s principal ideal theorem, 16, 23
 - statement, 11
- Künneth formula, 51
- Lefschetz $(1, 1)$ -theorem, 550
- Lefschetz decomposition, 558
- Lefschetz hyperplane theorem, 182, 222, 228, 553–554
 - applied to complete intersections, 554–557
 - extensions, 557
 - statement, 553
- Lefschetz principle, 245, 277
- lexicographic ordering, 207
- liaison, 71
- line bundle
 - generated by global sections, 362–363
 - on the projectivization of a bundle, 327–328
 - products of line bundles, 191
 - restriction to pullback, 527–528
 - tautological line bundle of a projective bundle, 171
 - twisting a vector bundle, 335, 355, 356
- linear series, 10, 566
 - complete, *see* complete linear series
 - discriminant of a very ample linear series, 258
 - maps to \mathbb{P}^n given by a linear series, 568–570
 - by a general series, 567
 - embeddings, 570–571
 - existence, 576
 - number, 577, 590
 - present on a general curve, 570–571
 - singular elements, 258–265
 - characterization of tangency of hyperplanes, 265
 - via the topological Hurwitz formula, 277
 - singular elements of a pencil, 259–260
- linear subspaces
 - characterized by Fano scheme, 197
 - characterized by Hilbert polynomial, 204
- linearization, 5, 166
- lines
 - and curves in \mathbb{P}^3 , 110–115
 - and surfaces in \mathbb{P}^3 , 122–125
 - have no sixth order contact with a general surface, 420
 - meeting a curve, 111–112
 - via specialization, 120–121, 128
 - meeting a curve in \mathbb{P}^4 , 161
 - meeting a line on a quadric, 286
 - meeting a smooth rational curve four times, 426
 - meeting a surface in \mathbb{P}^4 , 161
 - meeting a surface to high order, 390–391, 420
 - meeting four curves, 85, 112
 - transverse intersection of cycles, 127–128
 - meeting four lines, 85, 110–111, 127
 - meeting four planes, 131, 150, 162
 - on a complete intersection, 240
 - on a cubic surface, 199–201, 212
 - geometric viewpoint via first-order deformations, 218
 - the smooth case, 212
 - via the map α and Bertini’s theorem, 221
 - on a cubic with a double point, 234–236
 - on a hypersurface, 240
 - on a pencil of quartic surfaces, 231–233
 - alternative approach, 233–234
 - on a quadric, 122–123
 - on a quintic threefold, 193, 227–229
 - in algebraic geometry and string theory, 228
 - on a smooth cubic, 165, 166, 493
 - on a smooth hypersurface
 - odd behaviors, 243
 - on a smooth surface in \mathbb{P}^3 , 396–399
 - bounds, 399, 421

- number, 422
- on the intersection of two quadrics, 131, 157
- tangent to a surface, 161
- tangent to four quadrics, 85, 125
- tangent to four spheres, 480
- linked curves in \mathbb{P}^3 , 70–71
- Littlewood–Richardson coefficients, 143
 - appearing with multiplicity > 1 , 164
- locally closed subscheme, 9
- loci in space of plane cubics, 81–83
- locus of d -fold lines, 479
- locus of m -tuples of points of a curve lying on a plane, 377
- locus of asterisks, 65–66, 82
- locus of bundles of splitting type a , 495
 - as pullback of strata in a family of vector bundles, 496
- locus of Castelnuovo curves, 571
- locus of chords tangent to a curve, 113
 - alternative characterization, 129
 - class, 114
- locus of cones, 83, 257
- locus of conics containing a point in \mathbb{P}^3 , 360
- locus of cubics of the form $2L + M$, 82
- locus of curves tangent to a smooth curve
 - degree, 479
 - multiplicity along locus of d -fold lines, 479
- locus of curves with a triple point, 244, 257
- locus of degeneracy, *see* degeneracy locus
- locus of double lines, 81
 - class, 360
- locus of flecnodal curves
 - dimension and irreducibility, 422
- locus of hyperflex lines, 403
- locus of jumping lines, 494
 - as a nonsingular cubic curve, 509
 - degree in the even first Chern class case, 499
 - degree in the odd first Chern class case, 501
 - for a bundle of rank 2 on \mathbb{P}^3 , 508–509
 - of a bundle defined by a bilinear form, 509
 - scheme structure, 499, 501
- locus of matrices of rank $\leq k$
 - degree, 426, 433–436
- locus of planes on a quadric, 155–157
 - class, 157
- locus of reducible cubics, 64–65, 81
- locus of reducible cubics composed of a smooth conic
 - with a tangent line, 83
- locus of secant planes, 369
 - class, 378–380
 - improper secants, 385
- locus of singular conics, 81, 360
- locus of singular plane cubics, 62–66, 409
- locus of singular plane curves, 412, 423–424
- locus of smooth conics with a tangent line, 83
- locus of smooth curves
 - closure in the Hilbert scheme, 312
- locus of smooth curves with a hyperflex, 287
- locus of special lines, 444
- locus of triangles, 65, 82
- locus of triple lines, 82
- locus of trisecant lines, 443
- locus where global sections do not generate a bundle, 366
 - codimension and class, 363
- Macaulay2*, 2
 - applied to direct image complex, 534, 541–542
 - calculation of $c_1(\mathcal{E})$, 233
 - calculation of $c_4(\Phi(3, 3, 1))$, 232
 - calculation of $c_4(\text{Sym}^3 S^*)$, 200
 - calculation of $c_{d+1}(\text{Sym}^d S^*)$, 227
- maps from a curve to \mathbb{P}^n , 565–566
 - all curves vs. general curves, 565
 - birationally very ample maps, 569–571
 - correspond to pairs (\mathcal{L}, V) , 566
 - ease of finding high-degree maps and embeddings, 566
 - embedding in \mathbb{P}^3 , 564, 567
 - existence condition, 566, 576
 - to \mathbb{P}^1 , 564
 - to \mathbb{P}^2 , 564, 567
- Mayer–Vietoris, 25
- method of undetermined coefficients, 53–55, 79, 80, 107, 156
 - applied to the cycle of plane conics in \mathbb{P}^3 meeting a line, 349, 359
 - applied to the Grassmannian, 143–144
 - applied to the product of Grassmannians, 189
- minimal model program, 552
- minor, 90
- mirror symmetry, 228
- moduli space, *see also* fine moduli space
 - parametrizing smooth projective genus- g curves, 566
- moduli stack of stable curves, 503
- moving lemma, 7, 19–21
 - basic version, 511
 - via the cone construction, 512–517
 - bypassed via the Fulton–MacPherson approach, 512
 - direct proofs of the second part, 512
 - failure for singular varieties, 75
 - necessity of smoothness, 20
 - proof of basic version, 511–512
 - statement*, 19
 - strong version, 511
 - when one cycle is a first Chern class, 38
- multiplicity
 - of a hypersurface in smooth variety, 34
 - of a scheme, 15
 - of a scheme at a point, 33–36
 - connection with intersection multiplicity, 36
 - of a variety at a point, 411
 - of an intersection, 14
- Mumford relation, 504
 - in the case of a pencil of plane quartics, 505
 - proof via Grothendieck Riemann–Roch, 506–507
- Nakayama’s lemma, 101
- nested pairs of divisors, 55–56, 339–340
 - as projective bundle, 340
- net of cubic surfaces, 240
- net of curves

- binodal elements, 425
- geometric view, 417–419
- net of plane curves, 273–276
 - cusps, 413–419
 - discriminant, 274–276
 - flecnodes, 422
 - hyperflexes of quartics, 405
 - of cubics, 389, 409
 - singular points, 244, 275
- node, 411
- Noether's formula, 483
- nondegenerate map, 565
 - degree, 565
- nondegenerate quadratic form, 34
- nontransversality, 47–48
- normal bundle, 180, 208
 - computation, 209
 - definition*, 40
 - for arbitrary schemes, 209
 - lines on a hypersurface with prescribed normal bundle, 225
 - of k -planes on hypersurfaces, 219–220
 - of complete intersections, 212
 - of the diagonal, 249
 - of the exceptional divisor, 472
 - used for excess intersections, 447, 451
- numerical equivalence, 552, 565
 - and the Hodge conjecture, 552
 - yields ring structure, 552
- one-parameter family, 10
- order of contact, 265, 389
 - of a plane curve and a hyperflex, 422
- order of vanishing, 23
- ordinary m -fold point, 72
- ordinary cusp, 409–411
- ordinary double point, 34
- ordinary tacnode, 411
- ordinary triple point, 256
- oscnode, 414
- osculating plane, 267
 - rotation, 381, 388
- parallel transport, 561
- parameter space, 1, 62
 - alternative choices, 292–293
 - as projective bundle, 348–349
 - Chow ring of a parameter spaces, 62
 - desired attributes, 86
 - necessity of compactification, 289–290
 - of conics in \mathbb{P}^3 , 348–349
 - of curves, 310–317
 - utility in enumerative problems, 289
- pencil of cubic surfaces, 193
- pencil of curves
 - on a quadric surface, 285, 288, 510
 - on a surface, 260–262, 279–280
 - singular at a point, 286, 288
- pencil of plane curves, 422–423
 - curve traced out by flexes, 389, 405–409, 422–423
 - flex lines through a point, 422
 - hyperflexes, 389, 422
 - invariants, 504–505, 510
- pencil of plane sections of a smooth surface, 510
- pencil of quartic surfaces containing a line, 193, 231–233
- Pfaffians, 92
- Picard group, 37
 - of degree- d line bundles, 567
 - class of subvarieties, 585
 - is a fine moduli space, 573, 575
 - is an algebraic variety, 575
 - of the space of complete conics, 306
 - tangent space can be identified with cohomology, 544
- Pieri's formula, 145–147
 - for other special Schubert classes, 154–155
 - interpreted via Young diagrams, 154
 - statement*, 145
- pinch point
 - of a projection of the Veronese surface, 439
 - of a smooth surface in \mathbb{P}^3 , 442
 - of the projection of a cubic scroll, 442
 - of the projection of a rational normal surface scroll, 442
 - of the projection of a smooth surface, 436–440
- plane conics in \mathbb{P}^3
 - form a projective bundle, 347
 - locus of conics meeting a given line, 348
 - meeting eight general lines, 323
 - finite expected answer, 347
 - must be smooth, 352
 - outline of proof, 347–348
 - solution, 354
- plane curve
 - cusps, 389, 409
 - singularities, 410–412
 - triple points, 256–257
- plane sextic with four nodes, 592–593
- planes on the intersection of two quadrics, 157
- Plücker coordinates, 90
 - ratios as determinants of submatrices, 94
- Plücker embedding, 89–92, 131, 229
 - image, 90–91
 - Schubert cycles as intersections, 138
- Plücker formula, 268–270
 - analog in higher dimension, 272–273
- Plücker formula for plane curves, 418
- Plücker relations, 91
 - for the Grassmannian of 2-planes, 92
 - general case, 94
- Poincaré bundle, 575, 580–582
 - as direct image, 581
 - pushforward, 587–589
 - Chern class, 589–590
- Poincaré duality, 15
 - for the Chow ring of projective space, 45
- Poincaré's formula, 565, 585–586
- Poincaré–Hopf theorem, 181
- Porteous' formula, 427–429
 - applied to m -secant lines to curves, 444
 - applied to pinch points of a projection, 438

- applied to quadrisecant lines, 440
- as general case of Theorem 10.2, 364
- for $M_0(\varphi)$, 429–430
- geometric applications, 433–442
- linearizing the problem, 431–432
- reduction to a generic case, 430–431
- statement*, 429
- used to calculate class of the Brill–Noether locus, 591
- primary decomposition, 9
- primitive cohomology groups, 558
- principle of specialization, 462
- projection of a smooth curve
 - singularities, 128
- projection of a smooth surface, 436
 - pinch points, 426, 438–440
 - singularities, 436–437
- projection of the Veronese surface, 439
- projective bundle, *see also* vector bundle, projectivization
 - can be written as projectivization, 324
 - characterization, 329
 - Chow ring, 331–335
 - definition*, 323
 - is a Brauer–Severi variety, 344
 - is the projectivization of a bundle, 327–330
 - over \mathbb{P}^1 , 324–327
 - over an arbitrary scheme, 331
 - pushforward, 332
 - utility, 363
 - weakening of definition, 344
- projective tangent space, 10
- projectivization
 - of a bundle, 96, 324
 - degree, 335
 - identifying isomorphic projectivization, 330
 - recovering original data, 327–328
 - of a subbundle, 340–341
 - class, 341
 - line subbundle, 341
 - normal bundle, 341
 - of a vector space, 9
- projectivized tangent cone, *see also* tangent cone, 61
 - characterization via blow-ups, 36
 - extension to arbitrary schemes, 35
 - of a hypersurface, 34
- proper transform, 57, 72
 - class, 61
- Prym map, 471
- pullback, 29–31
 - existence and uniqueness, 30
 - flat, 31
 - flat pullback, 25
 - general definition via the excess intersection formula, 456
 - is not defined on a singular variety, 77
 - to a subvariety, 469–471
- push-pull formula, 30
- pushforward
 - for a projective bundle, 332
 - from the Grassmannian bundle, 432–433
 - of a cycle, 28
 - proper pushforward, 28–29
- quadric cone, 479
- quadric line complex, 192
- quadric surface
 - curves lying on a smooth quadric, 69–70
 - defined by a symmetric map $V \rightarrow V^*$, 297–298
 - intersection of quadrics containing a linear space, 445, 460–462
 - linear subspaces, 155–157
 - ruling, 53, 69
 - tangency of two smooth quadrics, 298
 - two rulings, 123
- quadrisecant
 - condition to be simple, 444
 - to a curve in \mathbb{P}^3 , 377
 - to a curve of higher genus, 441–442
 - to a general rational curve is simple, 444
 - to a rational curve, 387
 - to a rational space curve, 440–441, 444
- quartic curve
 - double at five specified points, 385
 - hyperflexes of a plane quartic, 389, 405
 - reducible quartics in \mathbb{P}^2 , 83
- quartic surface
 - containing a conic, 361
 - containing a line, 165
 - double at nine specified points, 386
 - in \mathbb{P}^3 , 242
- quasi-affine stratification, 27, *see also* affine stratification
- quasi-isomorphism, 534
- quintic surface
 - homological equivalence of curves on a smooth threefold, 551
 - lines on a quintic threefold, 3, 193, 227–229
 - lines with high-order contact, 389–391, 394–396
- ramification
 - divisor, 278
 - index, 278
 - points on \mathbb{P}^1 , 287
 - sequence, 266, 268
- rational curve
 - in projective space via the Kontsevich space, 317–321
 - is the projection of a rational normal curve, 373
 - on a hypersurface, 239
 - quadrisecant lines via Porteous’ formula, 440–441, 444
- rational equivalence, 16
 - definition*, 16
 - failure to preserve genera, 45
 - generation by divisors of rational functions, 23
 - of two 0-cycles on a curve, 24
 - preservation by pushforward, 29
 - via divisors, 22–24
- rational normal curve, 324, 355
 - abstract secant variety, 374
 - characterized by independence of divisors, 386
 - curve of pure d -th powers, 376
 - independence lemma, 373

- passing through seven general points, 386
 - secant variety
 - degree, 375–376
- rational normal scroll, 53, 324–327, 355
 - degree, 335
 - pinch points, 442
- rational quartic
 - containing 11 points, 289, 321
- rational quartic containing 11 points, 318
- reducible cubic, 64
- Rees algebra, 471
- reflexivity, 380–382
- regular 1-forms, 481
- relative duality, 328
- relative dualizing sheaf, 503
- relative Euler sequence, 394
- relative tangent bundle, 393
 - of a projective bundle, 393–394
 - Chern classes, 394
- representable functors, 205
- resultant, 427
 - formula via Δ_f^e , 428–430
- Riemann–Hurwitz formula, 68, 278
- Riemann–Roch theorem
 - applied to linear series on curves, 567–568
 - for smooth curves, 481–482
 - for smooth projective surfaces, 483
 - Grothendieck vs. Hirzebruch, 489
 - Grothendieck’s version, 489–490
 - applied to $\mathcal{E} = \text{Sym}^3 \mathcal{S}^*$, 490–493
 - applied to bundle of principal parts, 508
 - for a submersion, 490
 - Hirzebruch’s version, 488–489
 - reduces to classical versions, 508
 - motivation, 481
 - original formulation, 481–482
 - produces high-degree maps and embeddings, 566
- ruled surface, 341–343
 - containing curve of negative self-intersection, 323, 341–342
 - sections, 359
- Sard’s theorem
 - algebraic version, 519
- scheme, 9
- Schubert calculus, 2, 131
 - by static specialization, 115–117
- Schubert cell, 135
 - in $\mathbb{G}(1, 3)$, 102, 104
 - tangent space, 135
- Schubert class, 160
 - and the Chern class, 134
 - as fundamental invariant, 134
 - basis for Chow ring, 142–143
 - closed-form expression for multiplication, 148
 - combinatorial formula for multiplication, 149
 - counting via Young diagram, 153
 - definition, 132
 - notation, 132
 - product formula, 143
 - relation among special classes, 147
 - representation via Young diagram, 152
 - special classes, 145, 155
 - generate Chow ring of Grassmannian, 158, 183
- Schubert cycle, 4, 132
 - common cases, 132–133
 - definition, 132
 - equations, 138–139
 - generic transversality, 139
 - in $\mathbb{G}(1, 3)$, 102–103
 - intersection in complementary dimension, 141–142
 - notation, 103
 - benefits, 133–134
 - partial ordering, 133
 - relative to transverse flags, 140
 - special cycles, 133
 - tangent space, 108–109, 126
- Schubert index, 135
 - dual index, 142
- Schubert variety, *see* Schubert cycle
- scroll, 388
- secant line, 113, *see also* secant variety
 - stationary, 128
- secant plane, *see* secant variety
- secant plane map, 369
 - birationality, 370, 385
 - composed with the Plücker embedding, 378, 386
 - extends to an embedding, 374
 - improper secants, 385
 - is never regular for $n, m > 1$, 370
- secant variety, 367–371
 - abstract, *see* abstract secant variety
 - definition, 370
 - dimension, 370
 - expected dimension, 371
 - general point lies on unique secant plane to a curve, 386–387
 - of a curve of positive genus, 380
 - of a curve that is not a rational normal curve, 377–380
 - of a rational normal curve, 373–377
 - degree, 375–376
 - of an elliptic curve, 388
 - of the Veronese surface, 439
 - proper, 386
 - secant plane, 367
 - used to study pure d -th powers, 376–377
- second fundamental form, 243, 261–265
 - of a smooth hypersurface, 264
- section, 58
- Segre class, 350
 - as locus where global sections do not generate a bundle, 362, 363
 - definition, 363
 - generalized definition, 453–454
 - gives degree of the variety swept out by a linear space, 367, 385
 - is the reciprocal of the Chern class, 364
 - of the dual of a bundle, 364

- parallel with Chern class, 364
- used to obtain degrees of secant varieties, 375
- Segre map, 78
- Segre variety
 - as determinantal variety, 434
 - definition*, 52
 - degree, 52–53
 - Segre threefold, 286, 326
- self-intersection, 83
 - of a 2-plane on a fourfold, 478
 - of a curve on a surface, 70
 - of the zero section, 343–344
 - question of boundedness below, 359
- Serre duality, 328, 482
- Serre's coherence theorem, 530
- Serre's formula, 48
- sheaf, 9
- sheaf with specified fiber, 520
 - for 3-point ideal sheaf example, 522
- singular curve
 - conic meeting seven general lines in \mathbb{P}^3 , 360
 - cubic, 62
 - in a general pencil, 165, 244, 253, 279–280
 - of conics, 253
 - of higher degree, 64
 - on a quadric, 285
- singular elements
 - of linear series, *see* linear series, singular elements
- singular hypersurface, 34, 245–247
 - in a general pencil of hypersurfaces, 253
- singularity
 - of plane curves, 410–412
 - of plane sections of a general surface, 420
 - of the curve traced out by flexes of a pencil, 423
- smooth curve
 - conic tangent to a singular curve, 321
 - conic tangent to five conics, *see* five conic problem
 - conic tangent to five general curves, 321
 - conics degenerating to a double line, 294–295
 - conics degenerating to a rank-2 conic, 294
 - in \mathbb{P}^3 as intersection of three surfaces, 445, 452
 - quintic of genus 2 is the intersection of three surfaces, 477
 - quintics lying on surfaces, 83
 - with no inflection points is the rational normal curve, 287
- smooth hypersurface
 - cannot contain a plane of more than half its dimension, 222
 - containing a 2-parameter family of lines, 193, 238
 - quintic containing a 2-plane, 243
- smooth locus of discriminant, 284
- smooth plane curve
 - divisor of a conic, 298
 - genus, 69
 - smooth cubics, 62
- smooth rational curve
 - meeting lines four times, 426
 - quintic as intersection of three surfaces, 477
- smooth surface
 - class of a curve squared, 559
 - containing a curve
 - points of tangency, 445, 476
 - containing infinitely many irreducible curves of negative self-intersection, 343, 358
 - finitely many hyperplane sections with triple points, 420–421
 - finitely many lines on, 70, 83
 - of degree 3 in \mathbb{P}^4
 - lies on no smooth hypersurface, 556
- snake lemma, 187
- socle, 184
- space of complete conics, 290, 292
 - as compactification of smooth conics, 293
 - Chow ring, 306–309
 - classification of smooth conics, 296, 301
 - complete conic tangent to five general conics is smooth, 302–303
 - divisor class of complete conics tangent to a conic, 307–308
 - equations, 299–301
 - informal introduction, 293–296
 - is smooth and irreducible, 299
 - other divisor classes, 308–309
 - relation to blow-up, 301–302
 - rigorous description, 296–301
 - smooth complete conics, 296
 - used in solution of five conic problem, 302–308
- space of complete quadrics, 309–310
 - stratification, 310
- special divisors, 569
- special secant plane, 377
 - examples, 377–378
 - expected dimension, 387
- specialization, 62, 115–122
 - appearance of multiplicities, 121–122, 129
 - dynamic, *see* dynamic specialization
 - introductory example, 115
 - relations between singular plane cubics, 63
 - static vs. dynamic, 116–117
 - utility in projective space, 117
- specialization to the normal cone, 5
- specialization to the normal curve, 466–468
 - applied to excess intersection of cycles on a subvariety, 468–469
- sphere in complex projective space, 480
- splitting principle, 172–173
 - splitting construction, 172
 - statement*, 172
 - with Whitney's formula, 173–174
- splitting type, 494
- stability of fibers, 502, 503
- stable map, 313
- Steiner construction, 460
- stratification, 27
 - of \mathbb{P}^9 , 62–64
 - of the space of complete quadrics, 310
- strict transform, *see* proper transform

- swallowtail singularity, 247
- sweeping out
 - by a subscheme, 217
 - bound on dimension of tangent space, 217
 - by linear spaces, 144–145, 161, 366–367
 - degree via the Segre class, 367
 - by lines on a pencil of hyperplane sections of a cubic, 240
 - by lines on a quartic threefold, 240
 - by lines with specified order of contact, 421
 - by the 2-planes of an irreducible surface, 443
 - by trisecants, 362, 379–380
 - by twisted cubic, 131, 144–145, 162
- symmetric power, 367–369
 - affineness and projectivity, 368
 - as a projective bundle, 587–589
 - is a fine moduli space, 575–576
 - maps to Jacobian, 572
 - of \mathbb{A}^1 , 368
 - of \mathbb{P}^1 , 368
 - smoothness, 368–369
- tacnode, *see also* ordinary tacnode, 424–425
- tangent bundle, 179
 - of projective space, 179
 - of projective spaces
 - is not the sum of line bundles, 192
 - to hypersurface, 179–180
 - to the Grassmannian, 96–99
- tangent cone
 - extension to arbitrary schemes, 35
 - of a hypersurface, 34
 - to the discriminant, 284
- tangent hyperplane
 - to a quadric, 297
 - to two smooth quadrics, 298
- tangent lines to a surface, 123–125
- tangent space
 - of a cycle of tangent conics, 304
 - of the Fano scheme, 208
 - to a Schubert cycle, 108–109
 - to cycle of plane conics in \mathbb{P}^3 meeting a given line, 353
 - to the discriminant, 282–284
 - of degree- d polynomials, 285, 304
 - to the Fano and Hilbert schemes, 208–227
 - to the Grassmannian, 129
 - to the Picard group can be identified with cohomology, 544
- tangent vector
 - rank, 97, 126
- tangential variety of a surface, 161, 286, 387, 439, 443
 - is 4-dimensional, 440, 443
- tautological bundle, 177–179
 - on the universal k -plane, 336
- tautological class, 336
- tautological family of plane conics, 348, 350
- tautological quotient bundle
 - of a Grassmannian bundle, 347
- tautological subbundle, 324
 - of a Grassmannian bundle, 346
 - Chern class, 347
- Terracini's lemma, 372–373
- theorem on cohomology and base change, 520
 - necessity of flatness in version 3, 535
 - proof via approximation of a complex and the vector bundle criterion, 535–540
 - version 1, 526
 - version 2, 531–532
 - version 3, 533
 - alternative formulation, 541
- theta divisor, 573, 585
- Todd class, 487–488
 - as polynomial in the Chern classes, 508
 - multiplicativity, 488
- topological Chern class
 - algebraic Chern class results carry over, 563
 - and curvature, 561–563
 - as obstruction to a nowhere-zero section, 560–561
 - coincides with Chern class for an algebraic vector bundle, 561
 - definition, 560–561
- topological Euler characteristic, 39, 179–182, 280, 482
 - additivity, 277
 - and multiplicity of the discriminant, 281
 - determines the middle Betti number, 182
 - of a blow-up of a surface, 181
 - of a hypersurface, 181–182
 - of a smooth hypersurface of bidegree (a, b) , 192
 - of projective space, 181
 - via top Chern class, 181
- topological genus, 482
- topological Hurwitz formula, 277–285
 - applied to pencils of curves on a surface, 279, 280
 - statement, 278
- total inflection, 268
- transversality
 - generic, 18
 - of eight cycles corresponding to general lines, 354
 - of five cycles in the variety of complete conics, 303–306
- transverse flags, 139
- transverse intersection
 - definition, 18
 - of Schubert cycles, 141
- triangle, 65
- triple point, 34
 - of plane curves, 256–257
- triples of collinear points, 79
 - class via Porteous' formula, 442
- trisecant
 - surface in \mathbb{P}^3 swept out by, 362, 379–380
 - to a curve in \mathbb{P}^3 , 377
 - to a curve in \mathbb{P}^4 , 378
 - to a rational curve in \mathbb{P}^4 , 362, 378–379
 - to a space curve, 443
- Tschirnhausen transformation, 412
- tubular neighborhood theorem, 466
 - fails to generalize, 466, 478
- twisted cubic

- common chords of two twisted cubics, 122
- positive-dimensional component, 480
- is the unique curve whose secants sweep out \mathbb{P}^3 once, 374, 386
- tangent to 12 quadrics, 2
- two surfaces intersecting in a curve and a 0-dimensional scheme, 445, 459–460
 - dependence on geometry of the surfaces, 460
 - examples, 478
- 2-planes meeting three quadrics, 164
- universal divisor, 576, 581
- universal family of conics in the plane, 345
- universal family of subschemes, 204
- universal Fano scheme, *see also* Fano scheme, 230
 - and families of lines, 229–234
 - classes of universal Fano schemes of lines on surfaces in \mathbb{P}^3 , 231
 - defining equations, 197
 - definition*, 194
 - dimension, 194
 - global view as the zero locus of a section of a vector bundle, 229–231
 - of lines on cubic surfaces, 230
 - reducedness, 230
- universal flex, 423
- universal hyperplane, 382
 - Chow ring, 336
- universal hypersurface, 359
- universal line
 - Chow ring, 337, 394
 - in \mathbb{P}^n , 356
- universal line bundle, 575, 580–582
- universal plane, 96, 125, 144, 159
 - as projectivization of the universal subbundle, 336
 - Chow ring, 335–336
 - class via Porteous' formula, 442
- universal property of Proj, 327
- universal singular point, 245, 254, 260
 - class, 273–274
 - is a complete intersection, 245
- universality of a map, 484
- vanishing sequence, 266
 - geometric meaning, 267
- variety, 9
- vector bundle
 - complete classification over \mathbb{P}^1 , 223–224
 - generated by global sections, 362
 - mysteries on higher-dimensional projective space, 224
 - of rank 2 on \mathbb{P}^2
 - with even first Chern class, 497–499
 - with odd first Chern class, 499–501
 - on \mathbb{P}^1 splits, 223
 - on projective space
 - behavior in families, 494–497
 - projectivization, 171, 172
 - twisting by a line bundle, 335, 355, 356
- vector field, 99
- Veronese map, 48, 78
 - relation to discriminant, 246–247
 - used to prove the moving lemma, 514
- Veronese surface, 82
 - as variety of minimal degree, 335
 - hyperplane section with triple points, 421
 - in \mathbb{P}^5 as defective variety, 371–372
 - in \mathbb{P}^5 is the intersection of \mathbb{P}^5 with a Segre variety, 478
 - projection from a general line, 439
- Veronese variety, 48, 78
 - degree, 48–49
 - tangent planes, 385
 - which are m -defective, 373, 385–386
- very ample line bundle, 424
- web of quadrics in \mathbb{P}^3 , 444
- Weierstrass point, 287
- weight of a point with respect to a linear system, 268
- Weil divisor, *see* divisor
- Whitney's formula, 169
 - for globally generated bundles, 187–190
 - with the splitting principle, 173–174
- Young diagram, 132, 152–154
 - transposition and Grassmannian duality, 153–154, 163
- Zariski tangent space, 34, 208
 - algebraic descriptions, 100–102
 - identified with first-order deformations, 213
 - of a local ring, 209