

Example Sheet 4

Isaac Martin

EXERCISE 1. Let D be a degree 0 Weil divisor on \mathbb{P}_k^1 . Construct a rational function f , i.e. a section of the structure sheaf over the generic point, such that the divisor associated to f is D . Deduce that every degree 0 divisor on \mathbb{P}_k^1 is principal.

Proof: Let $D = D_1 - D_2$, where D_1 and D_2 are two effective Weil divisors.

Recall first that any codimension one subvariety $Y = V(\mathfrak{a}) \subset \mathbb{P}_k^1$ is determined by a single homogeneous equation. To see this, suppose $f \in \mathfrak{a}$ is some nonzero homogeneous equation. Then $V(\mathfrak{a}) \subseteq V(f) \subsetneq \mathbb{P}_k^1$, hence $Y = V(\mathfrak{a}) = V(f)$.

We now suppose that D is a Weil divisor on \mathbb{P}_k^1 as in the question. Write $D = D_1 - D_2$ for some effective divisors D_1 and D_2 , where

$$D_1 = \sum n_i B_i \quad \text{and} \quad D_2 = \sum m_j C_j$$

for prime divisors B_i and C_j . For each i , $B_i = V(f_i)$ for some homogeneous irreducible polynomial by our earlier comment, and likewise $C_j = V(g_j)$ with g_j an irreducible homogeneous polynomial for each j . Define $f = \prod_i f_i^{n_i}$ and $g = \prod_j g_j^{m_j}$ and let $h = \frac{f}{g}$. Because $\deg(D) = 0$ we have by design that $\deg(f) = \deg(D_1) = \deg(D_2) = \deg(g)$, and hence $h \in K^*$ as it is a quotient of two homogeneous function of equal degree. We have that

$$(h) = (f) - (g) = \sum_i n_i B_i - \sum_j m_j C_j = D_1 - D_2 = D,$$

and therefore D is principal.

By starting with a rational function $\frac{f}{g}$ in K^* and factoring both the numerator and denominator into irreducible components, we similarly see that $(f) = 0$ and can then conclude that a Weil divisor D has degree 0 if and only if it is principal. \square