

# CSC 600 HOMEWORK 1 - SYNTAX

February 13, 2018

Homework is prepared by: Ilya Kopyl.

It is formatted in LaTeX, using TeXShop editor (under GNU GPL license).

## 1. Using BNF write the syntax definitions of the following objects:

a) Natural number (1, 2, 3, ...). The answer:

$$\begin{aligned}\langle \textit{natural number} \rangle &::= \langle \textit{non-zero digit} \rangle \mid \langle \textit{natural number} \rangle \langle \textit{digit} \rangle \\ \langle \textit{digit} \rangle &::= 0 \mid \langle \textit{non-zero digit} \rangle \\ \langle \textit{non-zero digit} \rangle &::= 1 \mid 2 \mid 3 \mid 4 \mid 5 \mid 6 \mid 7 \mid 8 \mid 9\end{aligned}$$

b) Unsigned integer (0, 1, 2, 3, ...). The answer:

$$\begin{aligned}\langle \textit{unsigned integer} \rangle &::= 0 \mid \langle \textit{non-zero digit} \rangle \langle \textit{digits} \rangle \\ \langle \textit{digits} \rangle &::= \langle \textit{digit} \rangle \mid \langle \textit{digits} \rangle \langle \textit{digit} \rangle \\ \langle \textit{digit} \rangle &::= 0 \mid \langle \textit{non-zero digit} \rangle \\ \langle \textit{non-zero digit} \rangle &::= 1 \mid 2 \mid 3 \mid 4 \mid 5 \mid 6 \mid 7 \mid 8 \mid 9\end{aligned}$$

c) Integer (... , -2, -1, 0, 1, 2, ...). The answer:

$$\begin{aligned}\langle \textit{integer} \rangle &::= \langle \textit{sign} \rangle \langle \textit{unsigned integer} \rangle \\ \langle \textit{sign} \rangle &::= + \mid - \mid \langle \textit{empty} \rangle \\ \langle \textit{empty} \rangle &::= \\ \langle \textit{unsigned integer} \rangle &::= 0 \mid \langle \textit{non-zero digit} \rangle \langle \textit{digits} \rangle \\ \langle \textit{digits} \rangle &::= \langle \textit{digit} \rangle \mid \langle \textit{digits} \rangle \langle \textit{digit} \rangle \\ \langle \textit{digit} \rangle &::= 0 \mid \langle \textit{non-zero digit} \rangle \\ \langle \textit{non-zero digit} \rangle &::= 1 \mid 2 \mid 3 \mid 4 \mid 5 \mid 6 \mid 7 \mid 8 \mid 9\end{aligned}$$

d) Odd number (... , -3, -1, 1, 3, ..., 101, ..., 2047, ...). The answer:

$\langle \text{odd number} \rangle$	$::= \langle \text{sign} \rangle \langle \text{unsigned odd number} \rangle$
$\langle \text{sign} \rangle$	$::= + \mid - \mid \langle \text{empty} \rangle$
$\langle \text{empty} \rangle$	$::=$
$\langle \text{unsigned odd number} \rangle$	$::= \langle \text{odd digit} \rangle \mid \langle \text{number} \rangle \langle \text{unsigned odd number} \rangle$
$\langle \text{number} \rangle$	$::= \langle \text{non-zero digit} \rangle \mid \langle \text{number} \rangle \langle \text{digit} \rangle$
$\langle \text{digit} \rangle$	$::= 0 \mid \langle \text{non-zero digit} \rangle$
$\langle \text{non-zero digit} \rangle$	$::= 2 \mid 4 \mid 6 \mid 8 \mid \langle \text{odd digit} \rangle$
$\langle \text{odd digit} \rangle$	$::= 1 \mid 3 \mid 5 \mid 7 \mid 9$

e) Even number (... , -4, -2, 0, 2, 4, ..., 332, ..., 1022, ...). The answer:

$\langle \text{even number} \rangle$	$::= \langle \text{sign} \rangle \langle \text{unsigned even number} \rangle$
$\langle \text{sign} \rangle$	$::= + \mid - \mid \langle \text{empty} \rangle$
$\langle \text{empty} \rangle$	$::=$
$\langle \text{unsigned even number} \rangle$	$::= \langle \text{even digit} \rangle \mid \langle \text{number} \rangle \langle \text{unsigned even number} \rangle$
$\langle \text{number} \rangle$	$::= \langle \text{non-zero digit} \rangle \mid \langle \text{number} \rangle \langle \text{digit} \rangle$
$\langle \text{digit} \rangle$	$::= 0 \mid \langle \text{non-zero digit} \rangle$
$\langle \text{non-zero digit} \rangle$	$::= 1 \mid 2 \mid 3 \mid 4 \mid 5 \mid 6 \mid 7 \mid 8 \mid 9$
$\langle \text{even digit} \rangle$	$::= 0 \mid 2 \mid 4 \mid 6 \mid 8$

f) Integer divisible by five (... , -10, 5, 0, 5, 10, ...). The answer:

$\langle \text{integer div-by-5} \rangle$	$::= \langle \text{sign} \rangle \langle \text{unsigned int div-by-5} \rangle$
$\langle \text{sign} \rangle$	$::= + \mid - \mid \langle \text{empty} \rangle$
$\langle \text{empty} \rangle$	$::=$
$\langle \text{unsigned int div-by-5} \rangle$	$::= 0 \mid 5 \mid \langle \text{number} \rangle \langle \text{unsigned int div-by-5} \rangle$
$\langle \text{number} \rangle$	$::= \langle \text{non-zero digit} \rangle \mid \langle \text{number} \rangle \langle \text{digit} \rangle$
$\langle \text{digit} \rangle$	$::= 0 \mid \langle \text{non-zero digit} \rangle$
$\langle \text{non-zero digit} \rangle$	$::= 1 \mid 2 \mid 3 \mid 4 \mid 5 \mid 6 \mid 7 \mid 8 \mid 9$